

From: Judith Curry [curry.judith@gmail.com]
Sent: 5/20/2025 2:29:33 PM
To: Roy Spencer [roywspencer@hotmail.com]
CC: Ross McKittrick [ross.mckittrick@gmail.com]; Steve Koonin [steven.koonin@gmail.com]; John Christy [climatemanager60@gmail.com]; Travis Fisher [travis.scott.fisher@gmail.com]
Subject: new 3.3.2
Attachments: 3.3.2 curry.docx

ne

On Tue, May 20, 2025 at 4:34 AM Roy Spencer <roywspencer@hotmail.com> wrote:
...into the carbon cycle chapter where appropriate. As Judy pointed out, there are large variations among models. I will do the ocean models next.

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

3.3.2 The carbon cycle relating emissions and concentrations

Reservoirs of carbon in the earth system include the atmosphere, land ecosystems, the ocean, sediments and the Earth's interior. Excluding rocks, by far the largest reservoir of carbon is the ocean. Exchanges of carbon between reservoirs occurs via various chemical, physical, geological and biological processes. These exchanges establish a dynamical equilibrium over time in the absence of large external perturbations to the system. Since the advent of agriculture, humans have gradually influenced the carbon cycle by modifying the vegetation in land ecosystems. Since the industrial revolution, humans have modified the carbon cycle via production and burning of fossil fuels that emit CO₂ and other carbon-based pollutants into the atmosphere.ⁱ

Removal of carbon from the atmosphere is achieved via photosynthesis into leaves, roots, stems, and woody biomass. Carbon is returned to the atmosphere through the respiration of plants and the decomposition of plant detritus or dead organic material and also via soil carbon. Disturbances such as fire, insect outbreaks and timber harvesting provide substantial local perturbations to the land-atmosphere carbon exchange. In oceanic regions of ocean upwelling with nutrient-rich warm waters, carbon is outgassed into the atmosphere. By contrast, where there is cold, sinking water, such as in the North Atlantic and Southern Ocean, carbon is removed from the atmosphere. Carbon is exchanged between land and ocean reservoirs via river transport from interior regions to the coastal ocean.ⁱⁱ

The El Niño Southern Oscillation (ENSO) is a significant driver of tropical carbon flux variability for both the ocean and terrestrial ecosystems. During the warm phase of ENSO (El Niño), the ocean takes up more carbon because of reduced upwelling and outgassing from the eastern Tropical Pacific. On land, El Niño is associated with drought and warmer temperatures, resulting in outgassing from land ecosystems. There is also substantial decadal variability in the land and ocean carbon sinks, driven by atmospheric and oceanic circulation patterns in the extratropics.ⁱⁱⁱ

The annual estimation of global CO₂ emissions and sinks is a major effort by the international carbon cycle research community that includes compilation and synthesis of measurements, statistical estimates, and model results. The ocean CO₂ sink showed rapid growth during the past decade, after little to no growth during 1991–2002. The land CO₂ sink demonstrated a continued increase during the last decade (2011–20), driven predominantly by increased atmospheric CO₂.^{iv}

While there is good qualitative understanding of the relevant processes involved in the global carbon cycle, our quantitative understanding is relatively weak.^v There is persistent large uncertainty in the estimate of CO₂ fluxes from land-use changes. There is low agreement between the varied methods with respect to the magnitude of the land CO₂ flux in the northern extra-tropics. There is also substantial discrepancy between the different methods on the strength of the ocean sink over the last decade, particularly in the Southern Ocean.^{vi} There is also concern about permafrost feedback loops, that could release more carbon into the atmosphere with warming.

Global carbon cycle models relate concentrations to emissions using estimates of various land and ocean processes that exchange CO₂ with the atmosphere. Our understanding of these processes was reviewed by Crisp et al. (2021). The natural fluxes of CO₂ in and out of the Earth's surface are an order of magnitude larger than anthropogenic emissions, but the natural CO₂ sources and sinks were assumed to be in approximate balance in pre-industrial times, as inferred from relatively constant levels of atmospheric CO₂ found in air bubbles trapped in ice cores. This is one of several reasons why increasing atmospheric CO₂ levels are believed to be dominated by anthropogenic sources.

Specifically, fossil fuel burning (and to a lesser extent land use [biomass burning] and cement production) has led to steadily increasing CO₂ levels in the atmosphere, as shown in Fig. 3.1.3. CO₂ emissions from fossil fuel burning is a small fraction of the natural carbon cycle components. About 44,000 Gigatonnes of carbon-equivalent (GtC) are stored in the atmosphere and oceans, mostly in the form of CO₂. (Ciais et al. 2013). Between 150 and 200 GtC are released naturally each year through oceanic outgassing and land-based processes like plant decay and animal respiration. A variable but roughly equivalent amount is absorbed each year by the land and oceans, with a small net loss from the atmosphere because the formation of deep ocean sediment sequesters some carbon permanently. Global fossil fuel consumption and cement manufacture released about 10.3 GtC in 2023, up from 6.2 GtC in 1990 (Ritchie et al. 2024). Of this, about half is naturally sequestered through plant growth and oceanic uptake while the remainder, about half, adds to the atmospheric stock (Ciais et al. 2013).

As noted, each year natural processes remove a net amount of CO₂ equivalent to about 50% of global emissions. Thus, anthropogenic CO₂ emissions average twice what is required to explain the atmospheric rise in CO₂ concentration. The historical near-constancy of that 50% fraction also means that the more CO₂ humanity has produced, the faster Nature removed it from the atmosphere. That 50% fraction changes somewhat from year to year due to natural carbon cycle imbalances from El Nino, La Nina, and varying weather patterns. There was also a substantial additional reduction in atmospheric CO₂ after the 1991 eruption of Mt. Pinatubo, a curious result that has yet to be explained (Angert et al., 2004).

The main processes that remove excess CO₂ from the atmosphere are increased growth of land vegetation (especially at high latitudes), some increase in the sequestering of carbon in soils, and uptake of CO₂ by the ocean due to the increasing partial pressure of atmospheric CO₂ over that of CO₂ dissolved in the ocean. All twenty land carbon cycle models tracked by the Global Carbon Project (Friedlingstein et al., 2024) show land processes have been removing excess CO₂ at an increasing rate since 1959. This is consistent with a “global greening” phenomenon (Chapter 2.1) observed by satellites since monitoring of global greenness began in the 1982.

While land vegetation has been responding positively to more atmospheric CO₂, uptake of extra CO₂ by ocean biological processes remains too uncertain to be measured reliably. Our current understanding of these and many more carbon cycle processes was reviewed by Crisp *et al.* (2021).

CO₂ uptake by land processes

The uptake of extra CO₂ from the atmosphere by land surface processes (as also inferred from global greening) has been modeled with 20 different dynamic global vegetation models, the outputs of which are updated every year by the Global Carbon Project (Friedlingstein, 2024). As seen in Fig. 3.3.3, all of those models agree that vegetation and soils have been sequestering carbon from the atmosphere. But we also see that the long-term trends over 1959 to 2023 (65 years) vary widely between models, by nearly a factor of 7. This demonstrates that there remains considerable uncertainty regarding how fast CO₂ from the atmosphere is being removed by land processes, which in turn translates into uncertainty regarding future atmospheric CO₂ concentrations. Those uncertain future CO₂ concentrations then imply a source of uncertainty in climate model simulations of future climate change.




Figure 3.3.3 Decadal trends in the yearly CO₂ uptake by land processes during 1959-2023 simulated by 20 different dynamic global vegetation models periodically reported by the Global Carbon Project (Friedlingstein, 2024).

CO₂ uptake by ocean processes

The uptake of extra CO₂ from the atmosphere by ocean processes has been modeled with 10 different dynamic global vegetation models, the outputs of which are updated every year by the Global Carbon Project (Friedlingstein, 2024). Like the results from the land models, all of the ocean models agree that the global oceans have been sequestering carbon from the atmosphere at an increasing rate during 1959-2023 (Fig. 3.3.4). Unlike the land models, however, the ocean models show much better agreement with each other, with the model producing the fastest increasing CO₂ uptake being only 65% faster than the model with the most slowly increasing CO₂ uptake. Note that the average trend in CO₂ uptake across all land models in Fig. 3.3.3 is 25% larger than the average trend in ocean uptake. This suggests land

processes are increasing in their ability to remove CO₂ faster than ocean processes are increasing their CO₂ sequestration.




Figure 3.3.5 Decadal trends in the yearly CO₂ uptake by ocean processes during 1959-2023 simulated by 10 different ocean biogeochemistry models periodically reported by the Global Carbon Project (Friedlingstein, 2024).

New research continues to provide surprises. Phytoplankton, which live on the warm, light-filled ocean surface, use atmospheric CO₂ for food. Phytoplankton sequester carbon and once they die and sink, carry it to the ocean depths. A recent study found evidence that phytoplankton may become more efficient at sequestering carbon as the ocean warms. Another recent article reported the discovery of a widely circulated microbe variety that demonstrates the potential to sequester carbon.^{vii}

ⁱ “Chapter 1: Overview of the Global Carbon Cycle,” Second State of the Carbon Cycle Report (U.S. Global Change Research Program), accessed July 25, 2022, <https://carbon2018.globalchange.gov/chapter/1/>.

ⁱⁱ Ibid.

ⁱⁱⁱ Peter Landschützer et al., “Decadal Variations and Trends of the Global Ocean Carbon Sink,” *Global Biogeochemical Cycles* 30, no. 10 (October 5, 2016): 1396-1417, <https://doi.org/10.1002/2015gb005359>.

^{iv} Pierre Friedlingstein et al., “Global Carbon Budget 2021,” *Earth System Science Data* 14, no. 4 (April 26, 2022): 1917-2005, <https://doi.org/10.5194/essd-14-1917-2022>.

^v Ibid.

^{vi} Ibid.

^{vii} Nancy Averett, “The Ocean Is Still Sucking up Carbon-Maybe More than We Think,” *Eos*, May 3, 2022, <https://eos.org/articles/the-ocean-is-still-sucking-up-carbon-maybe-more-than-we-think>.

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Tuesday, May 20, 2025 6:33 AM
To: Ross McKittrick; Judith Curry
Cc: Steve Koonin; John Christy; Travis Fisher
Subject: New CO2 emissions scenarios graph
Attachments: SSP-emissions-scenarios-vs-historical-and-EIA-projections.jpg

The new citation for the observed and EIA-projected emissions is:

Friedlingstein, P., and 95 co-authors (2024): Global Carbon Budget 2024, Earth Syst. Sci. Data, <https://essd.copernicus.org/preprint>

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Monday, May 19, 2025 9:24 PM
To: Judith Curry <curry.judith@gmail.com>
Cc: Steve Koonin <steven.koonin@gmail.com>; John Christy <climateman60@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Travis Fisher <travis.scott.fisher@gmail.com>
Subject: Re: Albedo text for ch 5


Thanks Judy.

On Mon, May 19, 2025 at 7:30 PM Judith Curry <curry.judith@gmail.com> wrote:
Attached

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>



From: Ross McKittrick [ross.mckittrick@gmail.com]
Sent: 5/17/2025 5:36:13 PM
To: Travis Fisher [travis.scott.fisher@gmail.com]; Roy W. Spencer [roywspencer@hotmail.com]; John Christy [climatemanager60@gmail.com]; Steven Koonin [steven.koonin@gmail.com]; Judith Curry [curry.judith@gmail.com]
Subject: New econ ch with tipping points
Attachments: 12 Climate, Economy and SCC.May17.docx

There's a new version of the Econ-SCC chapter with a section added on tipping points that Judy and I assembled. It's dated May 17 and is attached and is in the 01.Drafts folder.

12 Climate change, the economy and the Social Cost of Carbon

12.1 Climate change and economic growth

12.1.1 Overview: climate is a relatively minor issue for a modern economy

It has long been noted that economies tend to do poorly in very cold and very hot regions, with the optimum somewhere in between (Nordhaus 2006). Temperature-sensitive economic activity migrates, whenever possible, to where it is best suited, and society always adapts to the local climate. Based in part on these observations, in 1992 Thomas Schelling, then President of the American Economic Association, argued that any effects of climate change on US economic activity would be small relative to the many other changes that would happen.

... Manufacturing rarely depends on climate, and where temperature and humidity used to make a difference, air conditioning has intervened. When Toyota chooses among Ohio, Alabama, and Southern California for locating an automobile assembly, geographical considerations are important, but not because of climate... Finance is little affected by climate; similarly for health care, or education, or broadcasting. Transportation can be affected, but improvements in all-weather landing and take-off in the last 30 years are greater than any differences that climate makes. If the average effect is a warming, iced waterways and snow removal may decline in importance. Construction is affected, mainly by cold, and if the average effect is in the direction of warming, construction may benefit slightly.

It is really agriculture that is affected. But even if agricultural productivity declined by a third over the next half century, the per capita GNP we might have achieved by 2050 we would achieve only in 2051. ...

I conclude that in the United States, and probably Japan, Western Europe, and other developed countries, the impact on economic output will be negligible and unlikely to be noticed.

(Schelling 1992, emphasis added)

Thirty years later virtually the identical point was made by the IPCC itself in the Fifth Assessment Report:

For most economic sectors, the impact of climate change will be small relative to the impacts of other drivers. Changes in population, age, income, technology, relative prices, lifestyle, regulation, governance, and many other aspects of socioeconomic development will have an impact on the supply and demand of economic goods and services that is large relative to the impact of climate change.

(Arent et al. 2014, emphasis added)

That statement is validated by experience. Since 1900, the average global surface temperature anomaly warmed 1.3°C, about as much as the IPCC predicts will occur in the next century under a moderate emissions scenario. But even as the globe warmed and the population quintupled, humanity prospered as never before. For example, global average lifespan went from thirty-two years to seventy-two years, economic activity per capita grew by a factor of seven, and the death rate from extreme weather events plummeted by a factor of fifty!

While extreme weather events are costly, in all modern economies they are becoming steadily less and less important (Formetta and Feyen 2019). Since 1990 weather-related disaster losses have declined as a proportion of global GDP (Pielke Jr 2018, 2020) as have mortality risks (Formetta and Feyen 2019). While economic weather-related insurance payouts are rising, this is fully explained by the growth in the size of the economy and the value of the insured asset stock. Past increases in episodes extreme weather have not had a significant effect (positive or negative) on the market value of insurance firms (Hu and McKittrick 2015). Nor have past extreme weather events had a significant effect on US banks' performance (Blickle et al. 2021).

For these reasons, economists have long been reluctant to endorse attempts to "stop" climate change or even aggressively reduce GHG emissions because the costs would not be worth it. As one critic of the economics of climate policy put it:

Mainstream climate economics takes global warming seriously, but perplexingly concludes that the optimal economic policy is to almost do nothing about it... The contrast is striking. While climate science is sending out loud-and-clear messages that fossil-fuel disinvestment must start now, letting go of coal and oil and diverting resources into renewable energy technology systems, to keep warming below the 2°C limit (IPCC 2014), mainstream climate economics claims that overly ambitious climate targets will unnecessarily hurt the economy and immediate de-carbonization is too expensive. Most climate economists thus recommend humanity to just wait-and-see.
(Storm 2017)

The mainstream economics position on the climate policy question is best represented by the findings over the past three decades from Integrated Assessment Models (IAMs) of climate change policy, for which Yale economist William Nordhaus was awarded the 2018 Nobel Memorial Prize in economics. IAMs combine economic, climate, and social data into a unified framework for simulating climate damages and evaluating the optimal policy response (Resources for the Future 2025). Nordhaus' work has generally supported modest global climate policy, with aggressive measures largely deferred to later in the century. Nordhaus developed the so-called Dynamic Integrated Model of the Climate and Economy model, or DICE, in the early 1990s (Nordhaus 1993) to study the interaction between climate change, climate policy, and global economic growth over long times. DICE assumes global emission control can be coordinated at no cost and asks what the optimal policy target should be. The climate component of the DICE model is based on simplified climatological modeling. The version of DICE at the time of Nordhaus' Nobel Prize award was described in Nordhaus (2018); it had a climate sensitivity parameter of 3.1°C for CO₂ doubling, consistent with the best estimate in contemporary IPCC reports (3.0°C).

The baseline (no policy) scenario incurs only \$0.4T in global abatement costs and leads to \$134.2T (trillion dollars) in global climate damages for a total cost of \$134.6T. The climate model component of DICE projects 4.1°C warming by 2100 relative to preindustrial temperatures. Note that this is a higher estimate of warming than many IPCC climate models. (AR6 offers 2.7°C in 2100 under a moderate emissions scenario.)

The Optimal Policy scenario barely deviates from the business-as-usual path. It aims for +3.5°C warming, in other words we scale back only the lowest-value fossil fuel use and otherwise just live with almost all the warming. This suggests that most CO₂ emissions are less harmful than the policies that would be necessary to abate them. Trying to prevent warming causes costs quickly to outstrip the benefits. Pursuing a goal of capping warming at 2.5°C creates total costs of \$177.8T,

which is \$43.2T worse than doing nothing at all. Nordhaus didn't evaluate trying to reach a Paris-type target of 1.5°C warming but it would be even worse.

A subsequent edition of DICE includes higher assumed damages from warming which, not surprisingly leads to more aggressive policy recommendations, as explained in the section below on the Social Cost of Carbon.

12.1.2 Empirical analysis of climate change and economic growth

Many other studies have used econometric methods applied to historical data, instead of IAMs, to study the potential impact of climate change on economic growth. Dell et al. (2012, herein DJ012) was an influential study that used a multi-country panel of national-level data spanning 1950 to 2005, in which they matched climate and economic data by averaging temperature from the local grid cell level up to the national level using population weights. They found that warming yields an insignificant positive effect on income growth in rich countries but a significant negative effect in poor countries. Moore and Diaz (2015) modified the DICE model to take that finding into account and concluded that it implied a dramatically higher Social Cost of Carbon due to the compounding effects of income loss over time.

A large subsequent literature has debated the robustness of the DJ012 findings. Burke et al. (2015) analyzed a global panel with temperatures averaged up to the national level and found a negative effect on growth from warming in rich and poor countries alike when the national average temperature is above 13°C. Zhao et al. (2018) used the G-Econ data set of Nordhaus (2006), which breaks economic activity down to the grid cell level, replicated DJ012-type results on the same subset of countries as used in DJ012 but then showed that on the full global sample warming increases growth in rich countries and poor countries alike, though the positive effect in the latter group is confined to where local temperatures are below about 16°C. Greßer et al. (2021) developed a regional economic data set for 1,542 sub-national regions around the world between 2005 and 2015 and found temperature had no effect on growth. Yang et al. (2023) updated the DJ012 data set and applied an estimator robust to mixed sample frequencies, finding that while temperature shocks exerted a temporary effect on income levels, they did not have a significant lasting effect on growth rates.

Newell et al. (2021) noted that there is no underlying theory to guide econometric model specification in this literature. Taking into account the arbitrary choices of which explanatory variables to include they identified over 800 possible model specifications. Using the Burke et al. (2015) data they used an estimation method that accounts for model uncertainty and found that the model form preferred by Burke et al. (2015), which implied negative effects of warming on growth even in rich countries, is explicitly excluded by the optimal model selection algorithm. Overall they could not detect a temperature effect on GDP or GDP growth, and they estimated the 95% confidence interval for the impact on global growth as of 2100 even under the exaggerated RCP8.5 warming scenario spans -86 percent to +388 percent. In other words the net effect is likely positive but too uncertain to distinguish from zero.

Barker (2023) criticized the DJ012 assumption that countries can be grouped into fixed “poor” and “rich” categories based on their incomes many decades ago. He noted that many countries were once poor but became rich over time, and if this is taken into account the original temperature effects reported by Dell et al. became small and insignificant.

Berg et al. (2023) argued that countries shouldn't be grouped into large Rich/Poor categories because they are too heterogeneous. They instead estimated country-specific temperature response

coefficients then grouped countries with similar response coefficients into small panels. They separately estimated responses to global and idiosyncratic local temperature shocks to better identify the climate signal in weather data. Overall they found countries experiencing negative effects of warming on growth outnumbered those experiencing positive effects, but only temporarily: eventually the effects reverse such that about twice as many countries experience a net positive growth effect. They also found global (as opposed to local) temperature changes are much more likely to benefit growth in poor countries than rich ones. In a simulation to 2100 even using the extreme RCP8.5 scenario they computed the average global GDP loss would be only 1.9% compared to a scenario with no warming. That is, instead of the economy growing 400% it would grow 398%. The implication of Nordhaus' earlier analysis is that trying to prevent the warming would lead to far less than 398% growth.

To summarize, economists have long considered climate a relatively unimportant factor in economic growth, a view echoed by the IPCC itself in the Fifth Assessment Report. Mainstream climate economics has recognized that CO₂-induced warming may have some negative economic effects but they are too small to justify aggressive abatement policy, and trying to “stop” or cap global warming even at levels well above the Paris target would be worse than doing nothing. An influential study in 2012 suggested that global warming would harm growth in poor countries but the finding has subsequently been found not to be robust. Studies that take full account of modeling uncertainties either find no evidence of a negative effect on global growth from CO₂ emissions or find poor countries as likely to benefit from it as rich countries.

The expectation that significant global warming would have a small impact on the U.S. economy was acknowledged quietly by the Biden Administration, even as the President was proclaiming a climate emergency. The following figure, from a 2023 OMB-CEA [report](#), shows the expected decrement in U.S. GDP as a function of temperature rise. The colored lines show the results of a dozen peer-reviewed published studies while the solid black line is their average. The figure could be summarized as “a few percent impact for a few degrees of warming”. Given that the economy’s annual growth rate is expected to be 1-2 percent, the impact of a warming globe is indeed negligible.




Figure 12.1: Decline in US GDP per degree of warming. Source: XX

12.2 Models of the Social Cost of Carbon

The Social Cost of Carbon (SCC) is a tool for quantifying the economic impact of carbon dioxide emissions, helping policymakers weigh the costs and benefits of climate policies. It estimates the damage caused by emitting one additional ton of CO₂, expressed in dollars. More formally, the SCC is the discounted present value of the current and future marginal loss of economic welfare due to an additional ton of CO₂ entering the atmosphere.

Sidebar: finding the optimal emissions price

On its own the SCC is not the appropriate price to attach to CO₂ emissions for the purpose of optimal emissions control but it can be converted into an optimal tax rate by scaling it by a parameter called the Marginal Cost of Public Funds (MCPF). The theory behind this was first derived by Sandmo (1975) and has been reiterated many times since then by other economists working on the topic (e.g. Parry et al. 1999). The MCPF is the cost in terms of lost private economic welfare of raising public revenue by one dollar and always exceeds unity. If a jurisdiction does not use a carbon tax, this formula still indicates the upper bound on the amount emitters should spend on abatement options at the margin along the optimal policy path. Regulations should be imposed only insofar as they do not impose marginal abatement costs in excess of the optimal tax rate.

12.2.1 Estimating the SCC

Although the literature refers to “estimates” of the SCC, it is not estimated in the way other economic statistics are estimated. For instance, data on market transactions including prices and quantities can be used to estimate the current inflation rate or the growth rate of per capita real Gross Domestic Product, and there are well-understood uncertainties associated with these quantities. But there are no market data available to measure many if not most of the marginal damages or benefits believed to be associated with CO₂ emissions, so these need to be imputed using economic models.

For example, an influential component of some SCC calculations is the perceived social cost associated with changed risk of future mortality due to extreme weather. There is no market in which people can directly attach a price to that risk. At best economists can try to infer such values by looking at transactions in related markets such as real estate or insurance, but isolating the component of price changes attributable to atmospheric CO₂ levels is very difficult.

Economists use IAMs to compute the SCC. Two of the best known are the Climate Framework for Uncertainty, Negotiation and Distribution (“FUND”, Tol 1997) and Nordaus’ DICE. IAMs embed a “damage function” or set of functions relating ambient temperature to local economic conditions. The assumptions embedded in the damage function will largely determine the resulting SCC. IAMs also assume a long-term discount rate or, as in DICE, compute the optimal internal discount rate as part of the solution.

One approach to developing a damage function is to begin with estimates of the costs (or benefits) of warming in specific sectors in countries around the world and aggregate up to a global amount. This was the approach used in FUND. An alternative approach is to develop a simple equation that penalizes global income according to a simple quadratic function of the average global temperature. This approach was used in DICE. In the case of the FUND model many hundreds of parameters had to be selected, whereas in DICE only three were needed and were originally chosen to assign a predetermined penalty (1.2%) to global output as of 3°C warming, with a quadratic term implying

that damages grow with the square of the global average temperature anomaly. Barrage and Nordhaus (2024) recently changed the parameters to increase the penalty at 3°C to 1.6%, and added a discrete additional 1.0% GDP penalty at 3°C warming to account for “tipping points” (discrete large-scale environmental changes triggered by crossing a warming threshold) and a “judgmental adjustment” of 0.5% for excluded impacts at 3°C warming. Not surprisingly the newer version of DICE generates much higher SCC estimates than before.

The concepts of estimation and uncertainty do not readily apply to SCC calculations. By comparison, the question of whether precipitation has trended up or down along the South Carolina coast for the past hundred years can be addressed by collecting rainfall data, estimating a trend parameter and calculating statistical confidence intervals using methods with well-understood mathematical properties. In this way uncertainty can, in principle, be resolved by collecting more data and applying more accurate methods. But no amount of data collection can change the fact that many components of the SCC are unknown and rely on judgment and opinion based on knowledge of the underlying literature on the physical effects of climate change.

SCC calculations are thus best thought of as “if-then” statements: *if* the following assumptions hold, *then* the SCC is \$X per tonne. The list of ‘if’ statements includes the premise that the world’s climate and economy work according to the representation in the IAM. One way this may fail to hold is in the timing of warming. Every IAM assumes a value of the Equilibrium Climate Sensitivity (ECS), which controls the amount of warming that results from CO₂ emissions and can be freely varied for the purpose of generating a distribution of SCC values associated with uncertainties over ECS. But as Roe and Baumann (2013) pointed out, time-to-equilibrium increases with the square of ECS, so an upward adjustment of the ECS parameter without an appropriate slowing down of the adjustment process can yield distorted present value damage estimates. In particular, the upper tail of warming associated with some commonly-used ECS distributions is physically impossible for even a thousand years into the future (Roe and Baumann 2013) yet in an IAM would be realized within a couple of centuries. Failure to align ECS with time-to-equilibrium will lead to an overestimate of the SCC value.

12.2.2 Variations in the SCC

Every level of the IAM calculation includes assumptions, some more influential than others. Key assumptions include the following.

- The discount rate: Climate damages accrue over a long time horizon, and costs a century or two in the future need to be discounted to the present. The higher the discount rate the smaller the present value of future damages and vice versa. The discount rate represents the opportunity cost of spending money today rather than investing it and then having more to spend tomorrow. Some economists have argued for the use of very low discount rates in SCC calculations, resulting in policy recommendations that favor relatively large immediate investments in CO₂ emission reductions. The downside is that other investments could potentially earn a larger rate of return for society.
- Equilibrium Climate Sensitivity: IAMs have customarily employed a value of 3.0°C or 3.1°C following the IPCC’s guidance. Empirically-estimated ECS values tend to be lower than this (see Chapter 9). Dayaratna et al. (2017, 2020, 2023) have shown that use of lower empirically-derived ECS values dramatically lower the resulting SCC estimate, even when low discount rates are used.

- Damage function coefficients: IAMs assume CO₂ and warming cause net harms that increase exponentially with temperature. More recently, IAMs have also incorporated effects from assumed potential climate tipping points. The FUND model took limited but explicit account of CO₂ fertilization effects in agriculture. Since the coefficients were selected prior the publication of the current evidence of global greening and the magnitude of benefits to crops from elevated CO₂ (see Chapters 2 and 10) the growth effects are likely understated. The DICE model (and others) did not explicitly include any CO₂ fertilization benefit, except to the extent it was taken into account in the literature consulted when picking the damage function coefficients. The damage function in FUND contains a region in which low warming yields net benefits in many regions, a finding which is supported by econometric models of warming and growth (Berg et al. 2023) and econometric simulations of agricultural changes (McKittrick 2025). The DICE damage function, by construction, rules out net benefits at any warming level.
- Emission scenarios: IAMs generate SCC estimates that increase as the pre-existing concentration of CO₂ increases. Consequently the value of damages later in the century will be higher, depending on the assumed baseline emissions over the coming decades.
- Abatement costs: IAMs represent the cost to the economy of reducing CO₂ emissions. If CO₂ emission reductions are assumed to be inexpensive, then the model will conclude that the optimal policy should aim for deeper emission cuts and vice versa.

It is informative to ascertain whether SCC results are invariant to changes in some assumptions. But when different assumptions result in higher or lower SCC values, the change in the SCC value does not provide *prima facie* evidence about the validity of the assumptions. For example, in 2023 the US Environmental Protection Agency raised its preferred SCC value about 5-fold over the estimates it had issued ten years earlier. This is not because new data had been collected or better mathematical methods had been invented, but because new assumptions had been used, and the validity of those assumptions was a separate question. One was that global agricultural damages were far higher than previously assumed, based on an analysis in Moore et al. (2017). But as discussed in Chapter 10, McKittrick (2025) showed that Moore et al. (2017) had used a database in which half the CO₂ change observations were missing. When as many of those observations as possible were recovered from underlying sources and the analysis was rerun, the projected crop yield losses disappeared and turned into gains at all warming levels. Hence the portion of the EPA's SCC revision attributable to agricultural yield losses was unwarranted.

12.2.3 Evidence for low SCC

Chapter 2 reviewed evidence on climate change and greening, and Chapter 10 looked at climate change and US agriculture. Evidence shows that CO₂ fertilization has a stronger beneficial effect on agriculture than was known at the time that IAMs like DICE and FUND were parameterized. In particular Haverd et al. (2020) found the observed CO₂ fertilization rate has been almost double what had been predicted by crop models. Dayaratna et al. (2023) used the updated empirical ECS distribution estimate of Lewis (2022), which assimilates modern instrumental and paleoclimatic temperature records, and allowed for a 30 percent gain in the CO₂ fertilization benefit in the FUND model, and found that, even at a low discount rate of 2.5 percent, the median SCC as of 2050 is only \$18.67, with a 24 percent probability of the true value being negative. At a five percent discount rate the median SCC value is negative until the mid-2040s and at 2050 was only \$0.37 with a 49 percent probability of being negative. Thus, under reasonable assumptions a mainstream IAM using

updated scientific inputs yields evidence consistent with the SCC not being significantly greater than zero.

It should also be noted that the SCC is focused on the social costs of CO₂ emissions from fossil fuel use. It does not measure the private marginal benefits to consumers and society from the availability of fossil fuels. Public willingness to pay for fuels of all types indicates the value to society of reliable, abundant fossil energy. Tol (2017) estimates that the private benefit of carbon is large relative to the social cost. This can be illustrated by noting that the price of a gallon of gas indicates the marginal value to the consumer of the fuel. Suppose we assume a relatively high Social Cost of Carbon of, say, \$75 per tonne. Deflated by a MCPF value of 1.5 that would result in a carbon tax of \$50 per tonne, which equates to about 44 cents per gallon of gas (Lavelle 2019). A pre-tax price of \$3.00 per gallon would imply the marginal social benefit of the fuel is nearly seven times the marginal social cost.

12.2.4 Tipping points

SCC calculations typically consider gradual impacts of a warming climate, such as slowly-melting glaciers and increasing average temperature. A driver of potentially high values of the Social Cost of Carbon (SCC) is the introduction into models of discrete catastrophic outcomes associated with abrupt changes (Dietz et al. 2021). They are often referred to as “tipping points.”

The term “tipping point” mingles two different physical concepts that pose different research challenges. Many physical systems are inherently stable unless acted upon with sufficient external energy. For example an ice cap might remain intact over a wide range of temperatures but once the temperature crosses the 0°C threshold it melts. Such discontinuities are ubiquitous in nature and require an external force. Whether the force needs to be large relative to the size of the system depends on the underlying stability of the system.

A different type of tipping point is called a *bifurcation* and arises from the study of the internal dynamics of nonlinear systems (Crawford 1991). Many systems have been observed to have more than one equilibrium point and can move between them with minimal or no external influence. For example a weather system might have two different equilibrium states: one calm and one with a tornado. A transition from one to another can happen either with no external force or with a minuscule change, such as a flap of the proverbial butterfly’s wings (Shen et al. 2014). The term “tipping point” is sometimes used to mean a bifurcation of this type and implies instability inherent in the system itself, which is not necessarily dependent on outside forces. It depends, instead, on parameters of the system taking values that support the emergence of bifurcations (Crawford 1991).

The two different concepts imply different research questions. Regarding the first we are interested in whether components of the climate system are susceptible to abrupt discontinuities in response to sufficiently-large anthropogenic forcing. Regarding the second we are interested in whether the Earth’s climate system has inherent bifurcations that imply the possibility of abrupt transitions with or without external forcing.

Models have been developed that imply the latter is a possibility. Kypke et al. (2020) presented a simple climate model in which the GHG concentration is one of the parameters that controls the emergence of bifurcations of the Arctic climate to one with both cold and warm equilibria. If sufficiently high GHG forcing combined with a sufficiently high rate of ocean heat transport are

imposed a bifurcation becomes possible. More generally GCMs have been observed to contain bifurcations and multiple equilibria (Brunetti and Ragon 2023).

The possible existence of bifurcations in the Earth's climate system implies abrupt transitions are possible, not just in response to large forcing but to small perturbations as well. This places tipping points into the category of low-likelihood and potentially catastrophic events, such as large meteor strikes. A key question to ask is whether those kinds of tipping points can be predicted. Current research has not resolved that question (Dakos et al 2024) and indeed may not be able to since one implication of the "butterfly effect" is the existence of predictability boundaries of nonlinear systems (Palmer et al. 2014). It is therefore not obvious how to incorporate such possibilities into SCC calculations. Small variations in assumptions will lead to arbitrarily large variations in the resulting SCC with no grounds for choosing among them. If such tipping points are possible the most appropriate stance for economic policy is to maximize resilience to any form of external catastrophe since it is unlikely we could predict or prevent it from happening.

The IPCC AR6 (Chapter 1) focuses mainly on the first type of tipping point, namely an abrupt change in response to external forcing. This is also the meaning associated with popular usage of the "tipping point" concept in discussions of climate change (see <https://report-2023.global-tipping-points.org/what-is-a-tipping-point/> for example). As summarized by the IPCC AR6, there is evidence of abrupt change in the paleoclimate record, and some of these events have been interpreted as tipping points. Some projections with Earth System Models for example have produced tipping points such as Amazon forest dieback in response to specified values of CO₂ concentration or temperature increases.

The alarm surrounding climate tipping points is reflected by *The Global Tipping Points Report* that was launched at the COP28 on 6 December 2023 (Lenton et al., 2023). It identifies more than 25 parts of the climate system said to constitute tipping points. What gets classified as a climate "tipping point" is a moving target. The most common examples in the literature and assessment reports include: Greenland ice sheet disintegration, West Antarctic ice sheet disintegration, summertime disappearance of Arctic sea ice, Amazon rainforest dieback, coral reef dieoff, thawing of permafrost and methane hydrates, Atlantic Meridional Overturning Circulation collapse, boreal forest shift, West African monsoon shift, and Indian Monsoon shift.

All such tipping points require a certain amount of system instability in order to exhibit an abrupt transition in response to warming. For this reason there seems to be very little to discriminate between a tipping event and natural climate variability. Natural climate variability has in the past produced shifts in the West African and Indian monsoons, Amazon forest and coral reef dieback, and disintegration of the Greenland and West Antarctic ice sheets. These impacts can be reversed on the decadal timescales associated with natural climate variability and ecosystem responses.

Some abrupt changes are potentially more consequential, including collapse of the West Antarctic ice sheet and collapse of the Atlantic Meridional Overturning Circulation (AMOC). The IPCC AR6 WG1 Summary for Policy Makers states:

"The Atlantic Meridional Overturning Circulation is very likely to weaken over the 21st century for all emission scenarios. While there is high confidence in the 21st century decline, there is only low confidence in the magnitude of the trend. There is medium confidence that there will not be an abrupt collapse before 2100." (C.3.4)

"There is limited evidence for low-likelihood, high-impact outcomes (resulting from ice-sheet instability processes characterized by deep uncertainty and in some cases involving tipping points) that would strongly increase ice loss from the Antarctic Ice Sheet for centuries under high GHG emissions scenarios (e.g., SSP5-8.5)." (B.5.2)

For the purpose of SCC calculations the research question implied by this type of tipping point is whether such events have been observed in the past in climate conditions similar to what we currently experience or will in the near future. The IPCC AR6 finds little evidence for impending collapse of the Atlantic Meridional Overturning Circulation or the West Antarctic ice sheet. It finds there is no tipping point associated with Arctic Sea ice (AR6 Technical Summary p. 76)

Dietz et al. incorporated several potential tipping points (abrupt changes) into an SCC model and found they added about 25 percent to the estimate, mainly associated with thawing permafrost and release of methane hydrates. However the IPCC considers this scenario *very unlikely* (AR6 Technical Summary p. 107).

In summary, there may be unknown bifurcation tipping points that are associated with natural climate processes, but this possibility does not translate into specific guidance on the SCC. There are potential abrupt change points in the climate system in response to warming, although the IPCC assigns low probabilities to most, including the largest ones. When these have been taken into account the result is only a modest increase in the SCC value in the 21st century.

12.3 Summary

In summary, SCC estimates are highly uncertain due to unknowns in future economic growth, socioeconomic pathways, discount rates, climate damages, and system responses. The SCC is not intrinsically informative as to the economic or societal impacts of climate change. It provides an index connecting large networks of assumptions about the climate and the economy to a dollar value. Some assumptions yield a high SCC and others yield a low or negative SCC. The evidence for or against the underlying assumptions needs to be established independently; the resulting SCC adds no additional information about the validity of those assumptions.

It is increasingly being argued that the SCC is too variable to be useful for policymakers. Cambridge Econometrics (Thoung 2017) stated it's "time to kill it" due to uncertainties. The UK and EU no longer use SCC for policy appraisal, opting for "target-consistent" carbon pricing (UK Department for Energy Security and Net Zero 2022, Dunne 2017). However the weaknesses of the SCC concept don't mean that other regulatory measures are inherently better or more efficient. Many climate regulations (such as electric vehicle mandates, renewable energy mandates, energy efficiency regulations and bans on certain types of home appliances) end up costing far more per tonne of abatement than a simple price on carbon emissions.

Barrage, Lint and William Nordhaus (2024) Policies, projections, and the social cost of carbon: Results from the DICE-2023 model. *Proceedings of the National Academy of Sciences* 121 (13) e2312030121 <https://doi.org/10.1073/pnas.2312030121>

Berg, Kimberly, Chadwick C Curtis and Nelson Mark. (2021) GDP and temperature: Evidence on cross-country response heterogeneity. NBER Working Paper 31327 June 2023 <http://www.nber.org/papers/w31327>

- Chris Thoung (2017) "Is It Time to Kill Off the Social Cost of Carbon?" Cambridge Econometrics Blog, September 14, 2017, <https://www.camecon.com/blog/time-kill-social-cost-carbon/>.
- Daisy Dunne (2017), "Q&A: The Social Cost of Carbon," Carbon Brief, February 14, 2017, <https://www.carbonbrief.org/qa-social-cost-carbon/>
- McKittrick, Ross R. (2025) Extended Crop Yield Meta-analysis Data do not Support Upward SCC Revision. *Scientific Reports* 15 article 5575 <https://www.nature.com/articles/s41598-025-90254-2>
- Moore FC, Baldos U, Hertel T, Diaz D (2017) New science of climate change impacts on agriculture implies higher social cost of carbon. *Nature Communications* 8:1607. <https://doi.org/10.1038/s41467-017-01792-x>
- Nordhaus, W. (1993) Optimal Greenhouse-Gas Reductions and Tax Policy in the "DICE" Model *American Economic Review* Vol. 83, No. 2, (Papers and Proceedings) pp.313-317.
- Parry, Ian, Roberton C. Williams III and Lawrence H. Goulder (1999). "When Can Carbon Abatement Policies Increase Welfare? The Fundamental Role of Distorted Factor Markets." *Journal of Environmental Economics and Management* 37: 52—84.
- Resources for the Future (2025) "Social Cost of Carbon Explorer," Resources for the Future website, accessed April 22, 2025, <https://www.rff.org/publications/data-tools/scc-explorer/>.
- Roe, Gerard H. and Yoram Bauman (2013) Climate Sensitivity: Should the climate tail wag the policy dog? *Climatic Change* (2013) 117:647–662 DOI 10.1007/s10584-012-0582-6
- Sandmo, Agnar (1975) "Optimal Taxation in the Presence of Externalities." *Swedish Journal of Economics* 77 (1), 86–98.
- Tol, R.S.J. (1997) On the optimal control of carbon dioxide emissions: an application of FUND *Environmental Modeling and Assessment* 2: 151—163.
- Tol, Richard S.J. (2017). "The Private Benefit of Carbon and its Social Cost," Working Paper Series 0717, Department of Economics, University of Sussex Business School. <https://ideas.repec.org/p/sus/susewp/0717.html>
- UK Department for Energy Security and Net Zero (2022). "Carbon Valuation." GOV.UK, last updated April 4, 2022. <https://www.gov.uk/government/collections/carbon-valuation--2>.
- US Environmental Protection Agency (2023) Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, "Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review", EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances, November 2023, https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf accessed February 28, 2024.
- Storm, S (2017) "How the Invisible Hand is Supposed to Adjust the Natural Thermostat: A Guide for the Perplexed." *Science and Engineering Ethics* vol 23: 1307—1331.
- Pielke Jr., Roger (2020) Economic 'normalisation' of disaster losses 1998–2020: a literature review and assessment," *Environmental Hazards*, 2020, pp. 1-19
- Pielke Jr., Roger (2018): Tracking progress on the economic costs of disasters under the indicators of the sustainable development goals, *Environmental Hazards*, DOI: 10.1080/17477891.2018.1540343
- Formetta, Giuseppe and Luc Feyen (2019) Empirical evidence of declining global vulnerability to climate-related hazards. *Global Environmental Change* 57 101920 <https://doi.org/10.1016/j.gloenvcha.2019.05.004>
- Hu, Bin and Ross McKittrick (2015) Climatic Variations and the Market Value of Insurance Firms. *Journal of Insurance Issues* Vol. 39, No. 1, pp. 92-111 <https://www.jstor.org/stable/43741103>
- Blickle, Kristian; Sarah N. Hamerling, and Donald P. Morgan (2021) : How bad are weather disasters for banks?, Staff Reports, No. 990, Federal Reserve Bank of New York, New York, NY <https://www.econstor.eu/bitstream/10419/247913/1/sr990.pdf>

Lavelle, Marianne (2019) Carbon Tax Plans: How They Compare and Why Oil Giants Support One of Them. *Inside Climate News* March 7, 2019 <https://insideclimatenews.org/news/07032019/carbon-tax-proposals-compare-baker-shultz-exxon-conocophillips-ccl-congress/>

Dietz, S., J. Rising, T. Stoerk, G. Wagner (2021) Economic impact of tipping points in the climate system. *PNAS* 2021 Vol. 118 No. 34 e2103081118

Shen, Bo-Wen, Roger A Pielke Sr and Xubin Zeng (2023) The 50th Anniversary of the Metaphorical Butterfly Effect since Lorenz (1972): Multistability, Multiscale Predictability, and Sensitivity in Numerical Models. *Atmosphere* 2023 14(8) <https://doi.org/10.3390/atmos14081279>

Crawford, John D (1991) Introduction to Bifurcation Theory. *Reviews of Modern Physics* 63(4) October 1991 991—1037.

Kypke, K. L., Langford, W. F., and Willms, A. R.: Anthropocene climate bifurcation, *Nonlin. Processes Geophys.*, 27, 391–409, <https://doi.org/10.5194/npg-27-391-2020>, 2020.

Brunetti, Maura and Charline Ragon (2023) Attractors and bifurcation diagrams in complex climate models. *Physical Review E* 107 054214 <https://journals.aps.org/pre/abstract/10.1103/PhysRevE.107.054214>

Dakos, Vasilis, Chris Boulton, Joshua Buxton et al. (2024) Tipping point detection and early warnings in climate, ecological, and human systems. *Earth System Dynamics* 15 1117—1135 <https://doi.org/10.5194/esd-15-1117-2024>

Palmer, TN, A Döring and G Seregin (2014) The real butterfly effect. *Nonlinearity* 27(9) 10.1088/0951-7715/27/9/R123 <https://iopscience.iop.org/article/10.1088/0951-7715/27/9/R123/meta>

Lenton, T.M., D.I. Armstrong McKay, S. Loriani, J.F. Abrams, S.J. Lade, J.F. Donges, M. Milkoreit, T. Powell, S.R. Smith, C. Zimm, J.E. Buxton, E. Bailey, L. Laybourn, A. Ghadiali, J.G. Dyke (eds), 2023, *The Global Tipping Points Report 2023*. University of Exeter, Exeter, UK.

From: Judith Curry [curry.judith@cfanclimate.com]
Sent: 5/18/2025 3:42:38 PM
To: John Christy [climateman60@gmail.com]; Roy W. Spencer [roywspencer@hotmail.com]; Ross McKittrick [ross.mckittrick@gmail.com]; Steven Koonin [steven.koonin@gmail.com]; Travis Fisher [travis.scott.fisher@gmail.com]
Subject: new text: declining planetary albedo
Attachments: Declining planetary albedo.docx

Ross suggested I write a section on the declining planetary albedo, for the attribution chapter. I'm sending the text for your comments before inserting into the chapter, since it is new material and I'm not sure how much of this we should actually include. In particular, I think Roy could be helpful here. Look forward to your comments.

Declining planetary albedo

Planetary albedo is the fraction of incoming solar radiation that is reflected back into space by the Earth's surface and atmosphere. Earth's planetary albedo is approximately 30%, meaning almost a third of the sunlight that reaches Earth is reflected back to space.

Arguably the most striking change to the Earth's climate system in the 21st century is a significant reduction in planetary albedo since 2015. Figure xxx shows the planetary albedo variations since 2000, when we have good satellite observations. The 0.5% reduction in planetary albedo since 2015 corresponds to an increase of 1.7 W/m² increase of solar energy averaged over the planet (Hansen and Karecha, 2025). For reference, doubling of CO₂ has a direct radiative impact of about 1.2 W/m².

Looking back prior to 2000 with less adequate data, Gossling et al. (2025) assessed that planetary albedo was relatively low around the 1940's and 50's, before industrial aerosol precursor emissions led to global dimming until the 1980's. The strongest planetary albedo excursions were high-albedo episodes caused by volcanic eruptions, such as after the Mount Pinatubo eruption in 1992. Although uncertain, the 2023 planetary albedo minimum may have been the lowest since at least 1940.




Figure xxx. Earth's albedo (reflectivity, in percent), with seasonality removed. From Hansen and Karecha (2025)

Changes in surface characteristics can't explain this decrease in planetary albedo since 2015:

- Arctic sea ice extent has declined by about 5% since 1980 (https://nsidc.org/data/seaice_index/images/s_plot_hires.png), although following 2007 there has been a pause in the Arctic sea ice decline (England et al., 2025)
- Regarding Antarctic sea ice, the IPCC AR6 concludes that "There has been no significant trend in Antarctic sea ice area from 1979 to 2020 due to regionally opposing trends and large internal variability." (SPM, A.1.5)
- Northern hemispheric snow cover has been slowly declining since 1967, with barely significant trends. The data show the Northern Hemisphere has snowier winters, accompanied by more rapid melt in spring and summer.
<http://climate.rutgers.edu/snowcover/>
- Global greening (Chapter 2) is contributing the decrease in planetary albedo, as forests have a lower albedo than open lands or snow. However, there is some evidence that forests increase

cloud cover (high reflectivity), which counteracts the direct albedo increase associated with increasing forested area.

Changes in surface reflectivity are contributing to a small, slow decline in planetary albedo. However, these changes cannot explain the sharp decline in planetary albedo beginning in 2015. Further, any changes in surface albedo are effectively damped by about a factor 3 on average, primarily due to cloud masking (Loeb et al. 2019). Surface albedo changes have thus contributed only weakly to the recent planetary albedo decline, particularly when averaged annually and globally.

This leaves the most likely explanation for the sharp albedo decline as being caused by changes in atmospheric aerosols and/or clouds.

Aerosols are small particles in the atmosphere that reflect sunlight and can interact with cloud processes to make the clouds more reflective. Aerosols have both natural origins (e.g dust from soils, wildfires) as well as human origins from combustion (including fossil fuels). Regions that are greening and/or have an increase in rainfall may be producing less dust. In terms of pollution aerosol, new ship fuel regulations aimed at reducing sulfur emissions were implemented in three phases, in 2010, 2015 and 2020. Sulfate particles reflect solar radiation. Cleaner air means less solar radiation is reflected, which contributes to surface warming. Indirect effects of sulfate aerosols include increasing the reflectivity of low-level clouds in the subtropics. There is controversy surrounding the importance of this change in sulfate emissions to the Earth's radiation balance (Hodnebrog et al., 2024; Yuan et al., 2024), but the restriction of this effect to major shipping routes suggests a small global impact (Schmidt, 2024).

This leaves changes in clouds as the primary candidate to explain the decline in global albedo since 2015. Two recent papers (Loeb et al., 2024; Gosling et al 2024) have addressed recent variations in cloud properties. By considering satellite and reanalysis data, Loeb et al. found that decreases in low and mid-level clouds since 2015 are the primary reason for decreasing planetary albedo in the northern hemisphere, whereas in the southern hemisphere the decrease in planetary albedo is primarily from decreases in mid-level clouds across all latitude zones. Gosling et al. (2025) found that cloud anomalies were mainly due to reduced low-level clouds. Regions with coherent low-level cloud reductions over the past decade include the warm pool region around the Maritime Continent and the northern extratropical western Pacific, as well as large parts of the Atlantic and adjacent land regions. The reduction of global cloud cover identified in these analyses since 2015 is 1-2 percent.

Two causes have been posited for the declining cloud cover over the past decade:

- Natural climate variability
- Changes in low cloud cover associated with warming sea surface temperatures, implying an emerging positive feedback to climate change (Hansen and Karecha, 2025)

The existence of a new positive low cloud feedback that began emerging in 2015 is not easy to justify. However, there are numerous natural climate signals during this period that are associated with atmospheric circulation changes that can influence the distribution of clouds:

- The 2014-2016 was the strongest El Nino event on record
- A cold anomaly beginning in 2015 in the subpolar gyre of the North Atlantic, reflect a shift in the ocean circulation pattern associated with decadal variability in the Atlantic (Frajka-Williams et al., 2017; Arthun et al. 2021).
- The Pacific Decadal Oscillation (PDO) positive index peaked in 2016, then declined and has been in negative territory since late 2019
- Eruption of the submarine Hunga-Tonga volcano in 2022

Interannual cloud anomalies associated with the El Niño Southern Oscillation (ENSO), which have a significant global signal and strong regional signals especially over the tropical Indian and Pacific Oceans.

The Hunga Tonga eruption is remarkable in its coincidence with the exceptionally low value of planetary albedo in 2023. Gossling et al. (2025) found that the cloud perturbations in 2023 showed a different pattern from the overall pattern since 2015, with reduced cloud cover being most pronounced in the northern hemisphere and tropics. Regional reductions in cloud cover were most pronounced over the eastern Indian Ocean, over South America and extending over the eastern Pacific, as well as over northern North America, in the Southern Ocean around 60S, in the subtropical and eastern North Atlantic, and in parts of the North Pacific.

The Hunga Tonga eruption was unusual in that it injected large amounts of water vapor into the stratosphere, in addition to sulfate particles. Early publications focused on the impacts to stratospheric circulations and the direct radiative impact. In 2025, papers are emerging that examine the indirect effects of Hunga Tonga on surface climate using ensemble simulations from earth system models (Bednarz et al, 2025; Kuchar et al., 2025). These papers found statistically significant impacts on regional climates that were driven by circulation changes driven by couplings between the stratosphere and troposphere. These papers indicate the complex interactions and between volcanic activity and climate dynamics. More research is needed to untangle any impacts of Hunga Tonga eruption on planetary albedo.

In summary, the decline in planetary albedo and the concurrent decline in cloudiness have emphasize the importance of clouds and their variations to global climate variability and change. A change of 1 to 2 percent in global cloud cover has a greater radiative impact on the climate than the direct radiative effect of doubling CO₂. While it is difficult to untangle causes of the trend, the competing explanations for the cause of the declining cloud cover have substantial implications for assessing the Equilibrium Climate Sensitivity and also the attribution of the recent warming. An additional 10 years of data should help clarify whether this is a strong positive cloud feedback associated with warming, or a temporary fluctuation driven by natural variability

References

- 35. S. A. Klein, A. Hall, J. R. Norris, R. Pincus, Low-cloud feedbacks from cloud-controlling factors: A review. *Surv Geophys* 38,1307-1329 (2017). doi:10.1007/s10712-017-9433-3
- 48. C. L. Boehm, D. W. J. Thompson, The Key Role of Cloud–Climate Coupling in Extratropical Sea Surface Tempe
- K. Li, F. Zheng, J. Zhu, Q.-C. Zeng, El Niño and the AMO Sparked the Astonishingly Large Margin of Warming in the Global Mean Surface Temperature in 2023. *Adv. Atmos. Sci.* 41, 1017-1022 (2024). doi:10.1007/s00376-023-3371-4

References

N. G. Loeb et al., Decomposing Shortwave Top-of-Atmosphere and Surface Radiative Flux Variations in Terms of Surface and Atmospheric Contributions. *Journal of Climate* 32(16), 5003-5019 (2019). doi:10.1175/JCLI-D-18-0826.1

G. Schmidt, Climate models can't explain 2023's huge heat anomaly — we could be in uncharted territory. *Nature* 627, 467 (2024).

Ø. Hodnebrog et al., Recent reductions in aerosol emissions have increased Earth's energy imbalance. *Commun Earth Environ* 5, 166 (2024). doi:10.1038/s43247-024-01324-8

13. T. Yuan, et al., Abrupt reduction in shipping emission as an inadvertent geoengineering termination shock produces substantial radiative warming. *Commun Earth Environ* 5, 281 (2024). doi:10.1038/s43247-024-01442-3

Mark England, Lorenzo M Polvani, James A Screen, et al. Surprising, but not unexpected, multi-decadal pause in Arctic sea ice loss. *ESS Open Archive* . March 29, 2025.

Hansen J and P Karecha (2025) Large cloud feedback confirms high sensitivity
<https://mailchi.mp/caa/large-cloud-feedback-confirms-high-climate-sensitivity>

G. Schmidt, Climate models can't explain 2023's huge heat anomaly — we could be in uncharted territory. *Nature* 627, 467 (2024).

Ø. Hodnebrog et al., Recent reductions in aerosol emissions have increased Earth's energy imbalance. *Commun Earth Environ* 5, 166 (2024). doi:10.1038/s43247-024-01324-8

13. T. Yuan, et al., Abrupt reduction in shipping emission as an inadvertent geoengineering termination shock produces substantial radiative warming. *Commun Earth Environ* 5, 281 (2024). doi:10.1038/s43247-024-01442-3

Loeb, N., S-H Ham, RP Allen, TJ Thorsen, B Meyssignac, S Kato, GC Johnson, JL Lyman,(2024) Observational Assessment of Changes in Earth's Energy Budget since 2000. *Surveys in Geophysics*, 45, 1757-1783

Desbruyeres, D, L Chafik and G Maze (2021) A shift in the ocean circulation has warmed the subpolar North Atlantic Ocean since 2016. *Communications Earth&Environment*.

Arthun, M, RCJ Willis, HL Johnson, L Chafik, HR Langehaug (2021) Mechanisms of Decadal North Atlantic Climate Variability and Implications for the Recent Cold Anomaly. *Journal of Climate*, Vol 34

Kuchar, A., Sukhodolov, T., Chiodo, G., Jörimann, A., Kult-Herdin, J., Rozanov, E., and Rieder, H. H.: Modulation of the northern polar vortex by the Hunga Tonga–Hunga Ha'apai eruption and the associated surface response, *Atmos. Chem. Phys.*, 25, 3623–3634, <https://doi.org/10.5194/acp-25-3623-2025>, 2025.

Bednarz, EW, AH Butler, X Wang, Z Zhou, W Yu, G Stenchikov, M Toohey, Y Zhu (2025) Indirect climate impacts of the Hunga eruption. Preprint, <https://doi.org/10.5194/egusphere-2025-1970>

Kuchar, A., Sukhodolov, T., Chiodo, G., Jörimann, A., Kult-Herdin, J., Rozanov, E., and Rieder, H. H.: Modulation of the northern polar vortex by the Hunga Tonga–Hunga Ha'apai eruption and the associated surface response, *Atmos. Chem. Phys.*, 25, 3623–3634, <https://doi.org/10.5194/acp-25-3623-2025>, 2025.

identify a decadal mode of atmosphere–ocean variability in the North Atlantic with a dominant time scale of 13–18 years. Large-scale atmospheric circulation anomalies drive SST anomalies both through contemporaneous air–sea heat fluxes and through delayed ocean circulation changes, the latter involving both the meridional overturning circulation and the horizontal gyre circulation.

From: John Christy <climateman60@gmail.com>
Sent: Thursday, August 14, 2025 2:30 PM
To: Steve Koonin; Judith Curry; Roy Spencer; Ross McKittrick
Subject: Now this is interesting

All

I know there is a plethora of press stories about our mental abilities and character, but I was not expecting this one.

<https://www.perplexity.ai/search/b62c4bc5-0657-4e6c-a467-7d43d756e153>

John C.

From: John Christy [climateman60@gmail.com]
Sent: 5/20/2025 3:55:47 PM
To: Ross McKittrick [ross.mckittrick@gmail.com]; Roy Spencer [roywspencer@hotmail.com]; Steve Koonin [steven.koonin@gmail.com]; Judith Curry [curry.judith@gmail.com]; Travis Fisher [travis.scott.fisher@gmail.com]
Subject: Obs and Models, global LT and MT, 1979-2024
Attachments: LT_MT_CMIP6_Obs.xlsx

Ross and Roy

Here are the annual anomalies of global LT and MT for observations and CMIP6 models. I have not put them in a common reference base, so you'll need to subtract out whatever reference period you use to put the anomalies relative to a common reference.

John C.

Global LT

	Sonde	Sonde	Sonde	Satellite	Satellite	Satellite	Reanalysis	Reanalysis
	RAOBCOR	RICH	RATPAC A	UAH_6.0	RSS	NOAA	ERA5	JRA3Q
1979	-0.360	-0.442	-0.414	-0.346	-0.495		-0.303	-0.405
1980	-0.213	-0.287	-0.275	-0.178	-0.344		-0.235	-0.303
1981	-0.240	-0.306	-0.282	-0.248	-0.358	-0.272	-0.261	-0.337
1982	-0.466	-0.530	-0.518	-0.433	-0.560	-0.393	-0.466	-0.481
1983	-0.267	-0.321	-0.362	-0.178	-0.327	-0.131	-0.235	-0.268
1984	-0.533	-0.586	-0.567	-0.374	-0.590	-0.402	-0.504	-0.547
1985	-0.493	-0.553	-0.530	-0.498	-0.645	-0.438	-0.535	-0.566
1986	-0.403	-0.481	-0.487	-0.356	-0.528	-0.274	-0.413	-0.430
1987	-0.125	-0.201	-0.242	-0.087	-0.233	0.033	-0.135	-0.130
1988	-0.112	-0.160	-0.196	-0.098	-0.264	0.012	-0.090	-0.105
1989	-0.373	-0.409	-0.446	-0.346	-0.467	-0.234	-0.318	-0.359
1990	-0.085	-0.113	-0.210	-0.124	-0.290	-0.015	-0.065	-0.119
1991	-0.083	-0.112	-0.245	-0.116	-0.245	0.006	-0.090	-0.130
1992	-0.414	-0.447	-0.512	-0.417	-0.518	-0.289	-0.376	-0.408
1993	-0.330	-0.366	-0.446	-0.339	-0.453	-0.275	-0.346	-0.392
1994	-0.266	-0.297	-0.293	-0.201	-0.318	-0.152	-0.255	-0.265
1995	-0.201	-0.227	-0.190	-0.067	-0.177	-0.053	-0.141	-0.146
1996	-0.324	-0.343	-0.369	-0.146	-0.278	-0.156	-0.246	-0.231
1997	-0.298	-0.316	-0.324	-0.144	-0.242	-0.159	-0.245	-0.228
1998	0.232	0.220	0.155	0.347	0.224	0.306	0.220	0.260
1999	-0.253	-0.259	-0.303	-0.153	-0.243	-0.211	-0.280	-0.215
2000	-0.284	-0.291	-0.326	-0.157	-0.248	-0.221	-0.315	-0.243
2001	-0.119	-0.137	-0.132	-0.022	-0.079	-0.061	-0.130	-0.063
2002	0.016	-0.006	-0.031	0.080	0.033	0.035	-0.001	0.033
2003	0.009	-0.008	0.024	0.050	0.048	0.014	-0.012	0.033
2004	-0.088	-0.091	-0.081	-0.056	-0.024	-0.088	-0.104	-0.049
2005	0.093	0.106	0.141	0.063	0.110	0.040	0.044	0.118
2006	-0.017	0.000	0.065	-0.023	0.020	-0.044	-0.035	0.018
2007	0.042	0.059	0.050	0.023	0.055	0.000	0.015	-0.005
2008	-0.254	-0.239	-0.199	-0.238	-0.153	-0.250	-0.240	-0.252
2009	-0.012	-0.001	0.013	-0.043	0.009	-0.043	-0.027	-0.049
2010	0.140	0.151	0.233	0.194	0.254	0.197	0.250	0.223
2011	-0.130	-0.115	-0.034	-0.120	-0.072	-0.109	-0.075	-0.071
2012	-0.059	-0.044	0.022	-0.084	-0.006	-0.084	-0.035	-0.036
2013	0.073	0.091	0.136	-0.001	0.069	0.004	0.068	0.058

2014	0.117	0.138	0.167	0.044	0.127	0.056	0.113	0.092
2015	0.270	0.287	0.302	0.137	0.260	0.154	0.221	0.197
2016	0.595	0.610	0.554	0.388	0.456	0.389	0.499	0.463
2017	0.355	0.373	0.382	0.265	0.332	0.240	0.359	0.307
2018	0.249	0.268	0.264	0.088	0.193	0.101	0.233	0.172
2019	0.470	0.492	0.481	0.296	0.401	0.293	0.437	0.368
2020	0.475	0.505	0.495	0.351	0.465	0.361	0.495	0.433
2021	0.238	0.275	0.298	0.138	0.271	0.184	0.290	0.234
2022	0.267	0.228	0.320	0.150	0.254	0.192	0.300	0.282
2023	0.515	0.530	0.573	0.432	0.551	0.475	0.585	0.571
2024	0.860	0.875	0.924	0.774	0.860	0.784	0.914	0.884
2025								
2026								
2027								
2028								
2029								
2030								

	ACCESS_C	ACCESS_E	AWI	BCC-CSM	CAMS	Can5	Can5-OE	CESM
1979	-0.190	-0.443	-0.313	-0.297	-0.210	-0.639	-0.613	0.026
1980	-0.269	-0.240	-0.356	-0.367	-0.092	-0.562	-0.634	0.047
1981	-0.193	-0.293	-0.293	-0.137	-0.159	-0.395	-0.552	-0.486
1982	-0.343	-0.479	-0.435	-0.123	-0.177	-0.262	-0.392	-0.469
1983	-0.458	-0.383	-0.433	-0.219	-0.178	-0.407	-0.596	-0.419
1984	-0.190	-0.292	-0.437	-0.179	-0.321	-0.464	-0.520	-0.373
1985	-0.130	-0.488	-0.268	-0.373	-0.522	-0.496	-0.186	0.052
1986	0.107	-0.458	-0.323	-0.263	-0.095	-0.465	-0.338	-0.263
1987	-0.319	-0.277	-0.140	-0.199	-0.017	-0.255	-0.359	-0.247
1988	-0.232	-0.328	-0.300	-0.190	-0.164	-0.237	-0.356	-0.079
1989	-0.130	-0.247	-0.040	0.002	0.056	-0.218	-0.037	-0.088
1990	0.036	-0.186	0.074	-0.049	0.142	-0.156	0.020	-0.167
1991	0.041	-0.261	-0.226	-0.263	0.014	-0.193	0.012	-0.257
1992	-0.517	-0.555	-0.545	-0.636	-0.397	-0.406	-0.595	-0.553
1993	-0.351	-0.325	-0.436	-0.377	-0.397	-0.559	-0.388	-0.270
1994	-0.178	-0.450	-0.198	-0.121	-0.168	-0.278	-0.452	0.034
1995	-0.193	-0.235	0.061	0.097	-0.060	-0.199	-0.224	-0.208
1996	-0.040	0.012	0.128	-0.258	-0.094	-0.038	0.166	-0.073
1997	-0.047	-0.014	-0.096	0.134	0.160	0.138	-0.003	0.183
1998	-0.029	0.206	-0.006	-0.028	0.247	0.197	-0.068	-0.111
1999	0.101	0.343	-0.020	-0.010	-0.019	0.333	0.128	0.069
2000	0.300	0.257	0.228	0.173	0.053	0.277	0.277	0.134
2001	0.199	0.458	0.272	0.324	-0.013	0.263	0.503	-0.051
2002	0.173	0.613	0.207	0.385	0.195	0.349	0.547	0.130
2003	0.366	0.393	0.368	0.309	0.318	0.387	0.464	0.442
2004	0.289	0.478	0.386	0.201	0.348	0.319	0.411	0.516
2005	0.334	0.443	0.228	0.300	0.371	0.405	0.306	0.335
2006	0.388	0.412	0.238	0.282	0.319	0.193	0.132	0.208
2007	0.097	0.231	0.406	0.307	0.287	0.306	0.326	0.327
2008	0.229	0.405	0.536	0.245	0.102	0.360	0.730	0.404
2009	0.238	0.493	0.485	0.342	0.093	0.552	0.486	0.524
2010	0.441	0.529	0.593	0.310	0.082	0.943	0.549	0.743
2011	0.555	0.550	0.352	0.326	0.231	0.878	0.577	0.858
2012	0.762	0.469	0.459	0.374	0.250	0.608	0.791	0.452
2013	0.387	0.666	0.551	0.443	0.498	0.807	0.586	0.500

2014	0.542	0.756	0.491	0.510	0.398	0.956	0.814	0.697
2015	0.789	0.744	0.450	0.484	0.400	0.812	0.865	0.610
2016	0.588	0.663	0.617	0.466	0.560	0.754	0.960	0.659
2017	0.743	0.832	0.716	0.608	0.353	1.030	0.862	0.841
2018	0.928	0.858	0.727	0.630	0.467	0.945	0.852	0.661
2019	0.778	0.656	0.773	0.506	0.562	0.875	1.173	0.613
2020	0.868	0.621	0.579	0.673	0.600	1.237	1.105	0.656
2021	0.864	0.561	0.817	0.694	0.811	1.107	1.291	0.599
2022	1.081	0.832	0.940	0.734	0.691	1.087	1.221	0.796
2023	0.952	0.906	0.961	0.791	0.397	1.237	1.275	1.226
2024	1.101	1.064	0.956	0.680	0.449	1.293	1.265	1.034
2025	1.176	0.961	0.908	0.637	0.680	1.497	1.475	1.173
2026	1.096	1.221	0.954	0.953	0.462	1.794	1.604	1.253
2027	1.271	1.319	0.900	0.940	0.657	1.622	1.507	0.679
2028	1.357	1.294	0.967	1.031	0.458	1.517	1.544	0.991
2029	1.352	1.426	1.002	0.925	0.303	1.465	1.785	1.064
2030	1.274	1.257	1.039	1.004	0.283	1.477	1.624	1.224

CESM2-W	CIESM	CIESM	CNRM-CN	CNRM_HI	CNRM-ES	EC-EARTH	EC-E3-VE	FGOALS-f	FGOALS-g
-0.288	-0.434	-0.250	-0.334	-0.282	-0.153	-0.018	-0.453	-0.416	-0.399
-0.459	-0.277	-0.048	-0.117	-0.232	-0.238	0.112	-0.262	-0.130	-0.158
-0.093	-0.155	-0.030	-0.194	-0.175	-0.339	-0.288	0.028	-0.151	-0.198
-0.222	-0.428	-0.246	-0.335	-0.051	-0.282	-0.372	-0.396	-0.395	-0.273
-0.554	-0.561	-0.169	-0.200	-0.421	-0.171	-0.472	-0.484	-0.290	-0.410
-0.285	-0.254	-0.125	-0.357	-0.389	-0.274	-0.058	-0.293	-0.217	-0.392
-0.325	-0.060	-0.298	-0.261	-0.175	-0.269	-0.416	-0.333	-0.622	-0.305
-0.501	-0.357	-0.406	-0.288	-0.330	0.001	-0.476	-0.273	-0.218	-0.497
-0.471	-0.231	-0.181	-0.048	-0.188	-0.191	-0.311	-0.335	-0.076	-0.108
-0.132	-0.083	-0.104	-0.037	-0.157	-0.061	-0.107	-0.328	-0.344	0.005
-0.038	-0.148	-0.148	-0.110	-0.053	-0.238	0.117	-0.174	-0.184	-0.197
-0.223	0.046	-0.047	0.023	-0.046	-0.165	0.205	-0.005	0.206	-0.064
-0.323	0.111	0.166	-0.056	-0.061	0.077	-0.163	0.215	-0.250	0.052
-0.199	-0.454	-0.454	-0.236	-0.209	-0.192	-0.755	-0.570	-0.547	-0.633
-0.147	-0.455	-0.433	-0.006	-0.145	-0.255	-0.663	-0.530	-0.057	-0.481
-0.158	-0.172	-0.550	-0.369	-0.028	0.176	-0.346	-0.214	-0.015	-0.158
-0.122	0.079	-0.463	-0.210	-0.062	-0.111	0.105	-0.134	-0.147	-0.301
-0.131	-0.151	-0.273	0.114	-0.023	-0.031	-0.255	0.058	0.026	-0.115
-0.024	0.015	-0.164	-0.183	0.063	0.116	-0.144	0.213	0.145	0.220
-0.089	0.185	-0.096	0.077	0.003	-0.146	0.007	0.315	0.110	0.024
0.046	0.249	-0.077	0.119	0.115	0.158	0.183	0.082	0.187	0.093
0.275	0.033	-0.031	0.241	-0.010	0.043	0.272	0.051	0.074	0.232
0.498	0.186	0.185	0.335	0.077	0.017	0.258	0.285	0.110	0.112
0.420	0.372	0.445	0.263	0.152	0.053	0.241	0.279	0.416	0.347
0.407	0.117	0.542	0.166	0.262	0.249	0.277	0.345	0.122	0.356
0.319	0.155	0.241	0.353	0.120	0.037	0.204	0.435	0.387	0.451
0.247	0.405	0.197	-0.058	0.251	0.310	0.639	0.262	0.513	0.395
0.211	0.546	0.357	0.137	0.142	0.074	0.687	0.345	-0.057	0.324
0.498	0.245	0.537	0.501	0.300	0.522	0.301	0.415	0.306	0.357
0.488	0.155	0.633	0.316	0.285	0.390	0.296	0.290	0.672	0.311
0.252	0.252	0.543	0.130	0.300	0.096	0.377	0.187	0.085	0.487
0.380	0.371	0.441	0.181	0.450	0.402	0.645	0.268	0.201	0.356
0.550	0.262	0.502	0.311	0.301	0.489	0.413	0.382	0.537	0.381
0.672	0.338	0.597	0.487	0.425	0.330	0.497	0.574	0.357	0.524
0.508	0.551	0.670	0.306	0.379	0.560	0.606	0.440	0.462	0.234

0.537	0.604	0.466	0.415	0.434	0.530	0.786	0.545	0.626	0.518
0.707	0.218	0.616	0.420	0.530	0.459	0.393	0.594	0.372	0.235
0.857	0.256	0.928	0.590	0.406	0.705	0.525	0.606	0.514	0.778
0.845	0.563	0.989	0.706	0.519	0.593	0.402	0.686	0.457	0.781
0.957	0.680	0.242	0.377	0.725	0.668	0.671	0.798	0.523	0.772
0.582	0.625	0.033	0.532	0.654	0.866	0.770	0.908	0.885	0.785
0.489	0.561	0.009	0.767	0.605	0.684	0.459	0.923	0.730	0.632
0.927	0.687	0.488	0.722	0.621	0.825	0.575	0.694	0.549	0.505
1.044	0.739	0.820	0.849	0.747	0.594	0.808	0.674	1.009	0.664
0.772	0.853	0.697	0.746	0.807	0.952	0.609	0.745	1.018	0.607
0.933	0.633	0.706	0.829	0.968	1.165	0.744	0.913	0.633	0.957
1.181	0.816	0.616	0.978	0.530	0.970	0.835	1.019	1.024	0.748
1.092	1.064	0.679	0.783	0.406	1.090	0.850	0.724	1.161	0.852
1.027	0.967	0.856	0.923	0.519	1.039	0.723	0.833	0.689	0.751
1.268	1.113	1.047	1.127	0.725	1.228	0.717	0.867	0.948	0.681
1.467	1.101	0.924	0.767	0.654	0.930	0.827	0.957	1.190	0.944
1.062	1.127	0.859	0.909	0.605	1.183	0.993	1.057	0.929	0.783

FIO	GFDL-CM	GFDL-ESM	GISS	HadGEM	IITM	INM-CM4	INM-CM5	IPSL-CM6	KACE
-0.498	-0.173	-0.194	0.009	-0.460	-0.199	-0.148	-0.136	-0.296	-0.472
-0.409	-0.212	-0.394	-0.077	-0.349	-0.308	-0.197	-0.104	-0.195	-0.321
-0.363	-0.538	-0.187	-0.289	-0.456	-0.273	-0.187	-0.237	-0.178	-0.207
-0.488	-0.322	0.057	-0.412	-0.350	-0.214	-0.245	-0.208	-0.569	-0.403
-0.600	-0.359	-0.389	-0.089	-0.497	-0.296	-0.667	-0.636	-0.695	-0.487
-0.267	-0.369	-0.440	0.013	-0.248	-0.092	-0.398	-0.458	-0.509	-0.239
-0.148	-0.504	-0.419	-0.425	-0.392	-0.322	-0.264	-0.401	-0.653	-0.363
-0.399	-0.250	-0.228	-0.572	-0.264	-0.171	-0.218	-0.195	-0.378	-0.216
-0.193	0.053	-0.234	-0.325	-0.345	0.036	-0.232	-0.213	-0.039	-0.193
-0.098	-0.104	-0.298	0.137	-0.303	-0.016	-0.120	-0.129	0.136	-0.024
-0.029	-0.079	-0.156	0.263	-0.227	-0.154	-0.026	-0.013	0.405	-0.150
0.184	-0.113	0.211	0.147	-0.047	0.010	0.069	0.001	-0.121	0.020
0.020	-0.135	0.144	-0.147	0.001	0.092	0.056	0.084	-0.278	-0.146
-0.289	-0.410	-0.635	-0.652	-0.336	-0.382	-0.461	-0.386	-0.342	-0.403
-0.307	-0.396	-0.554	-0.344	-0.377	-0.163	-0.468	-0.459	-0.340	-0.338
-0.301	-0.376	-0.118	0.166	-0.384	-0.280	-0.274	-0.237	-0.138	-0.174
-0.154	-0.217	-0.118	-0.111	-0.248	-0.143	-0.057	-0.091	-0.001	-0.141
-0.021	-0.063	-0.126	-0.390	-0.119	-0.110	0.021	-0.035	0.125	-0.064
-0.007	0.124	0.002	-0.091	-0.041	-0.106	0.078	0.116	0.139	0.027
0.061	0.105	0.027	0.212	0.023	-0.040	0.106	0.153	-0.048	0.021
0.119	-0.034	0.166	0.287	0.011	0.100	0.210	0.151	0.056	0.130
0.237	0.258	0.222	-0.142	0.122	0.241	0.268	0.191	0.268	0.065
0.393	0.331	0.322	-0.182	0.315	0.313	0.246	0.193	0.181	0.130
0.126	0.232	0.365	0.210	0.439	0.108	0.139	0.250	0.113	0.270
0.148	0.236	0.195	0.279	0.265	0.310	0.272	0.319	0.105	0.341
0.238	0.459	0.349	0.037	0.423	0.283	0.336	0.371	0.083	0.355
0.422	0.333	0.300	-0.010	0.405	0.369	0.273	0.323	0.435	0.324
0.369	0.447	0.109	0.289	0.614	0.235	0.253	0.326	0.497	0.243
0.333	0.479	0.213	0.623	0.542	0.362	0.261	0.325	0.176	0.351
0.300	0.469	0.352	0.569	0.305	-0.037	0.270	0.277	0.126	0.384
0.346	0.270	0.506	0.575	0.476	0.102	0.339	0.354	0.635	0.421
0.362	0.479	0.348	0.374	0.685	0.246	0.425	0.277	0.822	0.453
0.434	0.632	0.348	0.217	0.910	0.349	0.387	0.258	0.324	0.492
0.397	0.565	0.578	0.468	0.831	0.466	0.384	0.319	0.595	0.494
0.227	0.531	0.799	0.526	0.637	0.201	0.461	0.346	0.634	0.451

0.367	0.672	0.381	-0.041	0.886	0.372	0.352	0.309	0.378	0.579
0.476	0.763	0.379	0.026	0.968	0.427	0.379	0.454	0.340	0.768
0.612	0.666	0.576	0.566	1.007	0.594	0.356	0.469	0.442	0.778
0.546	0.529	0.777	0.813	0.945	0.278	0.425	0.432	0.812	0.967
0.650	0.625	0.902	0.422	1.067	0.445	0.356	0.397	0.874	0.864
0.853	0.645	0.990	0.154	0.989	0.597	0.461	0.502	0.627	1.002
0.681	0.740	0.499	0.410	1.075	0.336	0.544	0.459	0.758	1.003
0.837	0.757	0.579	0.943	1.242	0.522	0.606	0.634	0.836	1.197
0.767	0.975	0.796	0.916	1.536	0.533	0.595	0.676	0.591	1.233
0.824	0.926	0.968	0.862	1.348	0.199	0.603	0.600	0.716	1.240
0.655	0.923	0.977	0.682	1.385	0.517	0.842	0.755	0.960	1.302
0.730	1.018	0.861	1.069	1.503	0.636	0.799	0.700	1.029	1.396
0.968	1.196	1.057	1.206	1.541	0.938	0.830	0.703	1.184	1.410
0.948	1.047	0.904	0.659	1.544	0.507	0.788	0.746	1.036	1.431
1.000	1.080	0.844	0.338	1.626	0.779	0.799	0.858	0.931	1.493
0.928	1.219	0.864	0.681	1.627	0.704	0.820	0.808	0.996	1.498
0.925	1.354	1.065	1.170	1.504	0.566	0.823	0.924	0.810	1.498


KIOST	MCM-UA	MIROC6	MIROC6-2	MPI-HR	MPI-LR	MRI-ESM	NESM	NOR-LM	UKESM1
-0.248	-0.458	-0.170	-0.287	-0.274	-0.439	-0.237	-0.287	-0.450	-0.480
-0.488	-0.397	-0.410	0.019	0.043	-0.117	-0.366	-0.346	-0.285	-0.564
-0.348	-0.394	-0.475	0.161	-0.085	-0.178	-0.174	-0.221	-0.261	-0.433
-0.286	-0.452	-0.146	-0.458	-0.238	-0.351	-0.131	-0.217	-0.127	-0.390
-0.406	-0.557	-0.131	-0.626	-0.394	-0.466	-0.265	-0.733	-0.065	-0.484
-0.533	-0.327	0.191	-0.312	-0.309	-0.174	-0.407	-0.377	-0.203	-0.398
-0.323	-0.265	-0.097	0.112	-0.143	-0.062	-0.053	-0.503	-0.333	-0.508
-0.268	-0.290	-0.432	0.188	0.076	-0.030	-0.049	-0.407	-0.186	-0.314
-0.189	-0.312	-0.176	-0.371	0.001	-0.223	-0.213	-0.321	-0.384	-0.303
0.127	-0.190	-0.045	-0.373	-0.023	-0.306	-0.009	-0.159	-0.395	-0.437
-0.144	-0.262	-0.158	-0.231	-0.084	-0.205	-0.235	0.046	-0.163	-0.184
0.012	-0.178	-0.129	0.093	-0.018	0.004	-0.178	0.203	-0.074	-0.004
0.246	-0.285	-0.165	0.412	-0.164	0.020	-0.021	0.007	0.038	-0.260
-0.682	-0.397	-0.455	-0.530	-0.497	-0.406	-0.453	-0.569	-0.338	-0.698
-0.233	-0.301	-0.177	-0.713	-0.408	-0.136	-0.277	-0.514	-0.490	-0.434
-0.379	-0.248	0.128	-0.378	-0.064	-0.213	-0.212	-0.128	-0.367	-0.090
-0.181	-0.214	-0.307	-0.021	0.025	-0.339	-0.023	-0.161	-0.198	-0.128
-0.010	0.075	-0.474	0.168	-0.078	-0.064	-0.223	-0.045	-0.047	-0.082
-0.263	0.279	-0.210	0.383	-0.247	0.250	-0.097	-0.033	0.120	0.028
0.073	0.250	-0.076	-0.219	-0.023	0.127	0.111	0.202	0.205	0.288
0.176	0.280	0.416	-0.321	0.066	0.146	0.005	0.379	0.276	0.218
0.132	0.396	0.252	-0.070	0.087	0.107	0.093	0.214	0.446	0.252
0.372	0.252	0.086	0.384	0.251	0.035	0.331	0.061	0.529	0.305
0.012	0.220	-0.015	0.200	0.315	0.090	0.057	0.361	0.023	0.548
0.183	0.402	0.306	0.122	0.142	0.391	0.049	0.418	-0.032	0.263
0.380	0.637	0.628	0.659	0.109	0.319	0.200	0.325	0.171	0.313
0.338	0.618	0.211	0.658	0.117	0.107	0.520	0.263	0.145	0.417
0.217	0.185	0.049	-0.054	0.240	0.135	0.332	0.405	0.044	0.450
0.287	0.298	0.101	-0.115	0.176	0.329	0.319	0.326	0.248	0.478
0.664	0.332	0.537	0.156	0.356	0.451	0.451	0.443	0.460	0.467
0.646	0.257	0.690	0.324	0.461	0.319	0.339	0.395	0.502	0.517
0.371	0.180	0.062	0.773	0.340	0.325	0.201	0.334	0.445	0.596
0.708	0.329	0.039	0.383	0.416	0.390	0.194	0.683	0.593	0.642
0.803	0.654	0.240	0.164	0.521	0.437	0.380	0.780	0.484	0.615
0.664	0.744	0.799	0.265	0.598	0.293	0.412	0.885	0.592	0.687

0.813	0.731	0.724	0.519	0.494	0.275	0.338	0.522	0.870	0.826
0.613	0.895	0.044	0.894	0.312	0.400	0.522	0.912	0.597	0.797
0.409	0.856	-0.057	0.283	0.345	0.527	0.628	0.839	0.508	0.982
0.484	0.629	0.314	0.177	0.282	0.529	0.851	0.822	0.593	0.840
0.373	0.403	0.762	0.387	0.493	0.403	0.519	0.879	0.728	0.841
0.687	0.337	0.353	0.707	0.747	0.171	0.475	0.954	0.923	1.051
0.472	0.468	0.236	1.063	0.674	0.360	0.593	1.031	0.873	1.140
0.838	0.695	0.289	0.650	0.714	0.815	0.677	1.347	0.746	0.867
0.593	0.815	0.423	0.879	0.815	0.607	1.008	1.201	0.948	1.120
0.773	0.700	0.676	1.281	0.632	0.573	1.040	1.275	0.634	1.348
0.996	0.653	1.004	0.884	0.788	0.840	0.896	1.372	0.511	1.304
0.537	0.909	0.513	0.479	0.838	0.608	0.910	1.490	0.531	1.444
0.923	0.904	0.500	0.516	0.514	0.556	1.050	1.229	0.396	1.381
1.010	0.851	0.674	0.733	0.560	0.532	1.078	1.097	0.431	1.376
0.777	0.828	1.041	1.100	0.678	0.711	1.088	0.962	0.789	1.422
1.041	0.909	1.293	1.289	0.952	0.795	1.187	1.235	0.998	1.558
0.979	1.085	0.602	0.629	0.795	0.950	0.943	1.201	0.986	1.579

From: Judith Curry [curry.judith@gmail.com]
Sent: 6/9/2025 10:48:30 PM
To: Steven Koonin [steven.koonin@gmail.com]
CC: John Christy [climateman60@gmail.com]; Ross McKittrick [ross.mckittrick@gmail.com]; Roy W. Spencer [roywspencer@hotmail.com]; Travis Fisher [travis.scott.fisher@gmail.com]; Josh Loucks [loucksj14@gmail.com]
Subject: overconfidence
Attachments: Overconfidence when stating conclusions.docx

see attached text for overconfidence. I moved the hurricane example here. Could use some more examples

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

Overconfidence when stating conclusions

The NCA5 uses specific terms to convey information about scientific confidence associated with important finds, observations and projections. The Front Matter of the NCA5 states that “confidence in a finding is based on the type, amount, quality, strength, and consistency of evidence; the skill, range, and consistency of methods to detect, evaluate, attribute, and interpret climate trends; and the degree of agreement across scientific information sources.”

Table 1 provides calibrated language for confidence assessment:

Very high:	Strong evidence (established theory, multiple sources, well documented and accepted methods, etc.). High consensus
High:	Moderate evidence (several sources, some consistency, methods vary and/or documentation limited, etc.). Medium consensus
Medium:	Suggestive evidence (a few sources, limited consistency, methods emerging, etc.) Competing schools of thought.
Low:	Inconclusive evidence (limited sources, extrapolations, inconsistent findings, poor documentation and/or methods not tested, etc.). Disagreement or lack of opinions among experts.

This terminology is inflated: *high confidence* with just moderate evidence and medium consensus; *medium confidence* with only suggested evidence and competing schools of thought. This inflation in terminology leads to overconfidence in the conclusions.

Even within the context of this inflated terminology, NCA5 further inflates the confidence level of many of its conclusions to well beyond what the IPCC has concluded with more thorough justification, and also by incorrectly applying its own standards. Several illustrative examples are provided.

Key Message 3.1 Traceable Account states: “There is *very high confidence* that emissions of GHGs from human activities, fossil fuel use in particular, have unequivocally caused *all* global warming observed over the industrial era.” Compare the NCA5 statement with statements from the IPCC AR5 and AR6:

AR5: “It is *extremely likely* that *more than half* of the observed increase in global average surface temperature from 1951 to 2010 was caused by the anthropogenic increase in greenhouse gas concentrations and other anthropogenic forcings together.” (IPCC, 2013)

AR6: “The *likely* range of total human-caused global surface temperature increase from 1850–1900 to 2010–2019 is 0.8°C to 1.3°C, with a best estimate of 1.07°C. It is *likely* that well-mixed GHGs contributed a warming of 1.0°C to 2.0°C, other human drivers (principally aerosols) contributed a cooling of 0.0°C to 0.8°C, natural drivers changed global surface temperature by –0.1°C to +0.1°C, and internal variability changed it by –0.2°C to +0.2°C. It is

very likely that well-mixed GHGs were the main driver of tropospheric warming since 1979.” (IPCC, 2021)

The IPCC AR5 and AR6 statements provide likelihood assessments but not a confidence level. While the NCA5 claims with *very high confidence* that fossil fuels have unequivocally caused *all* global warming, the AR5 refers to “*more than half* of the observed warming” and the AR6 provides a more nuanced statement that relates to a *likely* range of warming contributed to by human drivers.

NCA5 Key message 2.2: “Hurricanes have been intensifying more rapidly since the 1980s (*high confidence*) and causing heavier rainfall and higher storm surges (*high confidence*).” No justification is provided in Chapter 2 for the *high confidence* conclusion on higher storm surges or hurricanes causing heavier precipitation. The Traceable Accounts section, Description of Confidence and Likelihood states: “Basic physical understanding and climate models both provide robust explanations for the links between climate change and observed changes in these extremes: this is why the authors also have *high confidence* that storms are delivering more rainfall and high confidence that storm surges are becoming higher.” In short, the NCA5 provides no observational basis or published research for their *high confidence* assertions that hurricanes are causing heavier rainfall and higher storm surges; and does not meet its own standards for a *high confidence level*.


From: Judith Curry <curry.judith@gmail.com>
Sent: Monday, June 16, 2025 10:55 AM
To: Steven Koonin
Cc: Ross McKittrick; John Christy; Roy W. Spencer; Travis Fisher
Subject: Pielke's latest

Excellent stuff in Pielke's latest substack, seems a shame not to integrate some of this into Ch 11, 12 of CWG report

<https://rogerpielkejr.substack.com/p/the-most-amazing-climate-policy-figure>

...

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Wednesday, May 7, 2025 12:40 PM
To: Travis Fisher; Judith Curry
Cc: Steven Koonin; John Christy; Ross McKittrick
Subject: Politico's article on those who have influenced Sec. Wright

For your entertainment and viewing pleasure, here's my blog response, which links to this morning's Politico article.

-Roy

<https://www.drroyspencer.com/2025/05/politicos-hit-piece-on-me-and-energy-secretary-wright/>

From: Travis Fisher <travis.scott.fisher@gmail.com>
Sent: Wednesday, May 7, 2025 10:37 AM
To: Judith Curry <curry.judith@gmail.com>
Cc: Roy Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>; John Christy <climateman60@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>
Subject: Re: Glossary

I'd be happy to take on the next version of the glossary. You don't need a PhD to do that, and I'm happy to take anything like that off your hands!

On Wed, May 7, 2025 at 10:31 AM Judith Curry <curry.judith@gmail.com> wrote:
thx for doing this. There are alot of acronyms used in the various chapters that should be included also

On Wed, May 7, 2025 at 3:45 AM Roy Spencer <roywspencer@hotmail.com> wrote:
Attached is my first cut at a glossary, 64 terms so far based upon my reading of the chapters.

I've omitted obvious things like EPA, CO2, NOAA, NASA...

I'm sure I missed a few terms, and feel free to make edits.

Of course, this could all wait until everything else is completed.

-Roy

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Tuesday, May 6, 2025 9:02 PM
To: John Christy <climateman60@gmail.com>
Cc: Ross McKittrick <ross.mckittrick@gmail.com>; Judith Curry <curry.judith@gmail.com>; Roy W. Spencer <roywspencer@hotmail.com>; Travis Fisher <travis.scott.fisher@gmail.com>
Subject: Re: Ch 8 Extreme weather

I was expecting to read/comment/edit Chapter 8 tomorrow (Wednesday), so I'll use this version. Roy, if you beat me to a summary, that's fine, else I'll take it on.

John, do decouple for at least a few days. I've made the mistake of not doing so on a few vacations, with great regrets afterward.

Steven Koonin

On May 6, 2025, at 21:53, John Christy <climateman60@gmail.com> wrote:

All

Here is a cleaned up version of chapter 8 which you will likely not see until 7 May, so that is the date.

My preference is for someone to read the thing and write the summary (8.4) based upon what a fresh set of eyes would conclude. Steve? Roy?

Ross. The panel charts of the individual station precipitation take up a lot of real estate and I think the value per acre is low. Since you calculated 3 statistics for each station, could you craft a bar chart or something to show that result? You would then have 9 charts, one for each of the 3 statistics and one each of the three regions. The nine charts may be panelized and fit on one plot. I know for us it's important for people to see the actual data (which is why I had those colored bar charts - now monochromatic), but it may be overkill for this document.

As to one of the statistics, I believe 3-day totals will be more meaningful and accurate than 1-day totals. I could send you a file of the heaviest 3-day total for each station and each year and you can do the analysis from there.

I especially like the way you parsed the Sidebar and I hope smart people will finally understand that when dealing with climate, our sample is pathetically small from which to draw hard conclusions.

I also enjoyed the added words on LTP. In fact do you have a few sentences that expand that a little to talk about the erratic nature of precipitation (spatial and temporal) in fancy statistical terms which means finding statistical significance in such a variable is a tough job?

I redid the number-of-event bar charts - all one color now.

I reran the Heatwave results to use 6-day periods as did NCA5. NCA5 was slick though. By choosing 1961-90 their background climatology was cold relative to decades before and since, so that results outside of those 30 years would produce a lot of days that exceed the percentile threshold. Then in the urban heatwave website, they started in 1960, assuring a result that more temps would exceed the threshold in post-1990 years.

That's all I've got.

Aloha

John C.

John Christy
Director, Earth System Science Center
Professor, Atmos and Earth Science

Alabama State Climatologist
The University of Alabama in Huntsville
climateman60@gmail.com

<8 Extreme weather.JRC.May07.docx>

On May 6, 2025, at 5:25 PM, John Christy <climateman60@gmail.com> wrote:

Steve

As Ross notes, the precipitation section is simply an update of our 2019 paper. The daily temperature work from the supplemented USHCN records is in a paper under review (but I suspect will have difficulty being accepted). The data are from the official USHCN site with my supplements adding about 10 percent to cover the missing data. There are about 41 million temperatures here, so I added about 4 million. Thus the results are dominated by publically-available data for others to check. If I get the go-ahead for publication, I can put the complete datafiles on a server for anyone to use.

John C.

John Christy
Director, Earth System Science Center
Professor, Atmos and Earth Science
Alabama State Climatologist
The University of Alabama in Huntsville
climateman60@gmail.com

On May 6, 2025, at 3:48 PM, Ross McKittrick
<ross.mckittrick@gmail.com> wrote:

The underlying analysis is all from John's and my Journal of Hydrology paper for which the code is already online. John updated and expanded the data sets and I ran them through the same code.

On Tue, May 6, 2025 at 4:41 PM Steven Koonin
<steven.koonin@gmail.com> wrote:

A thought about the analysis of long-term CONUS data, which as John knows I like very much:

If some (most?) of this work is as-yet unpublished in a peer-reviewed journal, I'd think you've got to make the expanded and in-filled database (and analysis code?) publicly accessible so that the work can be checked. Are you prepared to do that (at least for the "final", if not the "draft")?

SEK

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Tuesday, May 6, 2025 1:05 PM
To: John Christy <climateman60@gmail.com>
Cc: Judith Curry <curry.judith@gmail.com>; Roy W. Spencer <roywspencer@hotmail.com>; Travis Fisher <travis.scott.fisher@gmail.com>; Steven Koonin <steven.koonin@googlemail.com>
Subject: Re: Ch 8 Extreme weather

Here's another edition of Ch 8 on extremes taking Judy's and John's points into account. The IPCC/NCA review is ported to an Appendix. I added a paragraph about LTP to the intro. I've incorporated the NE data. The bulk of the chapter is John's US temperature and precip data analysis with some additional material on wildfires, drought and TCs. For the US temperature extremes I moved the range figure to come just after the hot/cold extremes count because I think they both make a coherent point namely that the US becoming less extreme because heat events are rising more slowly than cold events are warming. There will still need to be an overall chapter summary but I'm going to move on to other chapters now. I'll put this in the Dropbox.

On Tue, May 6, 2025 at 12:12 PM John Christy <climateman60@gmail.com> wrote:

All

We can fill in without much trouble the change, or lack thereof, of "high impact" weather events (mainly wind and flooding like hurricanes, tornadoes etc) in the CONUS Extreme chapter. The temp and precip metrics are important because they are just long enough to essentially tell us Mother Nature has a mind of her own.

1. I'll look at Ross's version of 8 soon, but my hope was that Ross would translate the statistical remarks into the correct language(eg are their more precise words that chaotic, episodic, stochastic, etc).
2. We have analyses of the CONUS with more complete datasets than the NCA used (recall NCA analyses depended essentially on a small group at NCEI using datasets that (a)

were not very consistent in time, (b) could promote alarm and (c) already easy for them to get to. Our datasets have literally hundreds of thousands of manually keyed-in observations for temporal consistency. In other words, this lets people know we can address these issues from soup to nuts - we're not armchair climate scientists ... we can reproduce what they did, but make the analysis more comprehensive to address the questions.

John C.

Sent from my iPhone

On May 6, 2025, at 4:14 AM, Ross McKittrick

<ross.mckittrick@gmail.com> wrote:

Just read your attribution chapter. I hadn't picked up on the gap between IPCC Ch 11 vs Ch 12 language but i see your point that rather than promoting one set of attribution findings we should explain that even within the same AR different authors assessed things differently. One option is to leave Ch 6 where it is but reverse the order of Ch 7 and 8 while retitling the latter as you suggest. That way, having established the extreme weather types for which there are no detectable trends it will more naturally follow that they won't be discussed in the attribution chapter.

On Tue, May 6, 2025 at 9:45 AM Judith Curry
<curry.judith@gmail.com> wrote:

Ok, but the summary table (12.12 from AR6) is inconsistent with conclusions in AR6 ch 11 (I point this out in section 6.4 of my attribution chapter).

In my mind, the main point is disagreement and ambiguity I would suggest chapter 8 going before chapter 6, retitling it as US Weather and Climate Extremes, and make it more of a data chapter, not an attribution chapter (I would put table 12.12 in the attribution chapter if you want to include the actual table)

On Tue, May 6, 2025 at 6:39 AM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

I like john's material too which is why I included it and am working on the extended regional analyses. But on this topic people are heavily primed to believe that extreme weather of all kinds is getting worse due to climate change. The assertion shows up everywhere climate change is discussed including in academic papers. I think it is very valuable to show people that neither the IPCC nor the NCA support such this view. We should at least lead off with the summary table. The library of quotes can go to an appendix but I have found in my public speaking that they have considerable persuasive force for most people. We also need to address floods, droughts, hurricane landfalls, tornadoes etc. I don't see any reason to delete all that material.

On Tue, May 6, 2025 at 9:22 AM Judith Curry <curry.judith@gmail.com> wrote:

I actually liked John's chapter. I would have retitled it "US temperature and rainfall extremes"

I think this is more relevant than a complete rehash of everything the last 3 IPCC reports have stated for every conceivable severe weather event.

John's version meshes very well with my version of chapter 6. Read especially section 6.4 of my chapter. I note a discrepancy between chapters 11 and 12 in terms of detecting and attributing extreme events. This is the main point in my opinion, there is ambiguity in all this. Lets focus on pointing out the problems/challenges, and topics that are particularly relevant for the US

On Mon, May 5, 2025 at 7:47 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

I have wrangled the extreme weather chapter into something I hope flows better and is more readable. I adopted Judy's editorial changes and then the new changes I made are in red. I added some new Figures showing results on PC and SE data and will add the NE chart when I get the data. I also made some labeling suggestions including for the 3 rainfall charts that are hard to understand. Section summaries and an overall summary are still needed.
Ross

--

Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>


<~WRD0765.jpg>

--

Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

<~WRD0765.jpg>

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

From: Ross McKittrick [ross.mckittrick@gmail.com]
Sent: 7/23/2025 2:01:49 AM
To: Judith Curry [curry.judith@gmail.com]; Roy W. Spencer [roywspencer@hotmail.com]; John Christy [climatemanager60@gmail.com]; Steven Koonin [steven.koonin@gmail.com]; Travis Fisher [travis.scott.fisher@gmail.com]
Subject: Post review edition
Attachments: DoE.Report.July22.docx

All amendments added, figures renumbered, metadata figure numbers updated, pages laid out so figures are placed appropriately etc.

Steve - can you add section citations in the Executive Summary?

Travis - any chance of a hi-res logo?

Other than that it's done I think. Everyone please feel free to take one more look.

Cheers,

Ross



A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate

Climate Working Group

United States Department of Energy

July 22, 2025

A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate

Report to U.S. Energy Secretary Christopher Wright
July 22, 2025

Climate Working Group:

John Christy, Ph.D.

Judith Curry, Ph.D.

Steven Koonin, Ph.D.

Ross McKittrick, Ph.D.

Roy Spencer, Ph.D.

Copyright © 2025 United States

Suggested citation:

Climate Working Group (2025) A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate. Washington DC: Department of Energy, July 22, 2025

Table of Contents

SECRETARY’S FOREWORD	VII
EXECUTIVE SUMMARY	VIII
PREFACE.....	IX
PART I: DIRECT HUMAN INFLUENCE ON ECOSYSTEMS AND THE CLIMATE.....	1
1 CARBON DIOXIDE AS A POLLUTANT	2
2 DIRECT IMPACTS OF CO₂ ON THE ENVIRONMENT	3
2.1 CO ₂ AS A CONTRIBUTOR TO GLOBAL GREENING	3
2.1.1 <i>Measurement of global greening.....</i>	<i>3</i>
2.1.2 <i>Photosynthesis and CO₂ levels</i>	<i>4</i>
2.1.3 <i>Rising CO₂ and crop water use efficiency.....</i>	<i>6</i>
2.1.4 <i>CO₂ fertilization benefits in IPCC Reports.....</i>	<i>6</i>
2.2 THE ALKALINE OCEANS	6
2.2.1 <i>Changing pH</i>	<i>6</i>
2.2.2 <i>Coral reef changes.....</i>	<i>7</i>
3 HUMAN INFLUENCES ON THE CLIMATE.....	11
3.1 COMPONENTS OF RADIATIVE FORCING AND THEIR HISTORY.....	11
3.1.1 <i>Historical radiative forcing.....</i>	<i>11</i>
3.1.2 <i>Change in atmospheric CO₂ since 1958</i>	<i>13</i>
3.2 FUTURE EMISSION SCENARIOS AND THE CARBON CYCLE	14
3.2.1 <i>Emission scenarios</i>	<i>14</i>
3.2.2 <i>The carbon cycle relating emissions and concentrations.....</i>	<i>17</i>
3.3 URBANIZATION INFLUENCE ON TEMPERATURE TRENDS	20
PART II: CLIMATE RESPONSE TO CO₂ EMISSIONS.....	24
4 CLIMATE SENSITIVITY TO CO₂ FORCING.....	25
4.1 INTRODUCTION	25
4.2 MODEL-BASED ESTIMATES OF CLIMATE SENSITIVITY	26
4.3 DATA-DRIVEN ESTIMATES OF CLIMATE SENSITIVITY	27
4.4 TRANSIENT CLIMATE RESPONSE	29
5 DISCREPANCIES BETWEEN MODELS AND INSTRUMENTAL OBSERVATIONS.....	31
5.1 INTRODUCTION	31
5.2 SURFACE WARMING	33
5.3 TROPOSPHERIC WARMING	34
5.4 VERTICAL TEMPERATURE PROFILE MISMATCH.....	36

5.5	STRATOSPHERIC COOLING	38
5.6	SNOW COVER MISMATCH	39
5.7	HEMISPHERIC SYMMETRY OF THE PLANETARY ALBEDO	41
5.8	U.S. CORN BELT	42
6	EXTREME WEATHER	46
6.1	INTRODUCTION	46
6.2	HURRICANES AND TROPICAL CYCLONES	48
6.3	TEMPERATURE EXTREMES.....	52
6.3.1	<i>Temperatures in the U.S. are becoming less extreme</i>	<i>53</i>
6.3.2	<i>Exceedances of a heat threshold.....</i>	<i>56</i>
6.3.3	<i>Heatwaves.....</i>	<i>57</i>
6.4	EXTREME PRECIPITATION.....	61
6.5	TORNADOES.....	66
6.6	FLOODING	67
6.7	DROUGHTS.....	68
6.8	WILDFIRES.....	69
7	CHANGES IN SEA LEVEL	75
7.1	GLOBAL SEA LEVEL RISE	75
7.2	U.S. SEA LEVEL RISE	75
7.3	PROJECTED SEA LEVEL RISE	79
8	UNCERTAINTIES IN CLIMATE CHANGE ATTRIBUTION	82
8.1	INTRODUCTION	82
8.2	ATTRIBUTION METHODS.....	83
8.3	ATTRIBUTION OF GLOBAL WARMING	84
8.3.1	<i>Natural climate variability.....</i>	<i>84</i>
8.3.2	<i>Optimal fingerprinting.....</i>	<i>88</i>
8.3.3	<i>Time series methods.....</i>	<i>89</i>
8.4	DECLINING PLANETARY ALBEDO AND RECENT RECORD WARMTH	90
8.5	ATTRIBUTION OF CLIMATE IMPACT DRIVERS.....	92
8.6	EXTREME EVENT ATTRIBUTION (EEA)	95
8.6.1	<i>Case study – 2021 Western North America heat wave</i>	<i>96</i>
	PART III: IMPACTS ON ECOSYSTEMS AND SOCIETY.....	103
9	CLIMATE CHANGE AND U.S. AGRICULTURE	104
9.1	ECONOMETRIC ANALYSES	104
9.2	FIELD AND LABORATORY STUDIES OF CO ₂ ENRICHMENT.....	105

9.3	CROP MODELING META-ANALYSES	107
9.4	CO ₂ FERTILIZATION AND NUTRIENT LOSS	108
10	MANAGING RISKS OF EXTREME WEATHER.....	110
10.1	SOCIOECONOMIC CONTEXT	110
10.2	DATA CHALLENGES	111
10.3	MORTALITY FROM TEMPERATURE EXTREMES.....	111
10.3.1	<i>Heat and cold risks.....</i>	<i>111</i>
10.3.2	<i>Mortality risks and energy costs</i>	<i>113</i>
11	CLIMATE CHANGE, THE ECONOMY, AND THE SOCIAL COST OF CARBON	116
11.1	CLIMATE CHANGE AND ECONOMIC GROWTH	116
11.1.1	<i>Overview.....</i>	<i>116</i>
11.1.2	<i>Empirical analysis of climate change and economic growth</i>	<i>119</i>
11.2	MODELS OF THE SOCIAL COST OF CARBON	120
11.2.1	<i>Estimating the SCC.....</i>	<i>121</i>
11.2.2	<i>Variations in the SCC.....</i>	<i>122</i>
11.2.3	<i>Evidence for low SCC</i>	<i>123</i>
11.2.4	<i>Tipping points</i>	<i>123</i>
11.2.5	<i>Are there alternatives?.....</i>	<i>125</i>
12	GLOBAL CLIMATE IMPACTS OF U.S. EMISSIONS POLICIES.....	129
12.1	THE SCALE PROBLEM	129
12.2	CASE STUDY: U.S. MOTOR VEHICLE EMISSIONS	129
12.3	CONCLUDING THOUGHTS	130
	GLOSSARY.....	132
	METADATA FOR FIGURES AND TABLES	136
	ABOUT THE AUTHORS.....	140

SECRETARY'S FOREWORD

Energy, Integrity, and the Power of Human Potential

Over my lifetime, I've had the privilege of working as an energy entrepreneur across a range of fields—nuclear, geothermal, natural gas, and more—and I now serve as Secretary of Energy under President Donald Trump. But above all, I'm a physical scientist who sees modern energy as nothing short of miraculous. It powers every aspect of modern life, drives every industry, and has made America an energy powerhouse with the ability to fuel global progress.

The rise of human flourishing over the past two centuries is a story worth celebrating. Yet we are told—relentlessly—that the very energy systems that enabled this progress now pose an existential threat. Hydrocarbon-based fuels, the argument goes, must be rapidly abandoned or else we risk planetary ruin.

That view demands scrutiny. That's why I commissioned this report: to encourage a more thoughtful and science-based conversation about climate change and energy. With my technical background, I've reviewed reports from the Intergovernmental Panel on Climate Change, the U.S. government's assessments, and the academic literature. I've also engaged with many climate scientists, including the authors of this report.

What I've found is that media coverage often distorts the science. Many people—even well-meaning ones—walk away with a view of climate change that is exaggerated or incomplete. To provide clarity, I asked a diverse team of independent experts to summarize the current state of climate science, with a focus on how it relates to the United States.

I didn't select these authors because we always agree—far from it. In fact, they may not always agree with each other. But I chose them for their rigor, honesty, and willingness to elevate the debate. I exerted no control over their conclusions. What you'll read are their words, drawn from the best available data and scientific assessments.

I've reviewed the report carefully and challenged its findings. I believe it faithfully represents the state of climate science today. Still, many readers may be surprised by its conclusions—which differ in important ways from the mainstream narrative. That's a sign of how far the public conversation has drifted from the science itself.

To correct course, we need open, respectful, and informed debate. That's why I'm inviting public comment on this report. Honest scrutiny and scientific transparency should be at the heart of our policymaking.

Climate change is real, and it deserves attention. But it is not the greatest threat facing humanity. That distinction belongs to global energy poverty. As someone who values data, I know that improving the human condition depends on expanding access to reliable, affordable energy. Climate change is a challenge—not a catastrophe. But misguided policies based on fear rather than facts could truly endanger human well-being.

We stand at the threshold of a new era of energy leadership. If we empower innovation rather than restrain it, America can lead the world in providing cleaner, more abundant energy—lifting billions out of poverty, strengthening our economy, and improving our environment along the way.

EXECUTIVE SUMMARY

This report reviews scientific certainties and uncertainties in how anthropogenic carbon dioxide (CO₂) and other greenhouse gas emissions have affected, or will affect, the Nation's climate, extreme weather events, and selected metrics of societal well-being. Those emissions are increasing the concentration of CO₂ in the atmosphere through a complex and variable carbon cycle, where some portion of the additional CO₂ persists in the atmosphere for centuries.

Elevated concentrations of CO₂ directly enhance plant growth, globally contributing to “greening” the planet and increasing agricultural productivity. They also make the oceans less alkaline (lower the pH). That is possibly detrimental to coral reefs, although the recent rebound of the Great Barrier Reef suggests otherwise.

Carbon dioxide also acts as a greenhouse gas, exerting a warming influence on climate and weather. Climate change projections require scenarios of future emissions. There is evidence that scenarios widely-used in the impacts literature have overstated observed and likely future emission trends.

The world's several dozen global climate models offer little guidance on how much the climate responds to elevated CO₂, with the average surface warming under a doubling of the CO₂ concentration ranging from 1.8°C to 5.7°C. Data-driven methods yield a lower and narrower range. Global climate models generally run “hot” in their description of the climate of the past few decades – too much warming at the surface and too much amplification of warming in the lower- and mid-troposphere. The combination of overly sensitive models and implausible extreme scenarios for future emissions yields exaggerated projections of future warming.

Most extreme weather events in the U.S. do not show long-term trends. Claims of increased frequency or intensity of hurricanes, tornadoes, floods, and droughts are not supported by U.S. historical data. Additionally, forest management practices are often overlooked in assessing changes in wildfire activity. Global sea level has risen approximately 8 inches since 1900, but there are significant regional variations driven primarily by local land subsidence; U.S. tide gauge measurements in aggregate show no obvious acceleration in sea level rise beyond the historical average rate.

Attribution of climate change or extreme weather events to human CO₂ emissions is challenged by natural climate variability, data limitations, and inherent model deficiencies. Moreover, solar activity's contribution to the late 20th century warming might be underestimated.

Both models and experience suggest that CO₂-induced warming might be less damaging economically than commonly believed, and excessively aggressive mitigation policies could prove more detrimental than beneficial. Social Cost of Carbon estimates, which attempt to quantify the economic damage of CO₂ emissions, are highly sensitive to their underlying assumptions and so provide limited independent information.

PREFACE

This document originated in late March 2025 when Secretary Wright assembled an independent group to write a report on issues in climate science relevant for energy policymaking, including evidence and perspectives that challenge the mainstream consensus. We agreed to undertake the work on the condition that there would be no editorial oversight by the Secretary, the Department of Energy, or any other government personnel. This condition has been honored throughout the process and the writing team has worked with full independence.

The group began working in early April with a deadline of May 28. The short timeline and the technical nature of the material meant that we could not comprehensively review all topics. Rather, we chose to focus on topics that are treated by a serious, established academic literature; that are relevant to our charge; that are downplayed in, or absent from, recent assessment reports; and that are within our competence.

While the report is intended to be accessible to non-experts, we have omitted some introductory or explanatory material that can easily be accessed elsewhere. Nor have we attempted to survey the entire literature related to the topics covered. We have focused as much as possible on literature published since 2020 and referenced previous IPCC and NCA assessment reports. We have also used data through 2024 where possible.

The writing team is grateful to Secretary Wright for the opportunity to prepare this report and for his support of independent scientific assessment and open scientific debate. We are also grateful to a team of anonymous peer-reviewers whose input helped improve the final report.

John Christy, Ph.D.

Judith Curry, Ph.D.

Steven Koonin, Ph.D.

Ross McKittrick, Ph.D.

Roy Spencer, Ph.D.

**PART I: DIRECT HUMAN INFLUENCE ON ECOSYSTEMS AND THE
CLIMATE**

1 CARBON DIOXIDE AS A POLLUTANT

Chapter summary:

Carbon dioxide (CO₂) differs in many ways from the so-called Criteria Air Pollutants. It does not affect local air quality and has no human toxicological implications at ambient levels. It is an issue of concern because of its effects on the global climate.

The Clean Air Act of 1970 defined six so-called Criteria Air Contaminants subject to regulation (EPA): particulate matter, ground-level ozone, sulfur dioxide, nitrogen dioxide, lead, and carbon monoxide. In 2007, the Supreme Court ruled that greenhouse gases (CO₂ among them) were also “pollutants” subject to regulation under Clean Air Act (*Mass. v. EPA*, 2007). While the definition of “pollutant” is ultimately a legal matter, there are important scientific distinctions between CO₂ and the Criteria Air Contaminants. The latter are subject to regulatory control because they cause local problems depending on concentrations that include nuisances (odor, visibility), damage to plants, and, at high enough exposure levels, toxicological effects in humans. In contrast, CO₂ is odorless, does not affect visibility and has no toxicological effects at ambient levels. It is a naturally occurring part of the atmosphere and a key component of human and plant respiration. CO₂ is essential for plant photosynthesis and higher levels are beneficial for vegetation. In these aspects, CO₂ is similar to water vapor.

Ambient outdoor air today contains about 430 parts per million (ppm) CO₂, increasing at about 2 ppm per year. The U.S. Occupational Safety and Health Administration issues guidelines for indoor workplaces in which elevated CO₂ might be encountered, such as where dry ice is used. The Permissible Exposure Limit is 5,000 ppm over 8 hours (OSHA, 2024). Allen *et al.* (2015) reported evidence of diminished performance on some cognitive tasks among workers in office cubicles when exposed to CO₂ levels above 1,000-1,500 ppm. These levels are far larger than any plausible ambient outdoor value through the end of the 22nd century.

The growing amount of CO₂ in the atmosphere directly influences the earth system by promoting plant growth (global greening), thereby enhancing agricultural yields, and by neutralizing ocean alkalinity. But the primary concern about CO₂ is its role as a greenhouse gas (GHG) that alters the earth’s energy balance, warming the planet. How the climate will respond to that influence is a complex question that will occupy much of this report.

References

- Allen, J., Macnaughton, P., Satish, U., *et al.* (2015). Associations of cognitive function scores with carbon dioxide, ventilation, and volatile organic compound exposures in office workers: A controlled exposure study of green and conventional office environments. *Environmental Health Perspectives*. 124. <https://doi.org/10.1289/ehp.1510037>
- Massachusetts v. Environmental Protection Agency, 549 U.S. 497 (2007). <https://www.oyez.org/cases/2006/05-1120>
- U.S. Environmental Protection Agency. (n.d.). Criteria air pollutants. <https://www.epa.gov/criteria-air-pollutants>
- U.S. Occupational Safety and Health Administration. (2024). OSHA occupational chemical database: Carbon dioxide. <https://www.osha.gov/chemicaldata/183>

2 DIRECT IMPACTS OF CO₂ ON THE ENVIRONMENT

Chapter summary:

CO₂ enhances photosynthesis and improves plant water use efficiency, thereby promoting plant growth. Global greening due in part to increased CO₂ levels in the atmosphere is well-established on all continents.

CO₂ absorption in sea water makes the oceans less alkaline. The recent decline in pH is within the range of natural variability on millennial time scales. Most ocean life evolved when the oceans were mildly acidic. Decreasing pH might adversely affect corals, although the Australian Great Barrier Reef has shown considerable growth in recent years.

2.1 CO₂ as a contributor to global greening

The growing CO₂ concentration in the atmosphere has the important positive effect of promoting plant growth by enhancing photosynthesis and improving water use efficiency. That is evident in the “global greening” phenomenon discussed below, as well as in the improving agricultural yields discussed in Chapter 10. Here we focus just on CO₂ fertilization; research on combined effects due to temperature and precipitation changes are discussed in Chapter 10.

2.1.1 Measurement of global greening

“Greening” refers to an increase in the fraction of the Earth’s surface covered by plants. It can be quantified by the “Leaf Area Index” (LAI) measured by satellite. Many studies over the past decade have confirmed a global greening pattern (increase in LAI) attributable in part to rising CO₂ levels. Zhu *et al.* (2016) was one of the first studies to report that global greening was detectable using satellite sensors. From 1982 to 2011 they detected greening over 25-50 percent of the Earth versus “browning” over only four percent and attributed 70 percent of the greening to rising CO₂ levels (see Figure 2.1). Other contributors included land-use changes, warming and nitrogen. The fraction attributable to CO₂ was largest in the tropics; other factors played more dominant roles in CONUS.

Zeng *et al.* (2017) confirmed the pattern of greening, noting that over thirty years it had added 8 percent to global leaf area and that greening was mitigating warming. Greening has been observed globally. Chen *et al.* (2019) show that in China and India much of it is driven by land management changes. Thus, while China accounts for only 6.6 percent of global vegetated area it accounts for 25 percent of global net increase in LAI. Piao *et al.* (2020) noted that greening was even observable in the Arctic. CO₂ fertilization effects are influenced by local temperature and nutrient and water availability, all of which vary regionally.

While plant models predict increased photosynthesis in response to rising CO₂, Haverd *et al.* (2020) reported a CO₂ fertilization rate much larger than model predictions. That is, CO₂ fertilization had driven an increase in observed global photosynthesis by 30 percent since 1900, versus 17 percent predicted by plant models. If true it would indicate that global models of the socioeconomic impacts of rising CO₂ have understated the benefits to crops and agriculture. Keenan *et al.* (2023), however, estimated a lower fertilization rate more in line with models. The connection between CO₂ fertilization and agriculture will be discussed in Chapter 9.




Figure 2.1: Trends in average Leaf Area Index (LAI). Source: Zhu *et al.* 2016 Figure 3.

Piao *et al.* (2020) and Chen *et al.* (2024) report that the greening trend continues with no evidence of slowdown, and CO₂ fertilization remains the dominant driver

2.1.2 Photosynthesis and CO₂ levels

Plants build biomass through *photosynthesis*, a process that converts carbon dioxide, water, and light into sugar. The plant enzyme responsible for photosynthesis is Ribulose-1,5-bisphosphate-carboxylase/oxygenase or “Rubisco”. Photosynthesis is initiated when CO₂ is available at the surface of the Rubisco enzyme where it is converted to a molecule with 3 carbon atoms and thereafter incorporated into plant mass. This is referred to as the “C3” process.

Rubisco is estimated to have evolved about 3 billion years ago. Over geological time the Earth’s atmospheric CO₂ levels were usually many times higher than they are today. About 400 million years ago CO₂ levels were an estimated 2,000-4,000 ppm and were at or above 1,000 ppm for much of the interval from 200 to 50 million years ago (Berner 2006, Judd *et al.* 2024). Over the past 35 million years the level of atmospheric CO₂ has been steadily declining, falling to as low as 170 ppm during glaciations (Gerhart and Ward 2010). While the modern rate of change in CO₂ may be high compared to prior intervals, the geological evidence is that plants and animals evolved under much higher CO₂ levels than at present.

In response to low-CO₂ conditions some plants evolved another photosynthetic pathway called C4, in which CO₂ is concentrated in the vicinity of Rubisco, allowing for the C3 process to function more efficiently. For agricultural purposes the plant categories are:

- C3: rice, wheat, soybeans and most other crops
- C4: maize (corn), sugar cane, millet, sorghum

Had atmospheric CO₂ levels continued declining, plant growth would have declined and eventually ceased. Below 180 ppm, the growth rates of many C3 species are reduced 40-60 percent relative to 350 ppm (Gerhart and Ward 2010) and growth has stopped altogether under experimental conditions of 60—140 ppm CO₂. Some C4 plants are still able to grow at levels even as low as 10 ppm, albeit very slowly (Gerhart and Ward 2010).




Figure 2.2: growth of *Abutilon theophrasti* after 14 days under identical conditions but for the indicated variations in CO₂ levels. Source: Gerhart and Ward (2010). Note “Current” corresponds to 1988 in image.

Current CO₂ levels are about 430 ppm, up from 280 ppm in the early 1800s. The positive response of plants to extra CO₂ is illustrated in Figure 2.2, reproduced from Gerhart and Ward (2010). It shows the growth effect of CO₂ on Velvetleaf (*Abutilon theophrasti*) seedlings over 14 days under controlled conditions where only the CO₂ exposure is varied. The gains induced by increasing CO₂ from 150 ppm to 350 ppm continue under a further doubling to 700 ppm.

Over the past 60+ years there have been thousands of studies on the response of plants to rising CO₂ levels. The overwhelming theme is that plants, especially C3 plants, benefit from extra CO₂. There are two mechanisms by which CO₂ confers a growth benefit:

- Enhanced photosynthesis via the metabolic pathways described above.
- Increased water use efficiency. This arises because plants draw in CO₂ by opening the stomata (pores) on the leaf surface. When CO₂ is scarce the stomata must be kept wide open for long periods, allowing water to evaporate. Under enriched CO₂ conditions the stomata remain closed for longer periods, thus helping the plant retain water longer, and so increasing water use efficiency.

Specific effects of climate change on U.S. agriculture will be reviewed in Chapter 9.

2.1.3 *Rising CO₂ and crop water use efficiency*

Deryng *et al.* (2016) surveyed evidence on crop water productivity (CWP), the yield per unit of water used, drawing attention to the potential for CO₂ both to enhance photosynthesis and to reduce leaf-level transpiration (water loss during leaf respiration). They surveyed all available FACE data (Free Air CO₂ Enrichment—see Chapter 9) on crop yield changes for maize (corn), wheat, rice, and soybean and combined it with crop model data simulating yield responses as of 2080 under the extreme RCP8.5 emissions scenario in four growing regions (Tropics, Arid, Temperate and Cold) each of which were split into rainfed and irrigated sub-regions. They reported that models without CO₂ fertilization predicted CWP losses in every region, but those were more than offset by CO₂ fertilization so that all regions showed a net CWP gain. Deryng *et al.* (2016) also reported that negative impacts of warming on wheat and soybean yields were fully offset by CWP gains and mitigated by up to 90 percent for rice and 60 percent for maize.

Similarly, Cheng *et al.* (2017) noted that increased Gross Primary Production from 1982 to 2011 due to rising CO₂ uptake was accompanied by such large gains in CWP that global water use by plants had not increased, despite the extra biomass.

Deryng *et al.* (2016) assumed that climate change would “exacerbate water scarcity”. Yet while models do predict that drylands will expand under climate warming, current data show the opposite: greening is happening even in arid areas. Zhang *et al.* (2024) report that due to increased CO₂ levels “increasing aridity in drylands won’t lead to a general loss of vegetation productivity”; at most only 4 percent of currently arid areas will see increased desertification.

2.1.4 *CO₂ fertilization benefits in IPCC Reports*

The IPCC has only minimally discussed global greening and CO₂ fertilization of agricultural crops. The topic is briefly acknowledged in a few places in the body of the IPCC 6th and earlier Assessment Reports but is omitted in all Summary documents. Section 2.3.4.3.3 of the AR6 Working Group I report, entitled “global greening and browning,” points out that the IPCC Special Report on Climate Change and Land had concluded with *high confidence* that greening had increased globally over the past 2-3 decades. It then discusses that there are variations in the greening trend among data sets, concluding that while they have *high confidence* greening has occurred, they have *low confidence* in the magnitude of the trend. There are also brief mentions of CO₂ fertilization effects and improvements in water use efficiency in a few other chapters in the AR6 Working Groups I and II Reports.

Overall, however, the Policymaker Summaries, Technical Summaries, and Synthesis Reports of AR5 and AR6 do not discuss the topic.

2.2 The Alkaline Oceans

2.2.1 *Changing pH*

A neutral aqueous solution has a pH of 7.0, while one with pH greater than 7.0 is alkaline (also termed basic) and with pH less than 7.0 is acidic. The modern-day global average pH of sea water is estimated to be 8.04 (Copernicus Marine Service 2025, Figure 2.3), down from an estimated value of 8.2 in pre-Industrial times (Gattuso and Hansson, 2011). As CO₂ concentrations in the atmosphere increased, the oceans absorbed more, which decreases their pH. Depending upon the oceans’ buffering capacity, they are expected to become somewhat less alkaline over time, consistent with the observed decrease in pH.




Figure 2.3: Ocean pH 1985 – 2022. Source: Copernicus Marine Service 2025

While this process is often called “ocean acidification”, it is a misnomer because the oceans are not expected to become acidic; “ocean neutralization” would be more accurate. Even if the water were to turn acidic, it is believed that life in the oceans evolved when the oceans were mildly acidic with pH 6.5 to 7.0 (Krissansen-Totton *et al.*, 2018). On the time scale of thousands of years, boron isotope proxy measurements show that ocean pH was around 7.4 or 7.5 during the last glaciation (up to about 20,000 years ago) increasing to present-day values as the world warmed during deglaciation (Rae *et al.*, 2018). Thus, ocean biota appear to be resilient to natural long-term changes in ocean pH since marine organisms were exposed to wide ranges in pH.

2.2.2 Coral reef changes

There are concerns that a decreasing pH of sea water will reduce the calcification rate of coral reefs. But coral reefs already endure large swings in pH, partly due to daily photosynthetic activity in the reef; measured pH values range from 9.4 during the day to 7.5 at night (Revelle and Fairbridge, 1957). De’ath *et al.* (2009) reported that a portion of Australia’s Great Barrier Reef (GBR, the world’s largest coral reef ecosystem) had experienced a 14 percent decline in calcification since 1990. This was tentatively attributed to increasing water temperature and decreasing pH. But Ridd *et al.* (2013) showed that report to have resulted from a biased data analysis that, when corrected, showed no change in calcification rates. Nevertheless, the alarm produced by the original paper has persisted as evidenced by the large number of published citations (541) to the original study compared to only 11 citations to the correction (as of 30 April 2025).

The most recent annual summary of GBR conditions from the Australian Institute of Marine Science indicates that coral production has rebounded strongly (AIMS, 2023). Figure 2.4 shows the results of the AIMS surveys of hard coral cover, expressed as a percentage of the reef area. Much of the decline in the GBR before 2011 turned out to be due to intense tropical cyclone activity (Beeden *et al.*, 2015) as well as a string of marine heatwaves, agricultural runoff and invasive species (Woods Hole, 2023). Given the reported declines in GBR calcification between 1990 and 2009 and the continued increase in atmospheric CO₂ levels the rebound has surprised some observers.




Figure 2.4 Hard coral cover of three regions of the Great Barrier Reef 1985 to 2023. Source: AIMS 2023.

It is being increasingly recognized that publication bias (alarming ocean acidification results preferred by high-impact research publications) exaggerates the reported impacts of declining ocean pH. An ICES Journal of Marine Science Special Issue addressed this problem with an article entitled, *Towards a Broader Perspective on Ocean Acidification Research*. In the Introduction to that Special Issue, H. I. Browman stated, “As is true across all of science, studies that report no effect of ocean acidification are typically more difficult to publish.” (Browman, 2016).

Similarly, a meta-analysis (Clements *et al.*, 2021) of the negative effects of ocean acidification on reef fish behavior found what they called a “decline effect”: initially dramatic conclusions published in prominent journals showing apparently large impacts of acidification tended to be followed up by subsequent studies on larger sample sizes yielding much smaller and typically non-existent effects. They call for their colleagues to improve research practices to counter the “decline effect”:

[The] vast majority of studies with large effect sizes in this field tend to be characterized by low sample sizes, yet are published in high-impact journals and have a disproportionate influence on the field in terms of citations. We contend that ocean acidification has a negligible direct impact on fish behavior, and we advocate for improved approaches to minimize the potential for a decline effect in future avenues of research (Clements *et al.*, 2021).

In summary, ocean life is complex and much of it evolved when the oceans were acidic relative to the present. The ancestors of modern coral first appeared about 245 million ago; as can be seen from Figure 2.2, CO₂ levels for more than 200 million years afterward were many times higher than they are today. Much of the public discussion of the effects of ocean “acidification” on marine biota has been one-sided and exaggerated.

References

- AR6: *Intergovernmental Panel on Climate Change Sixth Assessment Report* (2021) Working Group I Contribution. www.ipcc.ch.
- Australian Institute of Marine Science. (2022). Continued coral recovery leads to 36-year highs across two-thirds of the Great Barrier Reef. https://www.aims.gov.au/sites/default/files/2022-08/AIMS_LTMP_Report_on%20GBR_coral_status_2021_2022_040822F3.pdf
- Beeden, R., Maynard, J., Puotinen, M., Marshall, P., Dryden, J., Goldberg, J., and Williams, G. (2015). Impacts and recovery from Severe Tropical Cyclone Yasi on the Great Barrier Reef. *PLOS ONE*, 10, e0121272. <https://doi.org/10.1371/journal.pone.0121272>

- Berner, R. A. (2006). GEOCARBSULF: A combined model for Phanerozoic atmospheric O₂ and CO₂. *Geochimica et Cosmochimica Acta*, 70, 5653–5664.
- Browman, H. I. (2016). Applying organized scepticism to ocean acidification research. *ICES Journal of Marine Science*, 73(3), 529.1–536. <https://doi.org/10.1093/icesjms/fsw010>
- Chen, C., Park, T., Wang, X., Piao, S., Xu, B., Chaturvedi, R. K., and Myneni, R. B. (2019). China and India lead in greening of the world through land-use management. *Nature Sustainability*, 2, 122–129. <https://www.nature.com/articles/s41893-019-0220-7>
- Chen, X., Wang, Y., Liu, Y., and Piao, S. (2024). The global greening continues despite increased drought stress since 2000. *Global Ecology and Conservation*, 49, e02791. <https://www.sciencedirect.com/science/article/pii/S2351989423004262>
- Cheng, L., Zhang, L., Wang, Y. P., et al. (2017). Recent increases in terrestrial carbon uptake at little cost to the water cycle. *Nature Communications*, 8, 110. <https://doi.org/10.1038/s41467-017-00114-5>
- Clements, J. C., Sundin, J., Clark, T. D., and Jutfelt, F. (2022). Meta-analysis reveals an extreme “decline effect” in the impacts of ocean acidification on fish behavior. *PLOS Biology*, 20(2), e3001511. <https://doi.org/10.1371/journal.pbio.3001511>
- Copernicus Marine Service. (2025). Global ocean acidification – Mean sea water pH time series and trend from multi-observations reprocessing. https://data.marine.copernicus.eu/product/GLOBAL_OMI_HEALTH_carbon_ph_area_averaged/description
- De'ath, G., Lough, J., and Fabricius, K. (2009). Declining coral calcification on the Great Barrier Reef. *Science*, 323, 116–119. <https://doi.org/10.1126/science.1165283>
- Deryng, D., Conway, D., Ramankutty, N., Price, J., Warren, R., Jones, R., ... and Elliott, J. (2016). Regional disparities in the beneficial effects of rising CO₂ concentrations on crop water productivity. *Nature Climate Change*. <https://doi.org/10.1038/nclimate2995>
- Gattuso, J. P., and Hansson, L. (Eds.). (2011). *Ocean acidification: Background and history*. Oxford University Press.
- Gerhart, L. M., and Ward, J. K. (2010). Plant responses to low [CO₂] of the past. *New Phytologist*, 188, 674–695. <https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-8137.2010.03441.x>
- Haverd, V., B. Smith, J. G. Canadell, et al. (2020). Higher than expected CO₂ fertilization inferred from leaf to global observations. *Global Change Biology*, 26, 2390–2402. <https://doi.org/10.1111/gcb.14950>
- Keenan, T. F., X. Luo, B. D. Stocker, et al. (2023). A constraint on historic growth in global photosynthesis due to rising CO₂. *Nature Climate Change* 13(12): 1376-1381 DOI: 10.1038/s41558-023-01867-2.
- Judd, E. J., Scotese, C. R., Young, S. A., et al. (2024). A 485-million-year history of Earth's surface temperature. *Science*, 385(6715). <https://doi.org/10.1126/science.adk3705>
- Krissansen-Totton, J., Arney, G. N., and Catling, D. C. (2018). Constraining the climate and ocean pH of the early Earth with a geological carbon cycle model. *Proceedings of the National Academy of Sciences*, 115(6), 4105–4110. <https://doi.org/10.1073/pnas.1721296115>
- Piao, S., X. Wang, T. Park, et al. (2020). Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth & Environment* 1(1): 14-27 DOI: 10.1038/s43017-019-0001-x
- Rae, J. W. B., Burke, A., Robinson, L. F., et al. (2018). CO₂ storage and release in the deep Southern Ocean on millennial to centennial timescales. *Nature*, 562, 569–573. <https://doi.org/10.1038/s41586-018-0614-0>
- Revelle, R., and Fairbridge, R. W. (1957). Carbonate and carbon dioxide. In J. W. Hedgpeth (Ed.), *Treatise on marine ecology and paleoecology* (Vol. 1). *Geological Society of America*.
- Ridd, P., Silva, E., and Stieglitz, T. (2013). Have coral calcification rates slowed in the last twenty years? *Marine Geology*, 346, 392–399. <https://doi.org/10.1016/j.margeo.2013.09.002>
- Woods Hole Oceanographic Institution. (2023). Is the Great Barrier Reef making a comeback? <https://www.whoi.edu/oceanus/feature/is-the-great-barrier-reef-making-a-comeback/>

- Zeng, Z., Piao, S., Li, L., *et al.* (2017). Climate mitigation from vegetation biophysical feedbacks during the past three decades. *Nature Climate Change*. <https://doi.org/10.1038/nclimate3299>
- Zhang, Y., Liu, Y., Chen, X., *et al.* (2024). Less than 4% of dryland areas are projected to desertify despite increased aridity under climate change. *Nature Communications Earth and Environment*, 5. <https://www.nature.com/articles/s43247-024-01463-y>
- Zhu, Z., Piao, S., Myneni, R. B., *et al.* (2016). Greening of the Earth and its drivers. *Nature Climate Change*, 6, 791–795. <https://www.nature.com/articles/nclimate3004>

3 HUMAN INFLUENCES ON THE CLIMATE

Chapter Summary:

The global climate is naturally variable on all time scales. Anthropogenic CO₂ emissions add to that variability by changing the total radiative energy balance in the atmosphere.

The IPCC has downplayed the role of the sun in climate change but there are plausible solar irradiance reconstructions that imply it contributed to recent warming.

Climate projections are based on IPCC emission scenarios that have tended to exceed observed trends. Most academic climate impact studies in recent years are based upon the extreme RCP 8.5 scenario that is now considered implausible; its use as a business-as-usual scenario has been misleading.

Carbon cycle models connect annual emissions to growth in the atmospheric CO₂ stock. While models disagree over the rate of land and ocean CO₂ uptake, all agree that it has been increasing since 1959.

3.1 Components of radiative forcing and their history

3.1.1 Historical radiative forcing

A changing climate has been the norm throughout the Earth's 4.6-billion-year history. The Earth's temperature and weather patterns change naturally over timescales ranging from decades to millions of years. Natural variations in the surface climate originate in two ways. Internal climate fluctuations associated with circulations in the atmosphere and ocean exchange energy, water, and carbon between the atmosphere, oceans, land, and ice. External influences on the climate system include variations in the energy received from the sun and the effects of volcanic eruptions. Human activities influence climate through changing land use and land cover. Humans are also changing the composition of the atmosphere by emissions of CO₂ and other greenhouse gases and by altering the concentration of aerosol particles in the atmosphere.

The earth is warmed by the sunlight it absorbs and is cooled by the heat it radiates to space. Averaged over the Earth's surface, each of these processes involve power flows of about 240 Watts per square meter (W/m²). When they are in balance, there are no net external causes of warming or cooling. Both human and natural influences on the climate alter this balance and so cause the climate to change.

Influences on the Earth's energy balance at the top of the atmosphere are quantified by "radiative forcing", the extent to which they disturb the warming/cooling balance; a positive forcing warms while a negative forcing cools. The IPCC's estimated history of major components of radiative forcing since 1750 is shown in the following two figures from its AR6.




Figure 3.1.1: IPCC estimates of radiative forcing components over time. Shading indicates uncertainty ranges. Source: AR6 WGI Ch2 Fig. 10




Figure 3.1.2 IPCC estimates of radiative forcing component changes from 1750 to 2019. Source: AR6 WGI Ch 7 Fig. 7-6.

These graphs show that the total radiative forcing is comprised of both natural and anthropogenic components. Carbon dioxide is the largest human influence on the climate and the one most relevant to the influence of fossil fuel use. It exerts a warming influence by decreasing the cooling power of the atmosphere. Emissions of CO₂ are accumulating in the atmosphere, as described in the following section, so that the warming influence is growing. Other greenhouse gases (methane, nitrous oxide, halogens, and ozone) act similarly, currently adding another 75 percent to CO₂'s warming. Aerosols exert an overall cooling effect, although with large uncertainties in the way they catalyze the formation of reflective clouds. As a result, understanding the causes of recent warming requires not just identifying the warming effects of CO₂, but also the more uncertain cooling effects of aerosols.

The IPCC assesses the change in the radiative forcing by the sun to be negligible, based on their preference for data reconstructions that imply minimal solar change since preindustrial times. But Connolly *et al.* (2021) reviewed sixteen different Total Solar Irradiance (TSI) reconstructions in the literature

covering the years 1600-2000; the reconstructions vary from almost no change in TSI to a relatively large upward trend. Those authors note that the variation in TSI reconstructions combined with variations in surface temperature reconstructions allows for inferences consistent with either no or most 20th century warming being attributable to the sun.

A particularly thorny issue is the gap in TSI data between 1989 and 1991 due to a delay in the launch of a monitor following the Space Shuttle Challenger disaster on January 28 1986. This delay prevented a replacement satellite from being launched in time to overlap with, and its readings to be intercalibrated with, the prior system (Zacharias 2014, Scafetta *et al.* 2019). This is called the ACRIM (Active Cavity Radiometer Irradiance Monitor) gap problem. The question of whether there is an upward trend in TSI over 1978 to 2018 hangs on how the ACRIM data gap is filled. Connolly *et al.* (2021) found that the IPCC's consensus statements on solar forcing were formulated prematurely through the suppression of dissenting scientific opinions.

Another natural radiative forcing component is volcanic aerosols, which exert an episodic cooling influence. Box 4.1 in the IPCC AR6 Report addresses the climate impact of volcanic eruptions, noting three explosive volcanic eruptions that occurred in the first half of the 19th century. This included the 1815 Tambora eruption that resulted in the 'year without summer', with multiple harvest failures across the Northern Hemisphere. There is uncertainty about the sign of the relatively small forcing due to the submarine volcano Hunga Tonga which erupted in 2022 (Jenkins *et al.* 2023, Schoeberl *et al.* 2024).

Figure 3.1.1 shows that the anthropogenic forcing component was negligible before about 1900 and has increased steadily since, rising to almost 3 W/m² today. However, this is still only about 1 percent of the unperturbed radiation flows, making it a challenge to isolate the effects of anthropogenic forcing; state-of-the-art satellite estimates of global radiative energy flows are only accurate to a few W/m².

Natural sources of global energy imbalance other than volcanoes and total solar irradiance (TSI) are not included in these graphs because they remain largely unknown.

3.1.2 Change in atmospheric CO₂ since 1958

Carbon dioxide's warming influence depends on how much "extra" CO₂ accumulates in the atmosphere- *i.e.*, its concentration above the preindustrial value of 280 ppm. The CO₂ level as recorded at the Mauna Loa observatory in Hawaii, generally used as the representative global average concentration, is available online at <https://gml.noaa.gov/ccgg/trends/index.html>. The concentration was about 316 ppm at the start of the record in 1959 and is now about 430 ppm, a 36 percent increase. At the end of the last glaciation CO₂ levels had fallen to about 180 ppm. As discussed in Chapter 2, C3 plants begin dying at CO₂ levels below about 140 ppm and C4 plants at levels below 100 ppm, so if CO₂ levels had continued falling plant life would have been imperiled.




Fig. 3.1.3. Yearly average atmospheric CO₂ concentrations (1959-2025) in ppm measured at Mauna Loa (blue). C3 Threshold: Level below which C3 plants begin dying (140 ppm, see Chapter 2). C4 Threshold: Level below which C4 plants begin dying (100 ppm, see Chapter 2). Glacial minimum: Minimum level during recent glaciations (purple arrow). CO₂ data source: <https://gml.noaa.gov/ccgg/trends/index.html>

The annual increase in concentration is only about half of the CO₂ emitted because land and ocean processes currently absorb “excess” CO₂ at a rate approximately 50 percent of the human emissions. Future concentrations, and hence future human influences on the climate, therefore depend upon two components: (1) future rates of global human CO₂ emissions, and (2) how fast the land and ocean remove extra CO₂ from the atmosphere. We discuss each of these in turn.

3.2 Future emission scenarios and the carbon cycle

3.2.1 Emission scenarios

Assessing the dangers of future GHG emissions requires assumptions about what those emissions will be. Future emissions, and hence human influences on the climate, will depend upon future demographics, economic activity, regulation, and energy and agricultural technologies. Various assumptions about each of those lead to projections of greenhouse gas emissions and concentrations, aerosol concentrations, and changes in land use, which ultimately can be combined into assumptions about anthropogenic radiative forcing.

The great uncertainties about these many factors make it impossible to precisely predict future emissions. Instead, the IPCC has used various sets of scenarios meant to span a plausible range of possibilities for population, economy, and technologies. Recent versions of the scenarios are labeled by a number indicating the anthropogenic radiative forcing expected in 2100 under that scenario. Thus, a scenario labeled with a “6” corresponds to 6 W/m² of human-induced radiative forcing (warming) at the end of the century. (Recall current anthropogenic radiative forcing is about 2.7 W/m².)

Although the IPCC does not claim its emission scenarios are forecasts, they are often treated as such. Comparisons of past scenario groups against observations show that IPCC emission projections have tended to overstate actual subsequent emissions. For the IPCC Third and Fourth Assessment Reports a set of emission projections from the *Special Report on Emission Scenarios* was used; these were referred to as the SRES scenarios. McKittrick *et al.* (2012) showed that, when converted to per capita values, the SRES scenario emissions distribution was skewed upwards compared with observed trends. The bias of the SRES scenarios was confirmed by the later analysis of Hausfather *et al.* (2019) who showed that observed atmospheric CO₂ concentrations tracked the low end of the SRES range and also of subsequent IPCC scenario ranges (Figure 3.2.1).

For AR5 the IPCC developed a new set of scenarios called the Representative Concentration Pathways (RCPs). These were identified by a number representing the increase in forcing and were thus called RCP2.6, RCP4.5, RCP6.0 and RCP8.5. RCP2.6 (implying an anthropogenic radiative forcing in 2100 of 2.6 W/m²) describes a GHG concentration pathway leading to warming well below 2°C. At the other end of the scale RCP8.5 is an extreme outcome implying nearly 5°C warming from 1900 to 2100.

RCP8.5 came to be referred to as the no-policy baseline, or “business-as-usual” scenario in both the academic literature and popular media. It was therefore used to generate the reference outcome supposedly representing the 21st century world in the absence of increasingly stringent emission reduction policies. But RCP8.5 was intended as a low-probability high emissions scenario and its use as a business-as-usual baseline has been criticized as grossly misleading.¹ Hausfather and Peters (2020a) writing in a commentary in *Nature*, pointed out that RCP8.5 was developed as an extreme worst-case, and its misuse as a “business as usual” baseline has resulted in a large number of misleading studies and media reporting.

The implausibility of the RCP8.5 scenario was examined by Burgess *et al.* (2021). The implausibility of RCP8.5 should not be interpreted as *very unlikely* (e.g. 95th percentile) or a “worst case”, but rather as genuinely implausible owing to the implausibility of the inputs required to reach a forcing of 8.5 W/m². They noted that RCP8.5 has already diverged from observed trends in energy use and the near future trends diverge sharply from those of the International Energy Agency (IEA), which provides market-based projections of energy use for the coming decades. Pielke Jr. *et al.* (2022) further showed that the historic and projected IEA trends run near the bottom of the envelopes of both RCP projections and the more recent Shared Socioeconomic Pathway (SSP) scenario trends.

Schwalm *et al.* (2020) defended the use of RCP8.5 on the grounds that cumulative CO₂ emissions over 2005-2020 track it more closely than the lower RCP scenarios. They also argue that a modified version of the IEA scenarios closely track RCP8.5 in the coming decades. Hausfather and Peters (2020b) responded that the skill of RCP8.5 over those 15 years is due to offsetting errors in its representation of CO₂ from fuel use and land use change, and the apparent agreement with IEA in coming decades is due to Schwalm *et al.* adding in very high land use emissions. The IEA’s own projected CO₂ emissions track well below RCP8.5.

Widespread use of RCP8.5 as a no-policy baseline has created a bias towards alarm in the climate impacts literature. The extent of this problem was confirmed in a literature analysis by Pielke Jr. and Ritchie (2020). They found that some 16,800 scientific papers published between 2010 and 2020 used the RCP8.5 scenario, with about 4,500 of the articles linking RCP8.5 to the concept of “business-as-usual”. Their analysis showed how RCP8.5 was misused not only by individual researchers, but also by influential science agencies like the IPCC and the U.S. National Climate Assessment (USNCA), which has directly led to misleading coverage in prominent media outlets.

¹ This extreme scenario is useful for modelers, since a large forcing generates a large response (warming) making it easier to assess a model’s sensitivity. But that’s very different than claiming it is a plausible future outcome.




Figure 3.2.1 Since the 1970s successive families of emission and concentration projections (colored lines) have consistently overestimated observations (black line). Source: Hausfather *et al.* (2019) Figure S4.

Pielke and Ritchie (2020) reported new studies using RCP8.5 were published at a rate of about 20 per day, with about two per day specifically linking RCP8.5 and “business as usual.” They conclude that the climate research community has spent a decade “committing scientific resources to science fiction” and that “The scientific literature has become imbalanced in an apocalyptic direction.”

The IPCC developed a new set of scenarios for AR6, the “Shared Socioeconomic Pathway” (SSP) scenarios, which have continued the bias shown in the RCP and SRES scenarios. Figure 3.2.2 shows total global observed CO₂ emissions compiled by the International Energy Agency (IEA) merged with the emission projection from the IEA taking account of energy use projections and current policies. The other lines show the range of IPCC SSP scenarios (SSP1-SSP5). As of 2023, global CO₂ emissions have been tracking well below SSP7.0 and are even below SSP2-4.5.




Figure 3.2.2. Observed and projected CO₂ emissions. Source: IPCC (SSP scenarios) and Energy Information Administration (EIA). Green: observed historical emissions and EIA projections. Other lines: SSP1-5. Data source: Friedlingstein *et al.* (2024).

3.2.2 The carbon cycle relating emissions and concentrations

Carbon dioxide emissions from fossil fuel burning (and to a lesser extent deforestation and cement production) have led to steadily increasing CO₂ concentrations in the atmosphere, as shown in Fig. 3.1.3. The relation between emissions and concentration is determined by the global carbon cycle of land and ocean processes that exchange carbon with the atmosphere. Our understanding of these processes was reviewed by Crisp *et al.* (2021).

There are about 850 Gt of carbon (GtC) in the Earth's atmosphere², almost all of it in the form of CO₂. Each year, biological processes (plant growth and decay) and physical processes (ocean absorption and outgassing) exchange about 200 GtC of that carbon with the Earth's surface (roughly 80 GtC with the land and 120 GtC with the oceans). Before human activities became significant, removals from the atmosphere were roughly in balance with additions. But burning fossil fuels (coal, oil, and gas) removes carbon from the ground and adds it to the annual exchange with the atmosphere. That addition (together with a much

² Because CO₂ is chemically transformed through the course of the carbon cycle, it is more convenient to track atoms of carbon rather than molecules of CO₂. One gigatonne of carbon (GtC) is equivalent to about 3.7 Gt of CO₂.

smaller contribution from cement manufacturing) amounted to 10.3 GtC in 2023, or only about 5 percent of the annual exchange with the atmosphere.

The carbon cycle accommodates about 50 percent of humanity’s small annual injection of carbon into the air by naturally sequestering it through plant growth and oceanic uptake, while the remainder accumulates in the atmosphere (Ciais *et al.*, 2013). For that reason, the annual increase in atmospheric CO₂ concentration averages only about half of that naively expected from human emissions.

To project future CO₂ concentrations in the atmosphere, and hence future human influences on the climate, it is important to know how the carbon cycle might change in the future. The historical near constancy of that 50 percent fraction means that the more CO₂ humanity has produced, the faster nature removed it from the atmosphere. That 50 percent fraction changes somewhat from year to year due to natural carbon cycle imbalances from El Nino, La Nina, and varying weather patterns. There was also a substantial additional reduction in atmospheric CO₂ after the 1991 eruption of Mt. Pinatubo, a curious result that has yet to be explained (Angert *et al.*, 2004).

The main processes that remove excess CO₂ from the atmosphere are increased growth of land vegetation (especially at high latitudes), some increase in the sequestering of carbon in soils, and uptake of CO₂ by the ocean due to the increasing partial pressure of atmospheric CO₂ over that of CO₂ dissolved in the ocean. All twenty land carbon cycle models tracked by the Global Carbon Project (Friedlingstein *et al.*, 2024) show land processes have been removing excess CO₂ at an increasing rate since 1959. This is consistent with a “global greening” phenomenon (Chapter 2.1) observed by satellites since monitoring of global greenness began in 1982.

While land vegetation has been responding positively to more atmospheric CO₂, uptake of extra CO₂ by ocean biological processes remains too uncertain to be measured reliably. Our current understanding of these and many more carbon cycle processes was reviewed by Crisp *et al.* (2021).

CO₂ uptake by land processes

The uptake of extra CO₂ from the atmosphere by land surface processes (as also inferred from global greening) has been modeled with 20 different dynamic global vegetation models, the outputs of which are updated every year by the Global Carbon Project (Friedlingstein, 2024). As seen in Fig. 3.2.3, all of those models agree that vegetation and soils have been sequestering carbon from the atmosphere. But we also see that the long-term trends over 1959 to 2023 (65 years) vary widely between models, by nearly a factor of 7. This demonstrates that there remains considerable uncertainty in how fast land processes are removing CO₂ from the atmosphere, which in turn creates uncertainty in future atmospheric CO₂ concentrations, which then produce uncertainty in climate model simulations of future climate change.




Figure 3.2.3 Trends of annual CO₂ uptake (GtCO₂ per year per decade) by land processes during 1959-2023 simulated by 20 different dynamic global vegetation models periodically reported by the Global Carbon Project (Friedlingstein, 2024).

CO₂ uptake by ocean processes

The uptake of extra CO₂ from the atmosphere by ocean processes has been modeled with 10 different ocean biogeochemistry models, the outputs of which are updated every year by the Global Carbon Project (Friedlingstein, 2024). Like the results from the land models, all of the ocean models agree that the global oceans have been sequestering carbon from the atmosphere at an increasing rate during 1959-2023 (Fig. 3.2.4). Unlike the land models, however, the ocean models show much better agreement with each other, with the model producing the fastest increasing CO₂ uptake being only 65 percent faster than the model with the most slowly increasing CO₂ uptake. In spite of the relative agreement among models, Friedlingstein *et al.* (2022) notes that there is substantial discrepancy between the different methods on the strength of the ocean sink over the last decade, particularly in the Southern Ocean.

Note that the average trend in CO₂ uptake across all land models in Fig. 3.2.3 is 25 percent larger than the average trend in ocean uptake. This suggests land processes are increasing in their ability to remove CO₂ faster than ocean processes are increasing their CO₂ sequestration.




Figure 3.2.4 Trends of annual CO₂ uptake (GtCO₂ per year per decade) by ocean processes during 1959-2023 simulated by 10 different ocean biogeochemistry models periodically reported by the Global Carbon Project (Friedlingstein, 2024).

3.3 Urbanization influence on temperature trends

Historical temperature data over land has been collected mainly where people live. This raises the problem of how to filter out non-climatic warming signals due to Urban Heat Islands (UHI) and other changes to the land surface. If these are not removed the data might over-attribute observed warming to greenhouse gases. The IPCC acknowledges that raw temperature data are contaminated with UHI effects but claims to have data cleaning procedures that remove them. It is an open question whether those procedures are sufficient.

AR6 downplayed this issue by saying (WGI p. 235) that no recent evidence had emerged to alter the AR5 finding that urbanization causes an upward bias of no more than 10 percent in the global land surface warming trend. AR5 (WGI p. 189) also cited the 10 percent upper bound without citing a source. AR4 (WGI p. 244) cited Jones *et al.* (1990) and Peterson *et al.* (1999) as the basis of the claim. Peterson *et al.* had failed to find any difference in trends between rural and urban samples, although their definition of rural included local populations up to 10,000 persons while the relative influence of urbanization begins well below that (Spencer *et al.*, 2025). Jones *et al.* compared rural/urban warming in three regions: Eastern Australia, Eastern China and Western Soviet Union. Their definition of “rural” included towns of up to

10,000 in the former Soviet Union and up to 100,000 in China. They found relative warming biases greater than 10 percent in these areas but conjectured that the urbanization effect averaged over the areas they did not examine would bring the global land bias to under 10 percent of the observed warming trend.

Several papers appeared in print prior to the IPCC AR4 that argued that the warming effect of UHIs added a relatively large (30-50%) component to observed warming and was not simulated by climate models (de Laat and Maurellis 2006, McKittrick and Michaels 2007). These findings were based on correlations between locations of maximum warming over land with locations of maximum socioeconomic development. AR4 asserted (p. 244) that these correlations were an artefact of natural atmospheric circulations and were in fact statistically insignificant, and on that basis set the findings aside. Their claim was controversial because it was presented with no supporting evidence. McKittrick (2010) and McKittrick and Nierenberg (2010) showed that taking into account various conjectured alternative explanations for the correlations did not affect their significance. AR5 (p. 189) conceded that AR4 had provided “no explicit evidence” for its assessment and further acknowledged, based on these papers, that there was “significant evidence for such contamination of the record” i.e. a warming bias in the land record. However as already noted, elsewhere in the AR5 report they carried forward AR4 claim that it was less than 10 percent of observed warming. Further they provided no caution about using the land record for climate measurement despite conceding the evidence for UHI contamination. Recently Soon *et al.* (2023) estimated an urbanization bias in the Northern Hemisphere land record over 1850-2018 sufficient to increase the trend in the blended record from 0.55°C to 0.89°C per century.

Some studies providing evidence against UHI contamination compared warming rates between rural and urban locations (Jones *et al.* 1990, Peterson *et al.* 1999, Wickham *et al.* 2013). It is not known whether such methods would be capable of detecting UHI bias even when present. The influence of UHI warming is logarithmic in population, in other words it is strongest at low population density then levels off as local urbanization expands (Oke 1973, Spencer *et al.* 2025). Hence failure to find a difference in warming rates between urban and rural stations does not prove the absence of UHI contamination. McKittrick (2013) provided an empirical demonstration in which the rural/urban trends were not significantly different in a data set shown on other grounds to be contaminated with UHI bias.

Parker (2006) examined a sample of urban locations and found no difference in trends between subsets partitioned according to nighttime wind speed, concluding on this basis that urbanization could not be a significant factor. Here again the question is whether such a method would find UHI bias even if present. McKittrick (2013) presented an example in which UHI-contaminated data did not exhibit significant trend differences when stratified according to wind speed.

The challenge in measuring UHI bias is relating local temperature change to a corresponding change in population or urbanization, rather than to a static classification variable such as rural or urban. Spencer *et al.* (2025) used newly available historical population archives to undertake such an analysis and found evidence of significant UHI bias in U.S. summertime temperature data.

In summary, while there is clearly warming in the land record, there is also evidence that it is biased upward by patterns of urbanization and that these biases have not been completely removed by the data processing algorithms used to produce climate data sets.

References

- Angert, A., S. Biraud, Bonfils, C., Buermann, W. and I. Fung (2004). CO2 seasonality indicates origins of post-Pinatubo sink. *Geophysical Research Letters* 31. <https://doi.org/10.1029/2004GL019760>
- AR6: *Intergovernmental Panel on Climate Change Sixth Assessment Report* (2021) Working Group I Contribution. www.ipcc.ch.

- AR5: *Intergovernmental Panel on Climate Change Fifth Assessment Report* (2013) Working Group I Contribution. www.ipcc.ch.
- AR4: *Intergovernmental Panel on Climate Change Fourth Assessment Report* (2007) Working Group I Contribution. www.ipcc.ch.
- Burgess, Matthew et al (2021) *Environmental Research Letters* 16 014016 <https://iopscience.iop.org/article/10.1088/1748-9326/abcdd2/meta>
- Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, et al. (2013): Carbon and Other Biogeochemical Cycles. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, et al. (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA
- Connolly, Roman, Willie Soon, Michael Connolly et al. (2021) How much has the Sun influenced Northern Hemisphere temperature trends? An ongoing debate *Research in Astronomy and Astrophysics* 21(6) doi: 10.1088/1674-4527/21/6/131 <https://iopscience.iop.org/article/10.1088/1674-4527/21/6/131>
- Crisp, David & Dolman, Han (A.J.) & Tanhua, Toste & Mckinley, Galen & Hauck, Judith & Bastos, Ana & Sitch, Stephen & Eggleston, Simon & Aich, Valentin. (2022). How Well Do We Understand the Land-Ocean-Atmosphere Carbon Cycle?. *Reviews of Geophysics*. 60. 10.1029/2021RG000736.
- De Laat, A.T.J., and A.N. Maurellis (2006), Evidence for influence of anthropogenic surface processes on lower tropospheric and surface temperature trends, *International Journal of Climatology* 26:897—913.
- Friedlingstein, P., and 95 co-authors (2024): Global Carbon Budget 2024, *Earth System Science Data* 14(4), <https://essd.copernicus.org/preprints/essd-2024-519>
- Hausfather et al. (2019) “Evaluating the Performance of Past Climate Model Projections” *Geophysical Research Letters* 47(1) <https://doi.org/10.1029/2019GL085378>
- Hausfather, Z. and G. Peters (2020a) “Emissions – the ‘business as usual’ story is misleading” *Nature* 29 January 2020 <https://www.nature.com/articles/d41586-020-00177-3>
- Hausfather, Z. and G. Peters (2020b) RCP8.5 is a problematic scenario for near-term emissions. *Proceedings of the National Academy of Sciences* 117, 27791–27792 (2020)
- Jenkins, S., Smith, C., Allen, M. et al. Tonga eruption increases chance of temporary surface temperature anomaly above 1.5 °C. *Nature Climate Change*. 13, 127–129 (2023). <https://doi.org/10.1038/s41558-022-01568-2>
- Jones, P. D., P. Y. Groisman, M. Coughlan, N. Plummer, W.-C. Wang, and T. R. Karl (1990), Assessment of urbanization effects in time series of surface air temperature over land, *Nature*, 347, 169 – 172
- Liu, Pengfei et al. (2021) “Improved estimates of preindustrial biomass burning reduce the magnitude of aerosol climate forcing in the Southern Hemisphere” *Science Advances* 7(22) May 2021 <https://doi.org/10.1126/sciadv.abc1379>
- McKittrick, R.R. and P.J. Michaels (2007), Quantifying the influence of anthropogenic surface processes and inhomogeneities on gridded global climate data, *Journal of Geophysical Research*, 112, D24S09, doi:10.1029/2007JD008465.
- McKittrick, Ross R. (2010) *Atmospheric Oscillations Do Not Explain the Temperature-Industrialization Correlation. Statistics Politics and Policy* Vol 1. No. 1., July 2010
- McKittrick, Ross R. (2013) Encompassing Tests of Socioeconomic Signals in Surface Climate Data. *Climatic Change* doi 10.1007/s10584-013-0793-5. Volume 120, Issue 1-2. <http://link.springer.com/article/10.1007/s10584-013-0793-5>
- McKittrick, Ross R. and Nicolas Nierenberg (2010) *Socioeconomic Patterns in Climate Data. Journal of Economic and Social Measurement*, 35(3,4) pp. 149-175. DOI 10.3233/JEM-2010-0336
- McKittrick, Ross R., Mark Strazicich and Junsoo Lee (2012) “Long-Term Forecasting of Global Carbon Dioxide Emissions: Reducing Uncertainties Using a Per-Capita Approach.” *Journal of Forecasting*, Vol 32, Issue 5, pp 435-451 DOI: 10.1002/for.2248.
- Oke, T.R., 1973: City size and the urban heat island, *Atmospheric Environment* 7, 769-779

- Parker, D.E. (2006) “A Demonstration that Large-Scale Warming is not Urban.” *Journal of Climate* 19:2882—2895.
- Peterson, Thomas C., Kevin P. Gallo, Jay Lawrimore, Timothy W. Owen, Alex Huang, David A. McKittrick (1999) Global rural temperature trends. *Geophysical Research Letters* February 1999 <https://doi.org/10.1029/1998GL900322>
- Pielke Jr., Roger and Ritchie, Justin (2020) “Systemic Misuse of Scenarios in Climate Research and Assessment” Social Sciences Research Network April 2020, available at: <https://ssrn.com/abstract=3581777>
- Pielke Jr, R., Burgess, M. G., & Ritchie, J. (2022). Plausible 2005-2050 emissions scenarios project between 2 and 3 degrees C of warming by 2100. *Environmental Research Letters* 17 024027 <https://iopscience.iop.org/article/10.1088/1748-9326/ac4ebf/pdf>
- Scaffeta, Nicola, Richard C. Willson, Jae N. Lee and Dong Wu (2019) Modeling Quiet Solar Luminosity Variability from TSI Satellite Measurements and Proxy Models during 1980–2018. *Remote Sensing* 11(21) 2569 <https://doi.org/10.3390/rs11212569>
- Schoeberl, M.R., Y. Wang, G. Taha, D.J. Zawada, R. Ueyama and A. Dessler, 2024. Evolution of the climate forcing during the two years after the Hunga Tonga-Hunga Ha’apai eruption. *Journal of Geophysical Research.*, 129.
- Schwalm, C.R., S. Glendon, P. B. Duffy (2020) RCP8.5 tracks cumulative CO₂ emissions. *Proceedings of the National Academy of Sciences U.S.A.* 117, 19656–19657 (2020).
- Soon, W.; Connolly, R.; Connolly, M.; Akasofu, S.-I.; Baliunas, S.; *et al.* (2023) The Detection and Attribution of Northern Hemisphere Land Surface Warming (1850–2018) in Terms of Human and Natural Factors: Challenges of Inadequate Data. *Climate* 2023, 11, 179. <https://doi.org/10.3390/cli11090179>
- Spencer, Roy W, John R Christy and William D. Braswell (2025) Urban Heat Island Effects in U.S. Summer Surface Temperature Data, 1895–2023 *Journal of Applied Meteorology and Climatology* April 2025 <https://doi.org/10.1175/JAMC-D-23-0199.1>
- Wickham C, R Rohde , RA Muller, J Wurtele, J Curry, *et al.* (2013) Influence of Urban Heating on the Global Temperature Land Average using Rural Sites Identified from MODIS Classifications. *Geoinformatics and Geostatistics: An Overview* 1:2.
- Zacharias, Pia (2014) An Independent Review of Existing Total Solar Irradiance Records. *Surveys in Geophysics* 35 pp. 897—912 <https://link.springer.com/article/10.1007/s10712-014-9294-y>

PART II: CLIMATE RESPONSE TO CO₂ EMISSIONS

4 CLIMATE SENSITIVITY TO CO₂ FORCING

Chapter Summary

There is growing recognition that climate models are not fit for the purpose of determining the Equilibrium Climate Sensitivity (ECS) of the climate to increasing CO₂. The IPCC has turned to data-driven approaches including historical data and paleoclimate reconstructions, but their reliability is diminished by data inadequacies.

Data-driven ECS estimates tend to be lower than climate model-generated values. The IPCC AR6 upper bound for the likely range of ECS is 4.0°C, lower than the AR5 value of 4.5°C. This lowering of the upper bound seems well justified by paleoclimatic data. The IPCC AR6 lower bound for the likely range of ECS is 2.5°C, substantially higher than the AR5 value of 1.5°C. This raising of the lower bound is less justified; evidence since the AR6 finds the lower bound of the *likely* range to be around 1.8°C.

4.1 Introduction

The magnitude of the climate's response to increasing concentrations of CO₂ is central to the scientific debate on anthropogenic climate change, and so also to the public debate on “climate action.” The simplest measure of that response is the rise in the global average surface temperature, quantified by the Equilibrium Climate Sensitivity (ECS). ECS is defined as the amount of warming expected in response to a doubling of CO₂ from its pre-industrial concentration of 280 ppm, after all climate components have had time to adjust. Some components, like temperatures in the lower atmosphere (troposphere), adjust rapidly, while others such as the deep ocean and cryosphere might take as long as centuries. A related measure, the Transient Climate Response (TCR), better describes the shorter time scales; it is defined as the amount of warming when the CO₂ concentration is doubled by rising one percent annually for 70 years.

The 1979 Charney Report for the U.S. National Academy of Sciences (National Research Council 1979) proposed that the most likely ECS was $3.0 \pm 1.5^\circ\text{C}$. The IPCC repeatedly reaffirmed that range with only minor variations until its most recent AR6. AR5 termed 1.5–4.5°C as the *likely* range (66 percent probability) and stated that ECS is *extremely unlikely* (95 percent probability) to be below 1.0°C and *very unlikely* (90 percent probability) to exceed 6.5°C.

The uncertainty in ECS has remained stubbornly wide, despite many individual studies that claimed to narrow it (Hausfather 2023). Most recently, AR6 narrowed the *likely* range to 2.5–4.0°C and deemed the *very likely* range to be 2.0–5.0°C. This narrowing on the low end is disputed, as will be discussed below.

Uncertainties in ECS are highly consequential for policy making. As will be discussed in Chapter 11, economic models use ECS values to project the costs of CO₂ emissions. The traditional value (3.0 °C) has typically yielded modest global social costs of CO₂ emissions, sufficient to justify some policy actions, but mostly deferred to later in this century. If ECS is very high (above 4.5°C) immediate aggressive emission controls become more imperative, whereas no CO₂ emission controls are economically justifiable for ECS below 2.0°C (Dayaratna *et al.* 2017, 2020). Obtaining a precise estimate is impossible, so policy making needs to account for the uncertainty.

By itself, the equilibrium warming effect of a doubling of atmospheric CO₂ is slightly more than 1°C (Soden and Held 2006). Larger values of ECS arise from positive feedbacks that amplify the CO₂ warming. Water vapor feedback is positive: a warmer atmosphere might have more water vapor, which itself is a powerful greenhouse gas. Warmer temperatures also result in less snow and sea ice cover, allowing the

Earth to absorb more of the sun's radiation. Some simple estimates of these feedbacks increase the ECS to around 2°C (Sherwood *et al.*, 2020). Larger values of ECS are associated with positive cloud feedbacks.

Climate scientists use multiple lines of evidence to determine the Equilibrium Climate Sensitivity:

- Climate model simulations
- Historical observations
- Paleoclimatic reconstructions
- Process understanding of feedbacks

4.2 Model-based estimates of climate sensitivity

The ECS ranges given in IPCC AR4 and AR5 were obtained primarily by examining the behavior of large-scale climate models, also called General Circulation Models (GCMs). However, the IPCC changed course in its AR6 when it turned to a more direct data-driven methodology. Here we discuss some of the pitfalls of using GCMs to try to determine the Earth's climate sensitivity.

In principle, ECS is an emergent property of GCMs—that is, it is not directly parameterized or tuned, but rather emerges in the results of the simulation. Otherwise plausible GCMs and parameter selections have been discarded because of perceived conflict with an expected warming rate, or aversion to a model's climate sensitivity being outside an accepted range (Mauritsen *et al.* 2012). This practice was commonplace for the models used in AR4; modelers have moved away from this practice with time. However, even in a CMIP6 model, Mauritsen and Roeckner (2020) state the following regarding their Max Planck Institute (MPI) climate model (emphasis added):

“We have documented how we tuned the MPI-ESM1.2 global climate model to match the instrumental record of warming; an endeavor which has clearly been successful. Due to the historical order of events, the choice was to do this practically by **targeting an ECS of about 3 K using cloud feedbacks**, as opposed to tuning the aerosol forcing.”

In other words, the MPI modelers chose an ECS value of 3°C and then tuned the cloud parameterizations to match their intended result.

As noted, direct warming from CO₂ doubling is only about 1°C (Soden and Held 2006); further warming arises from climate feedbacks that are not explicitly resolved by the GCM but rely on parameterizations of physical processes. Higher values of ECS arise primarily from positive cloud feedbacks, whereas the magnitude and even the sign of the feedbacks are very uncertain. Elements of cloud feedback include changes in the latitudinal distribution of clouds, changes in the distribution of cloud height (changes in low versus high clouds), changes to the phase of clouds (ice versus liquid), changes in cloud particle size (associated with changes in concentration and/or composition of aerosol particles), changes in the precipitation efficiency of clouds, and even changes in how clouds are distributed over the daily solar cycle (Curry and Webster, 1999). It is difficult for GCMs to simulate any of these processes correctly owing to their small scale, let alone predict how they will change in the future. Further, cloud processes modulate the magnitudes of the water vapor, lapse rate, and the surface albedo feedbacks.

ECS can be determined from climate model simulations by doubling the concentration of CO₂ and allowing several centuries for the warming to equilibrate. To avoid the need for such long simulations, “effective climate sensitivity” is commonly evaluated from a 150-year simulation in response to a sudden quadrupling of CO₂.




Figure 4.1 Equilibrium Climate Sensitivities in °C of 37 climate models from the CMIP6 ensemble. Identifiers for the various models appear along the horizontal axis. From (Scafetta, 2021)

The spread of ECS values from the CMIP5 ensemble of climate models used in AR5 was 2.0–4.7°C; that range increased for the CMIP6 models used in AR6 to between 1.8 and 5.7°C (Chen *et al.*, 2021, Scafetta 2021, see Figure 4.1). Far from resolving the model-based climate sensitivity the range appears to be growing. The main cause of the overall upward shift in ECS in CMIP6 relative to CMIP5 is a larger positive cloud feedback, driven by changes to the cloud parameterizations in many CMIP6 models (Zelinka *et al.*, 2020)

Because of concerns about model tuning and the high sensitivity to cloud parameterizations, the IPCC AR6 (2021) did not rely on climate model simulations in their assessment of climate sensitivity, relying instead on data-driven methods.

4.3 Data-driven estimates of climate sensitivity

Climate sensitivity can also be estimated from instrumental records of surface temperatures and ocean heat content, combined with estimates of how climate forcings (e.g., greenhouse gases, solar, volcanoes, aerosols) have changed in the past (Otto *et al.*, 2013). Using this information, a simple empirical model called an Energy Balance Model can be employed. It requires estimating a feedback parameter whose uncertainties are highly amplified in the resulting ECS (Roe and Baker, 2007).

The accuracy of the data-driven methods depends on the quality of the input data. Assumptions are needed about ocean heat storage, and good data is only available for recent decades. The greatest source of

uncertainty is the amount and composition of aerosol particles and their interactions with cloud radiative properties (the so-called aerosol indirect effect; see Figures 3.1.1, 3.1.2). Climate models exhibit warming in response to GHGs but cooling in response to aerosols (Schwartz *et al.*, 2007). Observed 20th century warming can be shown to be consistent either with low ECS and low aerosol cooling, or high ECS and high aerosol cooling. Since fossil fuel use adds both GHGs and aerosols to the atmosphere, both effects need to be estimated to isolate the warming effect of CO₂.

Paleoclimate proxies are also used to evaluate the sensitivity of past climates by comparing paleoclimate changes in the Earth's temperatures to estimates of changes in forcings. The two most informative periods are the last glacial maximum (around 20,000 years ago), which was about 3–7°C colder than today, and a mid-Pliocene period (roughly three million years ago), which was 1–3°C warmer than today. The limits on cooling during the last glacial maximum give the best single evidence that high values of climate sensitivity are unlikely. However, paleoclimate estimates are associated with very large uncertainties in the estimated temperatures and forcings. Further, estimates of climate sensitivity based on past climate states might not be applicable to the current state of the climate system.

A recurring theme in the climate literature is that ECS estimates based on historical data are smaller than ECS estimates inferred from climate models (Sherwood and Forest 2024). About 15 estimates based on historical data appeared in the peer-reviewed literature between 2012 and 2024 yielding ECS best estimates between 1.0°C and 2.5°C, although critics have questioned some of the methods and the data quality. For AR6, the IPCC placed primary weight on the results of Sherwood *et al.* (2020) that combined historical data and paleoclimate proxies with the process-based approach and yielded a best estimate of 3.1°C with a *likely* range of 2.6–3.9°C. Lewis (2022) raised a number of concerns about this result, including methodological errors, outdated input values, and use of subjective Bayesian priors in the analysis. Lewis' analysis found that climate sensitivity is estimated to be much lower and better constrained than in the Sherwood *et al.* analysis – median 2.2°C (1.8–2.7°C in the 17–83 percent *likely* range, and 1.6–3.2°C in the 5–95 percent *very likely* range). The IPCC AR6 estimated only a 5 percent probability that ECS was below 2.3°C, whereas Lewis estimated it to be over 50 percent. The most recent publications on the debate between Sherwood *et al.* and Lewis further defend their respective positions: Sherwood and Foster (2024) and Lewis (2025).

An argument emphasized in AR6 is that data-driven ECS estimates might understate the future warming response to GHGs because of a so-called “pattern effect” (Forster *et al.*, 2021). The tropical Pacific is believed to strongly influence the overall efficiency with which the Earth radiates heat to space, but some regions remove heat more efficiently than others. If the west-to-east temperature gradient in the tropical Pacific is weakened in a warming climate, warming would concentrate where heat is less efficiently removed, raising ECS.

Most climate models simulate that rising GHGs will weaken the west-east temperature gradient, which led the IPCC in AR6 to conclude that data-driven ECS estimates understated the *future* ECS value. However, Seager *et al.* (2019) pointed out that, contrary to models, the west-east temperature gradient has been strengthening over time. They further argued that the mechanism predicting otherwise in climate models was based on a faulty characterization of oceanic dynamics and there is no reason to expect the gradient to weaken. A similar argument was recently made by Lee *et al.* (2024), who concluded that “the trajectory of the observed trend reflects the response to increasing GHG loading in the atmosphere”; in other words, GHG warming should lead to a future strengthening rather than a weakening of the temperature gradient. Increased efficiency of atmospheric cooling implies, if anything, that the future ECS in a warming climate might be *lower* than current estimates.

4.4 Transient Climate Response

The Transient Climate Response (TCR) provides a more useful observational constraint on climate sensitivity. TCR is the global temperature increase that results when CO₂ is increased at an annual rate of 1 percent over a period of 70 years (*i.e.*, doubled gradually). Relative to the ECS, observationally determined values of TCR avoid the problems of uncertainties in ocean heat uptake and the fuzzy boundary in defining equilibrium arising from a range of timescales for the longer-term feedback processes (*e.g.*, ice sheets). TCR is better constrained by historical warming, than ECS. IPCC AR6 judged the *very likely* range of TCR to be 1.2–2.4°C. In contrast to ECS, the upper bound of TCR is more tightly constrained. For comparison, the TCR values determined by Lewis (2023) are 1.25 to 2.0°C, showing much better agreement with AR6 values than was seen in a comparison of the ECS values.

References

- Carbon Brief. (2020, July 22). Explainer: How scientists estimate climate sensitivity. <https://www.carbonbrief.org/explainer-how-scientists-estimate-climate-sensitivity>
- Carbon Brief. (2020, July 22). Guest post: Why low-end ‘climate sensitivity’ can now be ruled out. <https://www.carbonbrief.org/guest-post-why-low-end-climate-sensitivity-can-now-be-ruled-out>
- Carbon Brief. (2021, August 9). In-depth Q&A: The IPCC’s sixth assessment report on climate science. <https://www.carbonbrief.org/in-depth-qa-the-ipccs-sixth-assessment-report-on-climate-science>
- Chen, D., *et al.* (2021). Framing, context, and methods. In V. Masson-Delmotte *et al.* (Eds.), *Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Curry, J. A., & Webster, P. J. (1999). Thermodynamic feedbacks in the climate system. In *Thermodynamics of atmospheres and oceans* (pp. 351–385). Academic Press.
- Dayaratna, Kevin, Ross McKittrick and David Kreutzer (2017) Empirically-Constrained Climate Sensitivity and the Social Cost of Carbon. *Climate Change Economics* April 2017 DOI: <http://dx.doi.org/10.1142/S2010007817500063>
- Dayaratna, Kevin, Ross McKittrick and Patrick J. Michaels (2020) Climate Sensitivity, Agricultural Productivity and the Social Cost of Carbon in FUND. *Environmental Economics and Policy Studies* <https://doi.org/10.1007/s10018-020-00263-w>
- Forster, P., *et al.* (2021). The Earth’s energy budget, climate feedbacks, and climate sensitivity. In V. Masson-Delmotte *et al.* (Eds.), *Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Hausfather, Z. (2020, July 22). Explainer: How scientists estimate climate sensitivity. Carbon Brief. <https://www.carbonbrief.org/explainer-how-scientists-estimate-climate-sensitivity>
- Lan, X., Tans, P., & Thoning, K. W. (2025). Trends in globally-averaged CO₂ determined from NOAA Global Monitoring Laboratory measurements. NOAA Global Monitoring Laboratory. <https://doi.org/10.15138/9N0H-ZH07>
- Lee, S., Byrne, M. P., Loikith, P. C., & O’Dell, C. W. (2024). Zonal contrasts of the tropical Pacific climate predicted by a global constraint. *Climate Dynamics*, 62(1–2), 229–246. <https://doi.org/10.1007/s00382-023-06741-7>
- Lewis, N. (2023). Objectively combining climate sensitivity evidence. *Climate Dynamics*, 61(9–10), 3155–3163. <https://doi.org/10.1007/s00382-022-06398-8>
- Lewis, N. (2025). Comment on “Can uncertainty in climate sensitivity be narrowed further?” by Sherwood and Forest (2024). *EGUsphere* [preprint]. <https://doi.org/10.5194/egusphere-2025-1179>
- Thorsten Mauritsen *et al.*, (2012) “Tuning the Climate of a Global Model,” *Journal of Advances in Modeling Earth Systems* 4, no. 3 <https://doi.org/10.1029/2012ms000154>

- Mauritsen, T., & Roeckner, E. (2020). Tuning the MPI-ESM1.2 global climate model to improve the match with instrumental record warming by lowering its climate sensitivity. *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002037. <https://doi.org/10.1029/2019MS002037>
- Otto, A., Otto, F. E. L., Boucher, O., Church, J., Hegerl, G., Forster, P. M., Gregory, J. M., & Johnson, G. C. (2013). Energy budget constraints on climate response. *Nature Geoscience*, 6(6), 415–416. <https://doi.org/10.1038/ngeo1836>
- Pachauri, R. K., & Meyer, L. (Eds.). (2015). Climate change 2014: Synthesis report. *Intergovernmental Panel on Climate Change*.
- Roe, G. and M. Baker (2007). Why is climate sensitivity so unpredictable? *Science* 318, 629–632. <https://doi.org/10.1126/science.1144735>
- Scafetta, N. (2021). Testing the CMIP6 GCM simulations versus surface temperature records from 1980–1990 to 2011–2021: High ECS is not supported. *Climate*, 9(11), 161. <https://doi.org/10.3390/cli9110161>
- Schwartz, S. E., R. J. Charlson, H. Rodhe (2007). Quantifying climate change—Too rosy a picture? *Nature Climate Change* 1, 23–24, <https://doi.org/10.1038/climate.2007.22>
- Seager, R., Cane, M., Ting, M., Naik, N., Clement, A., DiNezio, P., & Lee, D. E. (2019). Strengthening tropical Pacific zonal sea surface temperature gradient consistent with rising greenhouse gases. *Nature Climate Change*, 9(7), 517–522. <https://doi.org/10.1038/s41558-019-0505-x>
- Sherwood, S. C., & Forest, C. E. (2024). Opinion: Can uncertainty in climate sensitivity be narrowed further? *Atmospheric Chemistry and Physics*, 24(5), 2679–2686. <https://doi.org/10.5194/acp-24-2679-2024>
- Sherwood, S. C., Bony, S., Boucher, O., Bretherton, C. S., Forster, P. M., Gregory, J. M., & Stevens, B. (2020). An assessment of Earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics*, 58(4). <https://doi.org/10.1029/2019rg000678>
- Soden, B. J., and I.M. Held (2006). An assessment of climate feedbacks in coupled ocean–atmosphere models. *Journal of Climate*, 19(14), 3354–3360. <https://doi.org/10.1175/jcli3799.1>
- Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., & Taylor, K. E. (2020). Causes of higher climate sensitivity in CMIP6 models. *Geophysical Research Letters*, 47, e2019GL085782. <https://doi.org/10.1029/2019GL085782>

5 DISCREPANCIES BETWEEN MODELS AND INSTRUMENTAL OBSERVATIONS

Chapter summary

Climate models show warming biases in many aspects of their reproduction of the past several decades. In response to estimated changes in forcing they produce too much warming at the surface (except in the models with lowest ECS), too much warming in the lower-and mid-troposphere and too much amplification of warming aloft.

Climate models also produce too much recent stratospheric cooling, invalid hemispheric albedos, too much snow loss, and too much warming in the Corn Belt. The IPCC has acknowledged some of these issues but not all.

5.1 Introduction

Climate models are the primary tool used to project future climate changes in response to increasing atmospheric levels of anthropogenic greenhouse gases. To assess the fitness of climate models for this purpose, it is reasonable to ask how well they reproduce the current climate and its variations over the past century. The box “Climate modeling” gives some detail on how climate models work.

Of great concern is the fact that, after several decades of the climate modeling enterprise involving approximately three dozen models operated by research centers around the world, the range of future warming they produce in response to a hypothetical doubling of atmospheric CO₂ extends over a factor of three, as we discussed in the previous chapter. This range of disagreement among models has not decreased for decades.

Problems with climate models are not just in their disagreement over the future, but also in their ability to replicate the recent past. Here we review some of the most important metrics of climate model accuracy: ability to reproduce historical surface, tropospheric and stratospheric temperature trends; ability to reproduce the vertical warming profile; and ability to reproduce other climate features such as snowfall. In all cases a persistent finding is that models on average err on the side of too much warming in response to estimated historical forcings.

BOX: Climate modeling

Most climate models, except for the simplest ones, represent Earth's land surface using a grid of squares on the order of 100 km wide. To simulate the atmosphere, typically 30 or more grid boxes are stacked above these squares. The ocean is modeled using a similar but finer grid, resulting in tens of millions of grid boxes for the atmosphere and oceans.

The computer models, based on physical laws, calculate how air, water, and energy move between grid boxes over time. The time step can be as small as 10 minutes, and repeating this process millions of times allows the simulation of climate over centuries. Running these models, even on the most powerful supercomputers, can take months. Comparing simulation results with historical climate data helps assess the accuracy of a model, while projections into the future estimate climate changes under assumed human and natural influences.

Despite sounding straightforward, climate modeling is highly complex. Many critical processes occur at scales smaller than the grid size. For instance, the flows of sunlight and heat in the atmosphere depend strongly on cloud cover. Since tracking individual clouds is impractical, researchers must make “subgrid” assumptions about the distribution of clouds in each grid box. Snow and ice cover, which affects how much sunlight is reflected or absorbed by the surface, is another subgrid factor.

Each subgrid assumption requires numerical parameters, which must be carefully set. Modelers initially estimate these parameters based on physics and observed climate patterns, then run the model. Because early results often diverge significantly from real-world observations, they “tune” these parameters to better match observed climate features. Different modeling teams use distinct assumptions and tuning strategies, leading to varied outcomes. Tuning is a necessary but delicate aspect of climate modeling, as it is for any complex system. Poor tuning can result in inaccurate simulations, while excessive tuning risks artificially steering results toward predetermined conclusions.

The spread of model representations of the current climate is very wide. One of the most basic indicators—Earth's average surface temperature—varies by about 3°C across CMIP6 models prior to 1880 (Figure 5.1), narrows slightly until 2040 then diverges to over 4 °C. For comparison 20th century warming was only about 1.0°C. This variation suggests substantial differences among models' physical processes.




Figure 5.1 CMIP6 Average surface temperature range across 33 models and standard deviation using SSP5-85 scenario. Data from KNMI Climate Explorer website at <https://climexp.knmi.nl/start.cgi>.

Beyond the models' ability to reproduce features of today's climate, the critical issue for society is how well they predict responses to subtle human influences, such as greenhouse gas emissions, aerosol cooling, and land-use changes. The most crucial aspect that models must capture correctly is “feedbacks.” These occur when climate changes either amplify or suppress further warming. In general, the modeled net effect of all feedbacks doubles or triples the direct warming impact of CO₂.

5.2 Surface warming

A straightforward test of a climate model's validity is its ability to reproduce historical warming in response to known past changes in climate drivers such as greenhouse gases. Figure 5.2 is reproduced from Scafetta (2023), that groups the latest-generation (CMIP6) climate models into low ECS (1.5 to 3.0 degrees C), medium ECS (3.0 to 4.5 degrees C) and high ECS (4.5 to 6.0 degrees C), and compares their post-1980 global average temperature simulation ranges to those of three surface temperature records and one satellite-based lower troposphere temperature data product.

The leftmost column shows that the low-ECS models track the post-1980 historical warming record reasonably well, but the middle and right columns show that the medium- and high-ECS models conspicuously over-predict the warming.




Figure 5.2: Model-Observational comparisons for Earth's surface warming. The columns correspond to model groups showing low-ECS (13 models) medium-ECS (11 models) and high-ECS (14 models), while the rows correspond to widely-used observed temperature records, the first three showing surface averages and the fourth showing the lower troposphere average. In each panel the yellow area denotes the mean and range ($\pm$ one standard deviation) of climate model simulations for that group. The thick black line shows the observed annual average temperature in the indicated source. Source: Scafetta (2023) Fig.2.

Spencer (2024) has also provided a useful summary of the Model-Observation mismatch by comparing trends in surface temperature data products with those in individual climate models, as summarized in Figure 5.3; most climate models show substantially more warming than the observations since 1979.




Figure 5.3 Global surface air temperature trends (°C/decade), 1979–2024, from various CMIP6 climate models (red, 30-model average in orange); and the average of three thermometer datasets (HadCRUT5, NOAA Global Temp, and Berkeley 1 deg.) and two reanalysis datasets (ERA5 and NCEP/NCAR R1) in blue. Data source: <https://climexp.knmi.nl/start.cgi>.

5.3 Tropospheric warming

It has long been known that climate models on average overstate warming in the tropical troposphere. This region is an important test of climate models since this is where the signal of anthropogenic greenhouse warming emerges first and most strongly. Biases in tropospheric trends indicate model flaws in heat transfer processes that carry over to surface warming biases.

The discrepancy was flagged as a serious inconsistency in the first U.S. Climate Change Science Program report (Karl *et al.* 2006) and has been mentioned in every IPCC report since, but the discrepancy has gotten worse over time, and the bias is now global. McKittrick and Christy (2020) compared tropospheric warming trends in CMIP6 climate models to observed trends from satellites, weather balloons and reanalysis systems. Every model overpredicted the average observed warming trend over 1979-2014 in both Lower- and Mid-Troposphere layers, both globally and in the tropics. In most individual models the bias was statistically significant and on average across models it was highly significant.

Figure 5.4 presents the comparisons with data updated to 2024 (McKittrick and Christy 2025). The recent warm years moved the observed trend up slightly and widened the trend confidence intervals but the overall pattern remains the same: model bias is towards too much warming, in most cases the difference is statistically significant and on average the bias is statistically highly significant. McKittrick and Christy (2020) also showed that the bias is larger in high-ECS models, but even the models with lower average ECS predict too much warming. If future climate models were to realistically represent global tropospheric warming, they would likely be less sensitive than even the low-ECS members of the CMIP6 ensemble.




Figure 5.4: Observed versus CMIP6 modeled warming trends ($^{\circ}\text{C}/\text{decade}$ 1979-2024) in the global and tropical lower (LT) and mid-troposphere (MT). Uses methodology of McKittrick and Christy (2020) on data updated from 2014 to 2024. Blue dots: warming trends with 95 percent confidence intervals for 3 data products (radiosondes, reanalysis, and satellites). Blue dashed line: warming trend average for 3 observed series. Red dots: modeled warming trends with 95 percent confidence intervals in 35 models arranged lowest to highest.

As mentioned previously, the IPCC has long acknowledged the model-observation mismatch. For example, AR6 pp. 443-444 offers this on the tropical troposphere (it does not address the global comparison):

Several studies since AR5 have continued to demonstrate an inconsistency between simulated and observed temperature trends in the tropical troposphere, with models simulating more warming than observations (Mitchell *et al.*, 2013, 2020; Santer *et al.*, 2017a, b; McKittrick and Christy, 2018; Po-Chedley *et al.*, 2021)... Over the 1979–2014 period, models are more consistent with observations in the lower troposphere, and least consistent in the upper troposphere around 200 hPa, where biases exceed 0.1°C per decade. Several studies using CMIP6 models suggest that differences in climate sensitivity may be an important factor contributing to the discrepancy between the simulated and observed tropospheric temperature trends (McKittrick and Christy, 2020; Po-Chedley *et al.*, 2021), though it is difficult to deconvolve the influence of climate sensitivity, changes in aerosol forcing and internal variability in contributing to tropospheric warming biases (Po-Chedley *et al.*, 2021). Another study found that the absence of a hypothesized negative tropical cloud feedback could explain half of the upper troposphere warming bias in one model (Mauritsen and Stevens, 2015).

... In summary, studies continue to find that CMIP5 and CMIP6 model simulations warm more than observations in the tropical mid- and upper-troposphere over the 1979–2014 period (Mitchell *et al.*, 2013, 2020; Santer *et al.*, 2017a, b; Suárez-Gutiérrez *et al.*, 2017; McKittrick and Christy, 2018), and that overestimated surface warming is partially responsible (Mitchell *et al.*, 2013; Po-Chedley *et al.*, 2021). Hence, we assess with *medium confidence* that CMIP5 and CMIP6 models continue to overestimate observed warming in the upper tropical troposphere over the 1979–2014 period by at least 0.1°C per decade,

Notably, despite the accumulation of evidence of excess model warming the IPCC assigns only *medium confidence* to the existence of a warming bias.

5.4 Vertical temperature profile mismatch

Another important model-observational discrepancy is the excess amplification with altitude found in climate models. The comparison was in the IPCC AR5 Chapter 10, although only in the online supplement (Figure 10.SM.1) and only in a figure whose formatting obscured the point. Figure 10.SM.1 is not referenced in the main IPCC report nor in any summary so readers would not have been aware of it. Although not apparent at first glance, it shows that the 1979–2010 warming in the lower troposphere is so small as to be consistent with no GHG forcing at all and is inconsistent with the model runs that do have GHG forcing. In Figure 5.6 we adapt IPCC AR5 Figure 10.SM.1 to draw out this critical point.




Figure 5.5: Vertical warming pattern for tropics (20S to 20N). Source: Annotated version of IPCC AR5 Figure 10.SM.1

Figure 5.5 compares model and observational temperature trends by altitude between 20S and 20N (the tropics). In this region where the models say the warming should be strongest, the observations (shown here in white) lie within the blue “No CO₂” band and entirely outside the “with CO₂” red envelope. This means that in the entire tropical atmospheric column from the surface to the base of the stratosphere, observed warming trends are so small as to be consistent with the output of models that have no anthropogenic CO₂, and inconsistent with the entire envelope of warming trends generated by models forced with increased CO₂.

A similar comparison is shown in Christy and McNider (2017), an updated version of which (covering 1979-2024) is reproduced as Figure 5.6. Modeled temperature trends exceed observations from the surface through the top of the troposphere, with observed trends below the entire model range at most pressure levels. Also shown in Figure 5.6 is the tropical tropospheric temperature (TTT) average from three satellite data products (NOAA, UAH and RSS) compared to the same layer average from climate models for 1979-2024. Again, the observed trends lie below the entire model range.

The wide range of choices made by modelers to characterize the physical processes in the models (see Box: Climate Modeling in Section 5.1 above) is seen by the large spread of trends in the middle troposphere, ± 40 percent about the median (Figure 5.6). This vividly illustrates the uncertainties in attempts to model (parameterize) a complex system involving turbulence, moist thermodynamics, and energy fluxes over the full range of the tropical atmosphere’s time and space scales.




Figure 5.6: Modeled versus observed warming, tropical troposphere. Source: updated from Christy and McNider (2017) including data through 2024 and CMIP6 model outputs. Red line: model average. Green and blue lines: observational series (reanalysis).

This discrepancy has been the source of much controversy, with some arguing that even if there is very little observed warming aloft in the tropics, a “hotspot” still exists in the sense that the warming aloft is greater than at the surface (Santer et al. 2008). But there is good evidence that models also exaggerate the amplification rate. Klotzbach *et al.* (2009), showed that models project greater amplification with altitude than is observed. This result was subsequently confirmed by detailed time series analysis (Vogelsang and Nawaz 2016) which found that the model-observational difference is statistically significant.

The atmosphere’s temperature profile is a case where models are not merely uncertain but also show a common warming bias relative to observations. This suggests that they misrepresent certain fundamental feedback processes.

The IPCC AR6 did not assess this issue.

5.5 Stratospheric cooling

An important element of the expected general “fingerprint” of anthropogenic climate change is simultaneous warming of the troposphere and cooling of the stratosphere. The latter feature is also influenced by ozone depletion and recovery. AR6 acknowledged that cooling had been observed but only until the year 2000. The stratosphere has shown some warming since, contrary to model projections.

AR6 WG1 Ch 2 pp. 327-9 states:

Temperatures averaged through the full lower stratosphere (roughly 10–25 km) have decreased over 1980–2019 in all data products, with the bulk of the decrease prior to 2000. The decrease holds even if the influence of the El Chichon (1982) and Pinatubo (1991) volcanic eruptions on the trend, found by Steiner *et al.* (2020a) to have increased the 1979–2018 cooling trend by 0.06°C per decade, is removed. Most datasets show no significant or only marginally significant trends over 2000–2019, and the results of Philipona *et al.* (2018) show weak increases over 2000–2015 in the very lowermost stratosphere sampled by radiosondes....

It is virtually certain that the lower stratosphere has cooled since the mid-20th century. However, most datasets show that lower stratospheric temperatures have stabilized since the mid-1990s with no significant change over the last 20 years. It is likely that middle and upper stratospheric temperatures have decreased since 1980, but there is low confidence in the magnitude.

The cited source, Philipona *et al.* (2018), in an article entitled “Radiosondes Show That After Decades of Cooling, the Lower Stratosphere Is Now Warming”, stated:

In response to continued greenhouse gas increases and stratospheric ozone depletion, climate models project continued tropospheric warming and stratospheric cooling over the coming decades. Global average satellite observations of lower stratospheric temperatures exhibit no significant trends since the turn of the century. In contrast, an analysis of vertically resolved radiosonde measurements from 60 stations shows an increase of lower stratospheric temperature since the turn of the century at altitudes between 15 and 30 km and over most continents. Trend estimates are somewhat sensitive to homogeneity assessment choices, but all investigated radiosonde data sets suggest a change from late twentieth century cooling to early 21st century warming in the lower stratosphere.

Santer *et al.* (2023) show using updated data that a cooling trend has not re-emerged in the lower stratosphere.

A combination of tropospheric warming and stratospheric cooling is a commonly cited “fingerprint” of anthropogenic climate change. Stratospheric warming since 2000 coincides with continued surface and tropospheric warming, a pattern that is not found in climate model simulations and is not apparently consistent with the anthropogenic fingerprint.

5.6 Snow cover mismatch

Data compiled by the Rutgers University Snow Lab show that Northern Hemisphere winter snow cover is not decreasing (Figure 5.7); if anything, it shows an increasing trend.




Figure 5.7: Northern Hemisphere Winter Snow extent.

Source: https://climate.rutgers.edu/snowcover/chart_seasonal.php?ui_set=nhland&ui_season=1
(accessed May 27, 2025)

Yet models project declining Northern Hemisphere snow cover in a warming climate, as described by Connolly *et al.* (2019).

The climate models were found to poorly explain the observed trends [in Northern Hemisphere snow cover]. While the models suggest snow cover should have steadily decreased for all four seasons, only spring and summer exhibited a long-term decrease, and the pattern of the observed decreases for these seasons was quite different from the modelled predictions. Moreover, the observed trends for autumn and winter suggest a long-term increase, although these trends were not statistically significant.

AR6 largely confines its discussion of changing Northern Hemisphere snow cover extent (SCE) to the Spring season, for which models and observations agree on a downward trend. Regarding Winter changes it remarks as follows (AR6 WGI Ch. 2 p. 344):

Assessment of SCE trends in the NH since 1978 indicates that for the October to February period there is substantial uncertainty in trends with the sign dependent on the observational product. Analysis using the NOAA Climate Data Record shows an increase in October to February SCE (Hernández-Henríquez *et al.*, 2015; Kunkel *et al.*, 2016) while analyses based on satellite borne optical sensors (Hori *et al.*, 2017) or multi-observation products (Mudryk *et al.*, 2020) show a negative trend for all seasons.

AR6 WGI Chapter 9 (p. 1284) points out that the NOAA Climate Data Record showing increased fall and winter SCE is inconsistent with land-based observations and model-based data sets. It notes that using optical satellite imagery to infer SCE in winter is challenging due to cloud cover and decreased solar illumination in winter months. Focusing on the Pacific Coast states (CA, OR and WA) cold season mountain snowfall that melts in spring and summer provides a substantial portion of warm season water resources. A comprehensive reconstruction of snowfall for the main source regions (Cascades and Sierra Nevada Mountains) indicates no significant trends in annual totals since the late 19th century (Christy 2022).

In summary, the up-to-date Rutgers SCE database indicates a mismatch between models and observations. Further work to reconcile conflicting trends in observational data sets is required.

5.7 Hemispheric symmetry of the planetary albedo

The planetary albedo is the fraction of incoming solar radiation reflected by the Earth back to space. It is an important element of the radiative energy balance and influences whether the planet will warm or cool over time. The planetary albedo is typically estimated at around 0.30; variations on the scale of 0.01 correspond to changes in solar forcing of about 3 W/m^2 , an amount larger than current anthropogenic forcing. It has long been noted that models disagree with each other and with observations on the value of the global planetary albedo (Stephens *et al.* 2015).

An intriguing property of the Earth's albedo is that, on average, the Northern Hemisphere (NH) and Southern Hemisphere (SH) have had nearly the same albedo, at least throughout the fifty-year satellite record (Stephens *et al.*, 2015). This symmetry is surprising, because the SH has much more ocean than land. Since ocean is less reflective than land, the NH should have higher albedo. Clouds (which are highly reflective) are more common in the NH and so compensate the surface albedo imbalances of the two hemispheres. Datseris and Stephens (2021) show that this cloud compensation comes from the extra-tropical storm tracks of the SH, which are cloudier than those of the NH. While the mechanism for this hemispheric symmetry is unclear, it likely operates on large temporal and spatial scales.

The hemispheric symmetry of the albedo is a simple gross metric for climate models. Rugenstein and Hakuba (2023) defined that metric as the difference between the NH and SH annual mean albedos, expressed as Wm^{-2} of the reflected sunlight, and compiled it for the CMIP6 climate models, as shown in Figure 5.8. Most of the CMIP6 models do not reproduce the small observed asymmetry (about 0.1 Wm^{-2}) and even disagree as to which hemisphere is more reflective. Moreover, the magnitude of the asymmetry ranges up to 5 Wm^{-2} in some models, twice as large as the current anthropogenic forcing (about 2.5 Wm^{-2}).

The significance of unphysical albedo asymmetries in the climate models is not yet fully known. However, other model studies suggest that interhemispheric changes in albedo can alter poleward heat fluxes, meridional temperature gradients, storminess, and differences in hemispheric ocean heat storage. The discrepancy between models and observations further raises issues regarding cloud feedback processes and so more generally diminishes confidence in model projections of the future climate.




Figure 5.8. Differences in 20-year average reflectivity (albedo) between the Northern and Southern Hemispheres for CMIP6 models used in the most recent IPCC assessment (colored bars). The very small observed difference is indicated by the vertical black line. From Rugenstein and Hakuba (2023).

5.8 U.S. Corn Belt

One of the largest discrepancies between models and observations is in the U.S. Corn Belt, a region of particular importance to global food production. Figure 5.9 shows the warming trends for summertime (June, July, August) for the 12-state Corn Belt (IN, IA, IL, ND, SD, MO, MN, WI, MI, OH, KS, NE) during 1973-2022. All 36 climate models (red) warm far too rapidly compared to observations (blue).




Figure 5.9: Modeled versus observed warming trends in the U.S. Corn Belt, 1973-2022.

As discussed in Chapter 9, the anticipated negative effects of increasing temperatures on U.S. corn yields have not materialized, in contrast to widely publicized studies proclaiming that theoretical future impacts are already being experienced (*e.g.*, Seager *et al.*, 2018).

The IPCC acknowledges limitations in the accuracy of regional climate model outputs. This example shows that users need to assess model projections carefully on a case-by-case basis since local biases might be sufficiently large that the models are simply not fit for purpose. As has recently been noted by two leaders of the modeling community (emphasis added)

... for many key applications that require regional climate model output or for assessing large-scale changes from small scale processes, we believe that **the current generation of models is not fit for purpose**. (Palmer and Stevens 2019)

To summarize:

- Climate models show warming biases in many aspects of their reproduction of the past few decades.
- They produce too much warming at the surface (except in the models with lowest ECS), too much warming in the lower-and mid-troposphere and too much amplification of warming aloft

- They also produce too much stratospheric cooling, too much snow loss, and too much warming in the U.S. Corn Belt.
- The hemispheric albedo difference in individual climate models ranges widely in sign and magnitude compared to observations. The range in W/m^2 is three times larger than the direct anthropogenic forcing of CO_2 .
- The IPCC has acknowledged some of these issues but not others.

References

- Christy, J. R., R. T. McNider (2017). Satellite bulk tropospheric temperatures as a metric for climate sensitivity. *Asia-Pacific Journal of Atmospheric Sciences* 53, 511-518.
<https://doi.org/10.1007/s13143-017-0070-z>
- Christy, J. R. (2022). Time series construction of Oregon and Washington snowfall since 1890 and an update of California snowfall through 2020. *Journal of Hydrometeorology* 23, 1845-1860.
<https://doi.org/10.1175/JHM-D-21-0178.1>
- Connolly, R., M. Connolly, W. Soon, *et al.* (2019). Northern Hemisphere snow-cover trends (1967–2018): A comparison between climate models and observations. *Geosciences* 9, 135.
<https://doi.org/10.3390/geosciences9030135>
- Datseris, G., B. Stevens (2021). Earth’s albedo and its symmetry. *AGU Advances* 2, e2021AV000440. <https://doi.org/10.1029/2021AV000440>
- Karl, T.R., S. J. Hassol, C. D. Miller, W. L. Murray, eds. (2006). Temperature Trends in the Lower Atmosphere: Steps for Understanding and Reconciling Differences. U.S. Climate Change Science Program/Subcommittee on Global Change Research.
https://digital.library.unt.edu/ark:/67531/metadc12017/m2/1/high_res_d/sap1-1-final-all.pdf
- Klotzbach, P.J., R. A. Pielke, Sr., R. A. Pielke, Jr., J. R. Christy, R. T. McNider (2009). An alternative explanation for differential temperature trends at the surface and in the lower troposphere. *Journal of Geophysical Research — Atmospheres* 114(D21).
<https://doi.org/10.1029/2009JD011841>
- McKittrick, R. and J. R. Christy (2020). Pervasive warming bias in CMIP6 tropospheric layers. *Earth and Space Science* 7. <https://doi.org/10.1029/2020EA001281>
- McKittrick, R., and J.R. Christy (2025), “Data and Code for DoE Report 2025”, Mendeley Data, V1, doi: 10.17632/8n9ks93vn7.1
- Palmer, T., and B. Stevens (2019). The scientific challenge of understanding and estimating climate change. *Proceedings of the National Academy of Sciences* 116(49), 24390–24395.
<https://doi.org/10.1073/pnas.1906691116>
- Philipona, R., C. Mears, M. Fujiwara, *et al.* (2018). Radiosondes show that after decades of cooling, the lower stratosphere is now warming. *Journal of Geophysical Research—Atmospheres* 123.
<https://doi.org/10.1029/2018JD028901>
- Rugenstein, M., and M. Hakuba (2023). Connecting hemispheric asymmetries of planetary albedo and surface temperature. *Geophysical Research Letters* 50, e2022GL101802.
<https://doi.org/10.1029/2022GL101802>
- Santer B D, P.W. Thorne, L Haimberger *et al.* (2008) Consistency of modelled and observed temperature trends in the tropical troposphere *International Journal of Climatology* 28 1703–22
<https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/joc.1756>

- Santer, B.D. S. Po-Chedley, L. Zhao, C *et al.* (2023) Exceptional stratospheric contribution to human fingerprints on atmospheric temperature, *Proceedings of the National Academy of Sciences U.S.A.* 120 (20) e2300758120, <https://doi.org/10.1073/pnas.2300758120>.
- Scaffeta, N (2023) CMIP6 GCM ensemble members versus global surface temperatures. *Climate Dynamics* 60, 3091–3120 (2023). <https://doi.org/10.1007/s00382-022-06493-w>
- Seager, R., J. Feldman, N. Lis, *et al.* (2017). Whither the 100th meridian? The once and future physical and human geography of America's arid-humid divide. Part II: The meridian moves east. *Earth Interactions* 22. <https://doi.org/10.1175/EI-D-17-0012.1>
- Spencer, R. W. (2024). Global warming: Observations vs. climate models. Environment Backgrounder, The Heritage Foundation.
<https://www.heritage.org/environment/report/global-warming-observations-vs-climate-models>
- Stephens, G. L., D. O'Brien, P.J. Webster *et al.* (2015). The albedo of Earth. *Reviews of Geophysics*.
<https://doi.org/10.1002/2014RG000449>
- Vogelsang, T. and N. Nawaz (2016). Estimation and inference of linear trend slope ratios with an application to global temperature data: *Journal of Time Series Analysis* 38.
<https://doi.org/10.1111/jtsa.12209>

6 EXTREME WEATHER

Chapter Summary

Most types of extreme weather exhibit no statistically significant long-term trends over the available historical record. While there has been an increase in hot days in the U.S. since the 1950s, a point emphasized by AR6, numbers are still low relative to the 1920s and 1930s. Extreme convective storms, hurricanes, tornadoes, floods and droughts exhibit considerable natural variability, but long-term increases are not detected. Some increases in extreme precipitation events can be detected in some regions over short intervals but the trends do not persist over long periods and at the regional scale. Wildfires are not more common in the U.S. than they were in the 1980s. Burned area increased from the 1960s to the early 2000's, however it is low compared to the estimated natural baseline level. U.S. wildfire activity is strongly affected by forest management practices.

6.1 Introduction

High impact weather extremes, usually related to temperature, precipitation and/or high wind, can disrupt infrastructure and therefore endanger human health and wellbeing. The issue is not whether extremes occur. Rather, it is whether there are long-term (decadal scale) changes in the frequency or character of extremes (“detection”), as well as the extent to which such changes and the attendant changes in hazards are caused by anthropogenic emissions of greenhouse gases (“attribution”; e.g., AR6 Seneviratne *et al.* 2021).

It is naïve to assume that any recent extreme event is caused by human influences on the climate. Climate is about the statistical properties of weather over decades, not single events. Further, there are only about 130 years of reliable observational records that can be analyzed statistically. That brief interval does not begin to contain all the extreme events that the climate system can create on its own. Over geologic time the climate system has generated an (essentially) infinite variety of weather patterns and extremes that humans have never observed and thus are absent from the databases used to determine extreme statistics [see *Perils of short data records* below]. For that reason, attributing an extreme event unprecedented in the record requires assumptions about the magnitude of natural variations.

This chapter is concerned with detection of trends in extreme weather, while Chapter 8 considers causal attribution, with Section 8.4 specifically addressing extreme weather. If no trend is detected, then clearly there is no basis for attribution. But even where a trend is observed, attribution to human-caused warming does not necessarily follow. **Too often process-based reasoning is used without reference to observations...**

This is especially true for precipitation events. The hydrological literature has long noted the presence of long, slow and irregular oscillations in rainfall data (Hurst 1951, Cohn and Lins 2005, Markonis and Koutsoyiannis 2016). The characteristics of these natural patterns require long records to accurately estimate variability. Analysis of records that are short relative to the scale of natural variability will tend to misrepresent trends, therefore overstating the significance of apparent trends and underestimating the likelihood of extreme events (Cohn and Lins 2005).




Figure 6.1.1: The annual minimum depth of the Nile River near Cairo over more than 650 years from 622 to 1284 A.D. The data, measured in meters, shows a characteristic pattern of year-to-year fluctuations around longer-term trends. Data from Koutsoyiannis (2013)

A good example of this is the eight-century long record of the annual minimum height of the Nile River observed at Roda Island in Cairo shown in Figure 6.1.1. The Nile River is fed by precipitation over a 4 million square mile drainage basin, an area equal to about one third of CONUS. Since human influences on the global climate were negligible before the 20th century, the century-scale variability of the thirty-year average is entirely natural; Egyptians of the seventh and eighth centuries would have been incorrect to assume that the worsening drought during that time was the “new normal.”

With these caveats in mind, we examine the evidence for changes in selected weather and climate extremes. A recurring theme is the wide gap between public perceptions and scientific evidence. It has become routine in media coverage, government and private sector discussions, and even in some academic literature to make generalized assertions that extreme weather of all types is getting worse due to GHGs and “climate change.” Yet expert assessments typically have not drawn such sweeping conclusions and instead have emphasized the difficulty both of identifying specific trends and establishing a causal connection with anthropogenic forcing.

In the sections to follow we provide excerpts from various IPCC and NCA assessment reports denoting the sources as follows:

SREX: The IPCC Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation (2012)

AR6: The IPCC Sixth Assessment Report Working Group 1 (2021).

NCA4: The U.S. Climate Science Special Report of the Fourth National Climate Assessment (2017) Volume I.

NCA5: The U.S. Climate Science Special Report of the Fifth National Climate Assessment (2023) Volume I.

In the excerpts, *italics* are in the original whereas **boldface** emphasis has been added.

Additionally we present updated data series from standard government sources to provide up-to-date information through 2024 wherever possible.

6.2 Hurricanes and tropical cyclones

AR6 provides the following assessment of tropical cyclones (TCs; used here as a synonym for hurricanes):

AR6: There is *low confidence* in most reported long-term (multidecadal to centennial) trends in TC frequency or intensity-based metrics due to changes in the technology used to collect the best-track data. (IPCC, 2021 p. 1585)

AR6: It is *likely* that the global proportion of major (Category 3–5) tropical cyclone occurrence has increased over the last four decades . . . There is *low confidence* in long-term (multi-decadal to centennial) trends in the frequency of all-category tropical cyclones. (IPCC, 2023 SPM p. 9)

AR6: A subset of the best-track data corresponding to hurricanes that have directly impacted the United States since 1900 is considered to be reliable, and shows **no trend in the frequency of U.S. landfall events**. (IPCC 2021 p. 1585)

Since 1980, when satellite observations first fully covered the global oceans, we have confidence in the numbers of total global hurricanes and major hurricanes (Category 3 and higher). Figure 6.2.1 shows that on average, each year there are about 50 hurricanes, with about 25 reaching major hurricane status (Maue, 2025). There is substantial year-to-year and decadal variability, a weak decrease in the number of hurricanes, and a slight but insignificant increase in the number of major hurricanes. These two trends combine to create an overall increase in the proportion of major hurricanes.

Global hurricane statistics are dominated by the Northwest Pacific Ocean, which accounts for ~35 percent of total global hurricanes, whereas the Atlantic accounts for ~15 percent of global hurricanes (Colorado State University, 2025). Data in the Atlantic basin extends further back than in the other ocean basins and is most relevant to U.S. policy makers.




Figure 6.2.1: Global frequency of hurricanes and major hurricanes since 1980. Source: Updated from Maue 2011.

Figure 6.2.2 shows the frequency of Atlantic hurricanes and major hurricanes (Category 3 and higher) back to 1920. Data prior to 1965 (the onset of satellite observations in the Atlantic, shaded in blue) shows some undercounting, with data prior to 1920 showing substantial undercounting (Vecchi and Knutson, 2011). All measures of Atlantic hurricane activity show a significant increase since 1970. However, the period from 1971-1994 saw exceptionally low activity, with high activity (comparable to the past two decades, even with undercounting) also observed during the 1950's and 1960's, and even in the 1930's.




Figure 6.2.2: Atlantic frequency of hurricanes (HR) and major hurricanes (MHR) since 1920. Source National Hurricane Center (2024)

Figure 6.2.2 shows that Atlantic hurricanes vary strongly on decadal and multidecadal time scales. These variations are associated primarily with the Atlantic Multidecadal Oscillation (AMO), which is manifest in basin-wide sea surface temperature and sea level pressure fluctuations connected to large scale ocean circulation patterns. The AMO was in its warm phase during 1926-1970 and 1995-present, but in its cool phase during 1971-1994. It has its greatest impact on the number of major hurricanes (Category 3+), identified by Goldenberg *et al.* (2001) as associated with above normal SSTs and decreased vertical shear in the AMO warm phase (see also Bell and Chelliah, 2006; Klotzbach *et al.*, 2018).

Klotzbach *et al.* (2018) conducted a comprehensive evaluation of the landfalling hurricane data for the Continental U.S. since 1900. Figure 6.2.3 updates their analysis through 2024. While the largest numbers of landfalling hurricanes are from 2004, 2005 and 2020, there is no statistically significant trend since 1920. Figure 6.2.3 also shows the time series for major hurricane landfalls (Category 3-5). The largest year in the record is 2005, with 4 major hurricane landfalls. However, following 2005 there were no major hurricanes striking the U.S through 2016, the longest such period since 1920.




Figure 6.2.3: U.S. landfalling frequency of hurricanes (HR) and major hurricanes (MHR) since 1920. Source NOAA HRD(a) (2024)

Figure 6.2.3 shows substantial interannual to multidecadal variability in U.S. landfall activity. Klotzbach *et al.* (2018) examined how the landfall counts vary with ENSO (El Niño versus La Niña) and the warm versus cold phases of the Atlantic Multidecadal Oscillation (AMO).

Villarini *et al.* (2012) provide an analysis of U.S. hurricane landfalls back to 1878. While it is possible that some landfalls were missed in the late 19th century owing to sparsely populated regions on the Gulf Coast, it is remarkable that the highest year in the entire record is 1886, with 7 hurricane landfalls, when human influences on the climate were much smaller than they are today.

Table 6.2.1 shows the 10 strongest hurricanes (plus ties) to make U.S. landfall. Of the hurricanes that have made landfall with sustained winds greater than 150 mph, only one has occurred in the 21st century.

In summary, analyses of both global and regional variability and trends of hurricane activity provide the basis for detecting changes and understanding their causes. The relatively short historical record of hurricane activity, and the even shorter record from the satellite era, is not sufficient to assess whether recent hurricane activity is unusual relative to the background natural variability. Atlantic hurricane processes are influenced substantially by the natural modes of ocean circulation variability in the Atlantic, notably the Atlantic Multidecadal Oscillation. While it has long been hypothesized that a rising global sea surface temperature would cause an increase in hurricane intensity, identification of any significant trend in the hurricane data is hampered by a short data record and substantial natural variability.

Process-based understanding also suggests that storm surges and rainfall from hurricanes should be increasing with warmer temperatures. However, the relatively small number of hurricanes with varying landfall locations and the complex dynamics associated with each storm preclude meaningful detection of change.

Rank	Year	Landfall Wind (MPH)	Name
1	1935	185	"Labor Day"
2	1969	175	Camille
3	1992	165	Andrew
4	2018	160	Michael
5	1856	150	"Last Island"
5	1886	150	"Indianola"
5	1919	150	-----
5	1932	150	"Freeport"
5	2004	150	Charley
5	2020	150	Laura
5	2021	150	Ida
5	2022	150	Ian

Table 6.2.1 Strongest hurricanes to make landfall along the U.S. coast. Source (NOAA HRD(b), 2024)

6.3 Temperature extremes

The AR6 assessment focused on the period after 1950 and reported increasing trends in heatwave frequency and intensity. However, NCA4 noted that heatwave activity in the U.S. reached a peak in the 1930s (Figure 6.3.1).

AR6: It is *virtually certain* that hot extremes (including heatwaves) have become more frequent and more intense across most land regions since the 1950s, while cold extremes (including cold waves) have become less frequent and less severe (SPM, A3.1)

AR6: In North America, there is very robust evidence for a *very likely* increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes for the whole continent, though there are substantial spatial and seasonal variations in the trends. Minimum temperatures display warming consistently across the continent, while there are more contrasting trends in the annual maximum daily temperatures in parts of the USA. (Chapter 11, p 1550)

NCA4: Changes in warm extremes are more nuanced than changes in cold extremes. For instance, the warmest daily temperature of the year **increased** in some parts of the West over the past century, but there were **decreases** in almost all locations east of the Rocky Mountains. In fact, all eastern regions experienced a net **decrease**, most notably the Midwest (about 2.2°F [1.2°C]) and the Southeast (roughly 1.5°F [0.8°C]). (pp. 190-191)

NCA4: Since the mid-1960s, there has been only a very slight increase in the warmest daily temperature of the year (amidst large interannual variability). Heat waves (6-day periods with a maximum temperature above the 90th percentile for 1961–1990) increased in frequency until the mid-1930s, became considerably less common through the mid-1960s, and increased in frequency again thereafter. As with warm daily temperatures, **heat wave magnitude reached a maximum in the 1930s.** (pp. 190-191)




Figure 6.3.1: U.S. Heat waves since 1900. Source: NCA4 Figure 6.4

6.3.1 Temperatures in the U.S. are becoming less extreme

Daily maximum temperatures in the warm season (Tmax, May-Sep) and daily minimum temperatures in the cold season (Tmin, Dec-Mar) are available beginning in Dec 1898 (126 years). The dataset consists of 1,211 CONUS stations designated as United States Historical Climate Network or USHCN stations (see Figure 6.3.2; Quinlan *et al.* 1987, Karl *et al.* 1990). These stations were selected by NOAA as having the fewest problematic issues with gaps, station moves, and instrument changes. Where gaps still exist, nearby stations (bias-corrected) were merged so that the median volume of data available for a station is 98%. Although there are certainly errors in the dataset, including unresolved spurious warming due to UHI effects that especially bias Tmin records, this data set is sufficiently accurate for assessing trends in Tmax heat extremes.




Figure 6.3.2 Locations of USHCN temperature stations. Source: USHCN.

We begin with the question of whether the occurrence of daily record high or low temperatures has changed since Dec 1898. Each warm-season has 153 days (1 May to 30 Sep) and each cold-season has 122 days (1 Dec to 31 Mar). For each station and day, we calculated the year in which the record highest (lowest) temperature occurred. With 126 years of observations, if there were no temperature trends over time, the expected number of records for Tmax would be 1.21 ($=153/126$) per station per year and for Tmin 0.96 ($=122/126$) per station per year.

Figure 6.3.3 shows the observed distribution in time of the occurrence of these extreme events. There is a common feature in many metrics of warm-season extremes in the CONUS - the exceptional heat of the 1920s and especially the 1930s, peaking in 1936. On a per-station average, 60 percent of the Tmax records and 59 percent of the Tmin records occurred in the first half of the period (1899-1961).




Figure 6.3.3 Number of daily record High (red) and Low (blue) temperatures for warm and cold seasons in the CONUS. The lines represent the 15-year running, centered average. Source: 1,211 USHCN stations supplemented as needed to achieve a minimum of 92 percent of observations in the 126-year period since Dec 1898. US48: contiguous U.S. states. Tmax: maximum temperature. Tmin: minimum temperature.

On the cold side, the Valentine's Day Arctic outbreak in Feb 1899 stands as the most extensive cold extreme experienced by CONUS, with 1917 in 2nd place. The frequency of cold records has declined, especially over the last quarter of the period in which only 13 percent of the extreme cold events were measured. In contrast, 25 percent of the extreme Tmax records were achieved in the last quarter, in accordance with statistical expectations. These general features have been noted in past assessments (see above, IPCC AR6, NCA4). Combining the two histories, the overall reduction in numbers of both cold and hot extremes over the past century indicates a climate less prone to extremes.

This pattern is also shown in Figure 6.3.4. For each station and for each year the hottest warm season and coldest cold-season temperatures were calculated. Then the differences between these were computed by station and geographically-averaged over all stations, thus yielding an annual measure of the expected range of local extreme temperatures for each year. Figure 6.3.4 shows the 15-year trailing average of this measure, which has clearly declined over the past century.




Figure 6.3.4. 15-year trailing average of the difference each year of each station's hottest warm-season Tmax and coldest cold-season Tmin relative to the long-term average. Source: Author analysis of USHCN data.

The average difference for each station between the hottest summer Tmax and coldest winter Tmin has declined by about 5°F in the past 126 years. The decline is due mostly to warmer winter Tmin, but a decline in summer Tmax is also a factor. The rise in Tmin has been strongly related to the growing presence of manufactured surfaces around the weather stations over the last 100+ years (the so-called urban heat island effect; Section 3.3 and Spencer *et al.* 2025). **Need to add citations about UHI and Tmin**

In summary, while temperature extremes are regularly experienced in the U.S. and attract a great deal of media attention, long term records show the U.S. climate has become less extreme over time (milder) when measured by the range between warm season maxima and cold season minima.

6.3.2 Exceedances of a heat threshold

Under the heading of “The Risk of Temperature Extremes is Changing”, the most recent U.S. National Climate Assessment report (NCA5) notes the increase in a threshold metric of number of days at or above 95°F, stating,

The western U.S. has been particularly affected by extreme heat since the 1980s ..., experiencing a larger increase in days over 95°F, as would be expected given the greater warming in that region relative to the eastern US. Several major heatwaves have affected the U.S. since 2018, including a record-shattering event in the Pacific Northwest in 2021.

Are the occurrences of 95°F days changing? In a climate as varied as that of CONUS, threshold statistics can be misleading. A region with many stations that have near 95°F Tmax average temperatures in the summer might see large swings in the metric when only small changes in average temperature occur. Elsewhere with stations that either rarely or virtually always achieve 95°F Tmax temperatures, a small change will not have much impact on the results.




Figure 6.3.5. Total number of days $\geq 95^{\circ}\text{F}$ in 6-yr periods, U.S. 48 (bars) and regions (lines). A 6-year period is used as this evenly divides the 126-year record. Results are robust to using periods from 2 to 11 days. US48: contiguous U.S. states. See Figure 6.3.2 for region names. Source: Author analysis of USHCN data.

In the past 126 years, the average CONUS station experienced 129 days exceeding 95°C per 6-yr period, but the regional values range from 278 in the Southern Plains to 9 in the Northeast. Thus, such threshold analyses must be interpreted with caution. Figure 6.3.5 shows that only three of the nine regions, all in the West, have experienced upward trends in the number of 95°F or hotter days (dashed lines). The CONUS as a whole has not, and the other six regions have experienced declines.

The Pacific NW heatwave of 2021 referenced in the NCA5 quote will be examined more closely in Section 8.6.1. The evidence indicates that it was a single, unprecedented event in the record, not part of a pattern of increasing extreme heat. For example, the 5-day average tropospheric grid-point temperature anomaly over the Pacific NW during that event was $+10.8^{\circ}\text{C}$, the most extreme Northern Hemisphere grid point summer anomaly in the 46 years from over 4 million grid values. In contrast, the global temperature anomaly during that time was virtually zero ($+0.03^{\circ}\text{C}$, Mass *et al.* 2024).

6.3.3 Heatwaves

Heatwaves (consecutive days that exceed an extreme threshold) have a greater societal impact than a single daily record temperature. We measure “Heatwave Days” here as the count of all days in May-Sep each year that exceed the 90th percentile for that day and that lie within a period of at least six consecutive days. This is equivalent to the method used in NCA4, except that the reference period here is the entire record 1899-2024 while NCA4 truncated the reference period to 1961-1990, which was a cool interval. (The pattern of results shown below does not depend on the choice of reference period.) That truncation boosts positive results (days exceeding the 90th percentile) in years warmer than the reference period, especially starting in 1960 and moving to the present (see Figure 6.3.6 and discussion below).




Figure 6.3.6 15-year trailing average of number of heatwave days per year per station in the CONUS (black line) and two regions: West (red), Central-east (green).

Figure 6.3.6 indicates that there are regional variations in heatwave activity. The excessive heat of the first half of the 20th century occurred primarily in the eastern two-thirds of the nation, while the West has seen a recent increase of heatwave days (NCA5). This indicates that the background warm season circulation favored heatwaves in the eastern portions of the country in the first half of the 20th century, but in the 21st century the patterns have favored heatwaves in the West. For CONUS as a whole, heatwaves are no more common today than they were a century ago, consistent with the upper panel of our Figure 6.4.1 taken from NCA4.

This metric varies significantly with region. The four northern regions (Pacific NW, Northern Plains, Upper Midwest, and Northeast) on average experience 15 to 27 heatwave days per 15-year period. In contrast, the five southern regions (Pacific Southwest, 4-Corners, Southern Plains, Ohio Valley and Southeast) see 37 to 54 such days, essentially twice as many. This suggests the summer circulation pattern is more prone to stationary events in the southern regions while transient systems in the northern regions are more common and thus cut short these potentially longer events.

The analysis of heatwaves is an example of why it is important to consider complete datasets and appropriate metrics. The NCA5 directs readers to the website <https://www.globalchange.gov/indicators/heat-waves> (USGCRP 2023) to view a figure showing the number of urban heatwaves by decade from the 1960s, which we reproduce as Figure 6.3.7.




Figure 6.3.7: Average number of urban heatwaves per year for 50 large U.S. metropolitan areas, a misleading metric for reasons explained in the text. From <https://www.globalchange.gov/indicators/heat-waves> (accessed May 22, 2025).

The figure shows a monotonic increase in each decade from two occurrences per year in the 1960s to six in the 2020s. The definition of a heatwave used is an unusual but practical measure of human discomfort - a period of at least 2 consecutive days when the *minimum* apparent temperature (combination of temperature and humidity) exceeds the 85th percentile. Note too, the dataset is limited to the 50 largest U.S. cities.

Given the unusual heatwave definition and urban focus, these increasing values since 1960 presented in USGCRP (2023) are not informative about long term trends or the influence of GHG emissions for at least two reasons. First, as shown in Figures 6.3.1 and 6.3.6, the 1960s was the coldest decade and 1970s the second coldest decade since the 1910's, so this starting date preconditions the time series to show increases. Second, post-1960 urbanization in these cities is a major factor in the rise of Tmin relative to co-located Tmax and relative to Tmin at nearby rural stations (Karl *et al.* 1988, Runnals and Oke 2006, Christy *et al.* 2009, McNider *et al.* 2012). This does not dismiss the real rise in nighttime temperatures in major U. S. cities and the societal impacts associated with these changes. However, we note for a variety of reasons that summer Tmax (especially in rural areas) is a better metric for detecting changes in heatwaves influenced by changes in the background climate due, for example, to increasing GHGs (Christy *et al.* 2009). For CONUS as a whole, the evidence in this section suggests GHG emissions have had little-to-no effect on heatwaves against the background of urbanization and natural climate variability. Irrespective of the ultimate cause of regional trends, heatwaves have important effects on society which must be addressed, as we discuss in Chapter 10.

BOX: Perils of short data records

San Francisco provides a good case study of the limitation of using short historical samples to characterize natural variability of extreme events. Suppose we use a 130-year sample of daily San Francisco precipitation from 1895 to 2024 and we look for 3-day, 5-day, 14-day and 30-day rainfall records. The results are as shown in Table 7.2.

Event	Record (inches)	Year
3-day	6.94	2023
5-day	8.55	2023
14-day	12.62	2023
30-day	18.93	1998

Table 6.2: Extreme rainfall records, San Francisco, 1895-2024.

The records all cluster in the more recent years. 2023 appears to be an exceptional year and since it is near the end of the sample, it might suggest that the climate has shifted into a more hazardous state, perhaps because of human influences. [Note “2023” indicates the event occurred in the water-year of Aug 2022 to Jul 2023.]

But the picture is very different if we use a sample that begins 45 years earlier, in 1850. Table 6.3 shows that the record-setting events all happened in the 1860s. Furthermore 2023 is now not even in 2nd place but falls to 3rd or 4th place. And comparing the records of the two charts shows the extreme precipitation events in 1862 and 1867 involved considerably more rainfall than the 1998 and 2023 events, with 14- and 30-day totals about 50 percent higher.

Event	Record (inches)	Year	Rank of 130-yr extreme listed in Table 6.2
3-day	8.85	1867	3
5-day	9.80	1867	3
14-day	19.05	1862	4
30-day	28.25	1862	2

Table 6.3: Extreme rainfall records, San Francisco, 1850-2024.

The range of natural variability is made even more remarkable when paleoclimate evidence is examined. Porter *et al.* (2011) discovered that in the past 1,800 years at least six megastorms were more intense than the devastating 1861-62 ARkStorm that struck the region. Evidently such extreme events, “unprecedented” in our 1895-2024 sample, have impacted the region about every 300 years, though not since 1895.

This example illustrates the limitations of using relatively short climate periods (~130 years) to assess the character and range of natural variability in general and of extreme events in particular. Accurate representation of the full range of natural variability is necessary for any attribution analyses (Section 8.6). Infrastructure planners, emergency management institutions and attribution scientists would understand the significant mischaracterization of the magnitude of a future extreme if based only on the last 130 years. In this case, a single time-sample of 130 years provides an underestimation of the extreme value by up to 50 percent determined when adding only 45 more years of observations. Compared to millennial-scale paleoclimate evidence, an even greater underestimation would occur. An important lesson is that the climate can deliver great surprises on its own, even without human influences.

6.4 Extreme precipitation

AR6 assessed that an increase in heavy precipitation has been observed in data starting in the 1950s.

AR6: The frequency and intensity of **heavy precipitation events have increased** since the 1950s over most land area for which observational data are sufficient for trend analysis (high confidence). (SPM A3.2)

AR6: In North America, there is robust evidence that the **magnitude and intensity of extreme precipitation has *very likely* increased** since the 1950s. Both [one-day maxima] and [5-day maxima] have significantly increased in North America during 1950-2018. (Chapter 11, p. 1560)

The U.S. National Climate Assessments (NCA4, NCA5) have highlighted an increase in the occurrence of the heaviest precipitation events (defined in different ways) primarily in the eastern half of CONUS, especially the Northeast, when starting the analysis in either 1901 or 1958. Interestingly, the regional variations indicate that the largest increases in extreme precipitation events are in the Northeast and the smallest in the West, a pattern counter to the changes in temperature extremes (Figure 6.3.6).

McKittrick and Christy (2019) examined long-term and consistent station observations of extreme daily precipitation to test some of these NCA claims for the Southeast and West Coast using a trend model with a non-parametric variance estimator robust to the complex autocorrelation properties of precipitation data. When the time series were extended back in time (as far as 1872 in some cases) or started later (1978), there were no significant trends for either region.

These findings have been updated for this report (McKittrick and Christy 2025) with similarly constructed observations from 29 stations on the CONUS Pacific Coast (1893ff from San Diego CA to Blaine WA) and 24 stations in the humid Southeast (1872ff from Austin TX to Washington DC), also adding 27 stations in the Northeast (1888ff from Buffalo NY to Eastport ME). The locations are shown in Figure 6.4.1. The stations were selected based on availability of long-term high-quality records. The regions are each associated with important features of extreme precipitation behavior: the Pacific coast is associated with landfalling atmospheric rivers for which AR6 cites evidence of increasing activity since 1948 with further increases expected as the world warms (AR6 8.3.2.8.2); the NCA report indicates that the Northeast has experienced the greatest increase in extreme events, and the Southeast is also a place noted in the NCAs as having increased extreme events.

The results of applying the analysis of McKittrick and Christy (2019) were as follows for each region, then followed by further explanation.

Pacific Coast heavy rainfall events

- The average precipitation trend is statistically significant (downwards) in Astoria OR; insignificant elsewhere.
- The trend in rainfall variance is positive and significant in Big Sur CA; insignificant elsewhere.
- The trend in daily maximum precipitation is positive and significant in Aberdeen WA and Big Sur CA and negative and significant in Newport OR (insignificant elsewhere).

- Averaged over all stations in the region, none of these three trend parameters is statistically significant.

The Pacific Coast receives considerable precipitation from Atmospheric River (AR) events which often last more than a day or two (e.g., Gershunov *et al.* 2017, Pan *et al.* 2024). The worst series of such events in recent history was the so-called ARkStorm that occurred during December 1861 and January 1862; it dumped nearly 10 feet of rain in parts of California and submerged the entire Central Valley for weeks under as much as 15 feet of water (Brewer 1930, Null and Hulbert 2007). Additionally, Paleoclimate research has found six megastorms more severe than 1861–1862 in California during the last 1800 years, occurring at intervals of 300 years or so (Porter *et al.* 2011).



Figure 6.4.1: Locations of precipitation monitoring stations used in this report. Orange: Pacific coast. Blue: Northeast. Green: Southeast. Data from McKittrick and Christy (2025).

We examine occurrences of 5-day deluges as follows. Taking the Pacific coast as an example, a 130-year span contains 26 5-year intervals. At each location we computed the 5-day precipitation totals throughout the year and selected the 26 highest values across the sample. A single year might have more than one of the 26 heaviest. Each of those can be thought of as 1-in-5yr events. If there are no trends in precipitation, then the total number of these events across all stations should be evenly spread over the years. In Fig. 6.4.2 we show the distribution in time of these events for the Pacific coast. The deluges associated with the massive 1997-98 El Niño event are readily apparent. While erratic, as is typical of such precipitation metrics, there is no indication of a tendency to become more frequent over time.




Figure 6.4.2. The time distribution by 5-year periods of the 26 heaviest (1-in-5 yr) occurrences for 29 stations on the Pacific coast.




Figure 6.4.3. As in Fig. 6.4.2 but for the heaviest 30 (1-in-5yr) events for 24 stations in the humid Southeast from Austin TX to Washington DC in 5-year bins for 1875-2024.

Southeast heavy rainfall events

- The trend in average precipitation is positive and statistically significant in Mobile AL and Quitman GA but insignificant elsewhere.
- The trend in rainfall variance is positive and significant in Mobile AL but insignificant elsewhere.
- The trend in daily maximum precipitation is positive and significant in Vicksburg MS and Norfolk VA but insignificant elsewhere.
- Averaged over all stations in the region none of these three trend parameters is statistically significant

Figure 6.4.3 is analogous to Figure 6.4.2 for the last 150 years in the Southeast humid zone. The temporal pattern of 5-day totals of the 1-in-5yr heavy events is generally unremarkable, though a cluster of higher values appears in 1995 to 2019. The increase in those years is due largely to the 4 northeastern-most stations of Wilmington NC, Weldon NC, Washington DC, and Norfolk VA. This confirms the pattern indicated in NCA4 and NCA5 -- an increasing frequency of heavy events due to a temporal clustering of tropical storms from eastern NC to Maine discussed below. Otherwise, the remaining 20 stations show an unremarkable temporal distribution of heavy events.

Northeast heavy rainfall events

- The trend in average precipitation is positive and statistically significant in 12 of 27 locations and also in the regional average.
- The trend in rainfall variance is positive and significant in Portland ME, Albany NY, Buffalo NY and Eastport ME but insignificant elsewhere.
- The trend in daily maximum precipitation is positive and significant in Portland ME, Gardiner ME and Eastport ME but insignificant elsewhere.
- When averaged over all stations in the region, there is no statistically significant trend in either the precipitation variance or maximum

Fig. 6.4.4 is analogous to Figure 6.4.2 for the last 135 years in 27 stations in the Northeast (including Montreal Canada). We use 3-day totals here as this produced the largest temporal variations in time. In this region, 77 percent of events occur during June to October and are dominated by incursions of hurricanes, tropical storms, or tropical storms that transition to extratropical systems. According to NCA4 and NCA5 this region experienced the largest increases in extreme events, so it merits a closer examination.

There is a noticeable clustering of extreme events from 1995 to 2014. Howarth *et al.* (2019) examined a similar region as in Fig. 6.4.4 that includes PA and NJ and reported significant differences in various precipitation extremes between two 18-year periods, 1979-96 and 1997-2014. That included a 317 percent increase in 24-hr events exceeding 6 inches, while we find a 58 percent increase over the same years. However, Figure 6.4.4 shows that frequency drops sharply after 2014, returning to the long-term average in the subsequent 5-year intervals, again illustrating the perils of drawing conclusions from short-term trends in highly variable metrics.




Figure 6.4.4. As in Fig. 6.4.2 but for the heaviest 27 (1-in-5yr) 3-day precipitation events for 27 Northeast stations from NY to ME, including Montreal.




Figure 6.4.5. Average amount of precipitation falling in the 1-in-5yr events for the NE stations.

The high percentage increase in the Howarth *et al.* sample is associated with small numbers at relatively few locations: there were only 6 in the first period and 25 in the second across 58 stations. Most of the stations did not record such an event. If there is a region-wide increase in heavy events, it should be seen in the average across all stations. Figure 6.4.5 shows the average precipitation in the 1-in-5 yr events across the NE. The trend is only +0.04 inches/decade. The highest amount, 1935-39, includes the Great New England Hurricane of 1938, one of the rare major (Category 3 or higher) hurricanes to strike the region.

The results in Figs. 6.4.4 and 6.4.5 thus suggest that though there was a surge in the number of events at a few locations during the 1995 to 2014 interval, there was no regional pattern and the change did not persist beyond 2014.

Jong *et al.* (2024) document the increase of tropical influences on precipitation events in the Northeast since 1959 and concluded “The autumn extreme precipitation trend over the Northeast U.S. is primarily attributed to tropical cyclone-related events since the 1990s.” The question then becomes: “Was the temporal clustering of tropical systems in 1997-2014 which affected the Northeast a response to increasing GHGs?” Jong *et al.* examined CMIP-6 model output which suggests that there will be fewer such systems in the 21st century but that the intensity of the rainfall events might increase. This conjecture is not seen in Fig. 6.4.5 where the amount-per-event has remained steady over the 135-year period.

There is some evidence to indicate the heaviest rainfall events might be redistributed due to the impact of urban infrastructure on the local weather (e.g., Pielke Sr. *et al.* 2011, Zhang *et al.* 2018, Yang *et al.* 2024). Yang *et al.* state “Cities that experience compact development tend to witness larger increases in extreme rainfall frequency over downtown than their rural surroundings, while the anomalies in extreme rainfall frequency diminish for cities with dispersed development.” While this is an important insight to consider, the effect on the specific stations used in this analysis is unknown, or at least not detectable in Fig. 6.4.5.

In summary, some U.S. regions show short-duration increases in extreme precipitation events, consistent with natural variability. But analysis of long term, nationwide historical records that considers the autocorrelation properties of precipitation data does not support the claim that extreme short-duration rainfall events are becoming more frequent or intense.

6.5 Tornadoes

AR6 assesses tornado trends in the U.S. as follows:

[O]bservational trends in tornadoes, hail, and lightning associated with severe convective storms are not robustly detected due to insufficient coverage of the long-term observations. There is *medium confidence* that the mean annual number of tornadoes in the USA has remained relatively constant. (Chapter 11, section 11.7.3, p. 1594)

The monitoring of weak tornadoes has changed over time. The growth of rural populations and the increasing ability to take video with hand-held devices has led to more frequent reports of weak tornadoes that produce minimal damage. In contrast, strong to violent tornadoes have been observed more consistently over time. Note that tornado strength is measured by the damage it produces, not by the visual appearance of the funnel. Limited real-time observational capabilities in earlier decades did not prevent identification because strong to violent tornadoes leave much more damage which will be assessed later even if the tornado itself was not observed. Since statistics began in 1950, there has been a substantial decrease (by about 50%) in the number of strong to violent tornadoes as shown in Fig. 6.5.1a.

To summarize, there is a noticeable downward trend in the number of severe tornadoes in the U.S. since 1950. After 1990 the number of weak tornadoes in the U.S. has remained roughly constant; data before that are incomplete due to limited monitoring.




Figure 6.5.1. Annual U.S. tornado counts for (a) strong to violent tornadoes (EF3 to EF5), and (b) weak tornadoes (EF0 to EF2). Based upon NOAA Storms Prediction Center data, available at https://www.spc.noaa.gov/wcm/data/1950-2024_actual_tornadoes.csv

6.6 Flooding

Changes in floods were assessed as follows:

AR6: The SREX assessed *low confidence* for observed changes in the magnitude or frequency of floods at the global scale. This assessment was confirmed by the AR5 report. The SR15 found increases in flood frequency and extreme streamflow in some regions, but decreases in other regions. . . [H]ydrological literature on observed flood changes is heterogeneous, focusing at regional and sub-regional basin scales, making it difficult to synthesise at the global and sometimes regional scales. (Chapter 11.5)

AR6: [T]he seasonality of floods has changed in cold regions where snowmelt dominates the flow regime in response to warming (*high confidence*). Confidence about peak flow trends over past decades on the global scale is *low*. (Chapter 11.5)

NCA4: Trends in extreme high values of streamflow are **mixed** across the United States. Analysis of 200 U.S. stream gauges indicates areas of both increasing and decreasing flooding magnitude **but does not provide robust evidence** that these trends are attributable to human influences (pp. 240-241)

The absence of detectable US-wide trends in flooding is consistent with the findings in Section 6.4 of absence of coherent changes in extreme precipitation.

6.7 Droughts

Assessments of drought trends were as follows.

AR6: Few AR6 regions show observed increases in meteorological drought (Section 11.9, p. 1575),

AR6: Increasing trends in agricultural and ecological droughts have been observed on all continents (*medium confidence*), but decreases only in one AR6 region (*medium confidence*). Increasing trends in hydrological droughts have been observed in a few AR6 regions. (Chapter 11 Summary)

NCA4: As a consequence of this increased precipitation, **drought statistics over the entire CONUS have declined.** (p. 233)

NCA4: Recent droughts and associated heat waves have reached record intensity in some regions of the United States; however, by geographical scale and duration, the Dust Bowl era of the **1930s remains the benchmark** drought and extreme heat event in the historical record (very high confidence). (p.231)

SREX: From a paleoclimate perspective, **recent droughts are not unprecedented**, with severe ‘megadroughts’ reported in the paleoclimatic record for Europe, North America, and Australia. (p. 170)




Figure 6.7.1: Monthly percent of US classified as “Very Dry” 1895—2025. Data source: NOAA <https://www.ncei.noaa.gov/access/monitoring/uspa/wet-dry/0> least squares trendline added.

As shown in Figure 6.7.1, U.S. long-term data shows an insignificantly declining trend in extreme dryness (-0.001 percent per year)

Kogan *et al.* (2020) examines a 38-year high-resolution satellite-based drought measure and concludes that global drought has not intensified and is not connected to climate change: “it is possible to state firmly that global and main grain countries’ drought area and intensity trends have not been following global climate warming since 1980’s.”

In summary there is no evidence of increasing meteorological drought frequency or intensity in the U.S. or globally over recent decades.

6.8 Wildfires

The IPCC has not provided an attribution assessment of wildfires. As shown in Figure 6.8.1, global wildfire activity as measured by European Space Agency shows a downward trend in the 21st century.




Figure 6.8.1: Global wildfire area 2001-2018. Source: From Lizundia-Loiola *et al.* (2021) Figure 12. The different coloured lines represent data products derived from different satellites and algorithms

Global data show that wildfire coverage is constant or declining on every continent (Samborska and Ritchie, 2024). However there is evidence that the intensity of fires in some regions is worsening (Cunningham *et al.* 2024) and that wildfires resulted in a net loss of global forest cover over 2001-2019 (Tyukavina *et al.* 2022).

Active fire suppression since 1900 makes it difficult to establish a natural baseline for wildfire activity in the U.S. Paleoclimatic evidence indicates that past activity was much higher than today. Marlon *et al.* (2012) used sedimentary charcoal layers to reconstruct fire history of western U.S. for the past 1400 years and also fit a model to predict fire activity as a function of climatic conditions. Their results are summarized in Figure 6.8.2 below (from Figure 2 in their paper). There has been a growing wildfire deficit over the 20th century. In other words, however much fire was observed in the 20th century, it was less than what would have been observed in previous centuries based on the climatic conditions. Parks *et al.* (2025) likewise find that despite the recent increase in wildfire burn area in North America, a significant wildfire deficit remains relative to historical wildfire regimes.




Figure 6.8.2 Fire frequency and fire deficit in the US. The red line shows the smoothed charcoal record and the black dotted line shows the predicted charcoal record based climatic conditions. Source: Marlon *et al.* (2012) Figure 2.

U.S. data from the National Interagency Fire Centre (NIFC) from 1926 to 2023 are shown in Figure 6.8.3. The NIFC has removed the pre-1960 data from its current website on the grounds that measurement methods changed after 1960 making the comparison unreliable. Nonetheless just focusing on the post-1985 interval the number of fires is not increasing. The area burned did increase but only until about 2007.

Forest fires have always been a part of nature, and they can certainly create conditions that are inhospitable in the short term for all life, including humans. Science has confirmed the overall benefit and necessity of forest fires. While recent high-profile fires and seasons serve as a reminder of the potential destructive impact, the highest profile U.S. forest fire remains the 1910 Big Blowup fire in the U.S. west, which destroyed over three million acres and eliminated entire towns like Taft, MT (Apple, 2020). The 1910 fire reshaped the U.S. Forest Service (National Forest Foundation 2022) leading to a focus on fire suppression with a primary goal of defeating all forest fires (Forest History Society, 2022). This led to the “10 am rule” in 1935 requiring that all fires spotted on any day had to be controlled by 10 am the following day (National Forest Foundation, 2022).

While defeating all fires seemed a noble goal, questions began to arise as to whether this behavior “followed the science” (U.S. Forest Service, 2022). Over time the U.S. Forest Service has begun to rethink its goals, recognizing that new approaches such as prescribed burns, fuel elimination, and controlled wildfires are more appropriate (Sommer, 2016). Recent research is validating this approach and recognizing that more frequent smaller fires likely result in healthier forests, water ecosystems and biodiversity (Stephens *et al.*, 2021).




Figure 6.8.3: U.S. wildfires 1926 to 2023. Source: Post-2018: National InterAgency Fire Center data <https://www.nifc.gov/fire-information/statistics/wildfires>. Pre-2017 webarchive.org (n.d.).

References

- Apple, C. (2020, January 21). 1910 fire – The big burn across Montana and Idaho. *The Spokesman-Review*. <https://www.spokesman.com/stories/2020/jan/19/1910-fire-big-burn-across-montana-and-idaho/>
- Bell, Gerald D. and Muthuvel Chelliah, “Leading Tropical Modes Associated with Interannual and Multidecadal Fluctuations in North Atlantic Hurricane Activity,” *Journal of Climate* 19, no. 4 (February 15, 2006): 590–612, <https://doi.org/10.1175/JCLI3659.1>.
- Brewer, W. H. (1930). *Up and down California in 1860–1864: The journal of William H. Brewer*. University of California Press.
- Christy, J. R., Norris, W. B., and McNider, R. T. (2009). Surface temperature variations in East Africa and possible causes. *Journal of Climate*, 22(12), 3342–3356. <https://doi.org/10.1175/2008JCLI2726.1>
- Cohn, T. A. and Lins, H. F. (2005). Nature’s style: Naturally trendy. *Geophysical Research Letters*, 32(23). <https://doi.org/10.1029/2005GL024476>
- Colorado State University. (2025). *Realtime tropical cyclone guidance*. <https://tropical.atmos.colostate.edu/Realtime/index.php>
- Cunningham, C.X., G.J. Williamson, and D. Bowman (2024) Increasing frequency and intensity of the most extreme wildfires on Earth. *Nature Ecology and Evolution* 8, 1420–1425. <https://doi.org/10.1038/s41559-024-02452-2>
- Forest History Society (2022). *The 1910 fires*. <https://foresthistor.org/research-explore/us-forest-service-history/policy-and-law/fire-u-s-forest-service/famous-fires/the-1910-fires/>
- Gershunov, A., and K. Guirguis (2012). California heat waves in the present and future. *Geophysical Research Letters*, 39(18). <https://doi.org/10.1029/2012GL052979>
- Gershunov, A., T. Shulgina, F.M. Ralph, D.A. Lavers J.J. and Rutz (2017). Assessing the climate-scale variability of atmospheric rivers affecting western North America. *Geophysical Research Letters*, 44(15), 7900–7908. <https://doi.org/10.1002/2017GL074175>
- Goldenberg, Stanley *et al.*, “The Recent Increase in Atlantic Hurricane Activity: Causes and Implications,” *Science* 293, no. 5529 (July 20, 2001): 474–479, <https://doi.org/10.1126/science.1060040>.

- Howarth, M. E., C.D. Thorncroft, and L.F. Bosart (2019). Changes in extreme precipitation in the Northeast United States: 1979–2014. *Journal of Hydrometeorology*, 20(4), 673–689.
<https://doi.org/10.1175/JHM-D-18-0155.1>
- Hurst, H. E. (1951). Long term storage capacities of reservoirs. *Transactions of the American Society of Civil Engineers*, 116, 776–808.
- Intergovernmental Panel on Climate Change (IPCC). (2021). *Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (V. Masson-Delmotte *et al.*, Eds.). Cambridge University Press.
<https://www.ipcc.ch/report/ar6/wg1/>
- Jong, B.-T., H. Murakami, T.L. Delworth and W.F. Cooke (2024). Contributions of tropical cyclones and atmospheric rivers to extreme precipitation trends over the Northeast US. *Earth's Future*, 12, e2023EF004370. <https://doi.org/10.1029/2023EF004370>
- Karl, T.R., H.F. Diaz, and G. Kukla. (1988): Urbanization: Its detection and effect on the United States climate record. *Journal of Climate*, 1, 1099–1123
https://journals.ametsoc.org/view/journals/clim/1/11/1520-0442_1988_001_1099_uidaei_2_0_co_2.xml
- Karl, T. R., C.N. Williams Jr., F.T. Quinlan, and T.A. Boden (1990). *United States Historical Climatology Network (HCN) serial temperature and precipitation data* (Publication No. 3404). Oak Ridge National Laboratory.
- Klotzbach, Phil *et al.* (2018) Continental U.S. Hurricane Landfall Frequency and Associated Damage: Observations and Future Risks, *Bulletin of the American Meteorological Society* 99, no. 7 (July 2018): 1359–1376, <https://doi.org/10.1175/BAMS-D-17-0184.1>.
- Kogan, Feliz, Wei Guob and Wenze Yang (2020) “Near 40-year drought trend during 1981–2019 earth warming and food security” *Geomatics, Natural Hazards and Risk* Vol 11(1)
- Koutsoyiannis, D. (2013) Hydrology and Change. *Hydrological Sciences Journal* 58(6)
<http://dx.doi.org/10.1080/02626667.2013.804626>
- Lizundia-Loiola, J., *et al.* (2020) A spatio-temporal active-fire clustering approach for global burned area mapping at 250 m from MODIS data. *Remote Sensing of the Environment* 236, 111493,
<https://doi.org/10.1016/j.rse.2019.111493>
- Markonis, Y., and D. Koutsoyiannis (2016). Scale-dependence of persistence in precipitation records. *Nature Climate Change*, 6, 399–401. <https://doi.org/10.1038/nclimate2894>
- Marlon, Jennifer, Patrick J Bartlein, Daniel G. Gavin *et al.* (2012) Long-term perspective on wildfires in the western USA *Proceedings of the National Academy of Sciences* February 14 2012
<https://www.pnas.org/doi/pdf/10.1073/pnas.1112839109>
- Maue, R. N. (2025). *Climatlas: Tropical cyclones*. <https://climatlas.com/tropical/>
- Maue, R. N. (2011) Recent historically low tropical cyclone activity. *Geophysical Research Letters* 38, L14803, doi:10.1029/2011GL047711
- McKittrick, R., and J.R. Christy (2019). Assessing changes in US regional precipitation on multiple time scales. *Journal of Hydrology*, 578, 124074. <https://doi.org/10.1016/j.jhydrol.2019.124074>
- McKittrick, R. and J.R. Christy (2025), “Data and Code for CWG2025 Report”, Mendeley Data, V1, doi: 10.17632/by76s7rxfb.1
- McNider, R.T., G.J. Steeneveld, A.A.M. Holtslag, R.A. Pielke Sr., S. Mackaro, A. Pour-Biazar, J. Walters, U. Nair, and J.R. Christy, 2012: Response and sensitivity of the nocturnal boundary layer over land to added longwave radiative forcing. *Journal of Geophysical Research*, 117, D14106, doi:10.1029/2012JD017578.
- Null, J. and J. Hulbert, “California Washed Away: The Great Flood of 1862,” *Weatherwise* 60, no. 1 (2007): 26–30, <https://doi.org/10.3200/wewi.60.1.26-30>
- National Forest Foundation. (2022). *Blazing battles: The 1910 fire and its legacy*.
<https://www.nationalforests.org/our-forests/your-national-forests-magazine/blazing-battles-the-1910-fire-and-its-legacy>

- National Interagency Fire Center. (2024). *Wildfire statistics*. <https://www.nifc.gov/fire-information/statistics/wildfires>
- NCA4. (2017). *Climate science special report: Fourth National Climate Assessment, Volume I* (D. J. Wuebbles, D. W. Fahey, and K. A. Hibbard, Eds.). U.S. Global Change Research Program. <https://doi.org/10.7930/J0J964J6>
- NCA5. (2023). *Fifth National Climate Assessment* (A. R. Crimmins *et al.*, Eds.). U.S. Global Change Research Program. <https://doi.org/10.7930/NCA5.2023.CH1> (accessed May 22, 2025)
- NOAA Hurricane Research Division. (2025a). *All U.S. hurricanes (1851–2019)*. https://www.aoml.noaa.gov/hrd/hurdat/All_U.S._Hurricanes.html
- NOAA Hurricane Research Division. (2025b). *Most intense (3, 4, 5) continental United States hurricanes: 1851–1970, and 1983–2023*. https://www.aoml.noaa.gov/hrd/hurdat/most_intense.html
- Overland, J. E. (2021). Causes of the record-breaking Pacific Northwest heatwave, late June 2021. *Atmosphere*, 12(11), 1434. <https://doi.org/10.3390/atmos12111434>
- Pan, M., M. Lu and U. Lall (2024). Diversity of cross-Pacific atmospheric river main routes. *Communications Earth and Environment*, 5, 378. <https://www.nature.com/articles/s43247-024-01552-y>
- Parks, Sean, Christopher Guiterman, Ellis Margolis *et al.* (2025) A fire deficit persists across diverse North American forests despite recent increases in area burned. *Nature Communications* 16 <https://www.nature.com/articles/s41467-025-56333-8>
- Pielke, R. A., *et al.* (2011). Land use/land cover changes and climate: Modeling analysis and observational evidence. *Wiley Interdisciplinary Reviews: Climate Change*, 2, 828–850. <https://doi.org/10.1002/wcc.144>
- Porter, K., *et al.* (2011). *Overview of the ARkStorm scenario*. U.S. Geological Survey.
- Quinlan, F. T., T. R. Karl, and C.N. Williams Jr. (1987). *United States Historical Climatology Network (USHCN) serial temperature and precipitation data* (NDP-019). Oak Ridge National Laboratory.
- Runnalls, K.E. and T.R. Oke, 2006: Technique to detect microclimate inhomogeneities in historical records of screen-level air temperature. *Journal of Climate*, 19, 959–978. <https://doi.org/10.1175/JCLI3663.1>
- Samborska, V., and H. Ritchie (2024). Wildfires. *Our World in Data*. <https://ourworldindata.org/wildfires>
- Seneviratne, S. *et al.* (2021) “Chapter 11: Weather and Climate Extreme Events in a Changing Climate.” In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by V. Masson-Delmotte, P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, *et al.*, 1513–1766. Cambridge: Cambridge University Press, 2021. <https://doi.org/10.1017/9781009157896.013>.
- Sommer, L. (2016, November 7). Let it burn: The Forest Service wants to stop putting out some fires. *KQED*. <https://www.kqed.org/science/1134217/let-it-burn-the-forest-service-wants-to-stop-putting-out-some-fires>
- Stephens, Scott *et al.*, (2021) “Fire, Water, and Biodiversity in the Sierra Nevada: A Possible Triple Win,” *Environmental Research Communications* 3, no. 8 (August 6, 2021): p. 081004, <https://doi.org/10.1088/2515-7620/ac17e2>.
- Tyukavina Alexandra, Peter Potapov, Matthew Hansen *et al.* (2022) Global Trends of Forest Loss Due to Fire From 2001 to 2019, *Frontiers in Remote Sensing*. Volume 3 <https://www.frontiersin.org/journals/remote-sensing/articles/10.3389/frsen.2022.825190/full>
- U.S. Forest Service. (2022). *Managing fire*. <https://www.fs.usda.gov/science-technology/fire>
- USGCRP. (2023). *USGCRP indicators platform: Heat waves*. <https://www.globalchange.gov/indicators/heat-waves>
- Vecchi, Gabriel A. and Thomas R. Knutson (2011) “Estimating Annual Numbers of Atlantic Hurricanes Missing from the HURDAT Database (1878–1965) Using Ship Track Density,” *Journal of Climate* 24, no. 6 (2011): 1736–1746, <https://doi.org/10.1175/2010JCLI3810.1>.

- Villarini, Gabriele, Gabriel A. Vecchi, and James A. Smith, “U.S. Landfalling and North Atlantic Hurricanes: Statistical Modeling of Their Frequencies and Ratios,” *Monthly Weather Review* 140, no. 1 (January 2012): 44–65, <https://doi.org/10.1175/MWR-D-11-00063.1>.
- Web Archive. (n.d.). *Archive of NIFC fire data*.
https://web.archive.org/web/20200212033452/https://www.nifc.gov/fireInfo/fireInfo_stats_totalFires.html
- Yang, L., Y. Yang, Y. Shen, *et al.* (2024). Urban development pattern’s influence on extreme rainfall occurrences. *Nature Communications*, 15, 3997. <https://doi.org/10.1038/s41467-024-48533-5>
- Zhang, W., G. Villarini, G.A. Vecchi, and J.A. Smith, J. A. (2018). Urbanization exacerbated the rainfall and flooding caused by Hurricane Harvey in Houston. *Nature*, 563, 384–388.
<https://doi.org/10.1038/s41586-018-0676-z>

7 CHANGES IN SEA LEVEL

Chapter Summary

Since 1900, global average sea level has risen by about 8 inches. Sea level change along U.S. coasts is highly variable, associated with local variations in processes that contribute to sinking and also with ocean circulation patterns. The largest sea level increases along U.S. coasts are Galveston, New Orleans, and the Chesapeake Bay regions - each of these locations is associated with substantial local land sinking (subsidence) unrelated to climate change.

Extreme projections of global sea level rise are associated with an implausible extreme emissions scenario and inclusion of poorly understood processes associated with hypothetical ice sheet instabilities. In evaluating the IPCC AR6 projections to 2050 (with reference to the baseline period 1995–2014), almost half of the interval has elapsed by 2025, with sea level rising at a lower rate than predicted. U.S. tide gauge measurements reveal no obvious acceleration beyond the historical average rate of sea level rise.

7.1 Global sea level rise

Global sea level rise is arguably the most important climate impact driver that is unambiguously associated with increasing temperatures. At the global level, warming raises sea level through thermal expansion of sea water and through melting of glaciers and ice sheets. Variations in land water storage are another important factor. At the regional scale, sea level change is influenced by large-scale ocean circulation patterns, and geologic processes and deformation from the redistribution of ice and water. Locally, vertical land motion from geologic processes, ground water withdrawal, and fossil fuel extraction are also important.

The IPCC AR6 estimates that global mean sea level increased by 7.9 (5.9–9.8) inches between 1901 and 2018, with the rate of sea level rise accelerating in recent decades. At the ocean basin scale, sea levels have risen fastest in the Western Pacific and slowest in the Eastern Pacific over the period 1993–2018 (Fox-Kemper *et al.*, 2021). The rate of global sea level rise is estimated to be 0.12 inches/year, about the height of two stacked pennies (NASA, 2020).

The observing systems for global sea level rise have advanced significantly in the satellite era, particularly with the advent of satellite altimeters in 1993. Local tide gauges have provided useful data for the past century, and even longer for a few locations. Following the end of the Little Ice Age in the mid-nineteenth century, tide gauges show that the global mean sea level began rising during the period 1820–1860, well before most anthropogenic greenhouse gas emissions.

7.2 U.S. sea level rise

Observed and predicted rates of mean global sea level rise might have little scientific relevance for specific locations, owing to local processes (NOAA, 2025). Figure 7.1 shows that in Canada and Alaska (and also northern Washington), sea level is decreasing, owing to uplift from glacial rebound. Most of the Pacific coast tide gauges show low rates of sea level rise, while largest U.S. rates are on the Gulf coast (Louisiana and Texas) and in the mid-Atlantic states (Chesapeake Bay region).




Figure 7.1. Map of rates of relative sea level rise along the U.S. coast (NOAA, <https://tidesandcurrents.noaa.gov/sltrends/>). For reference, 3 mm = 0.12 in.

Measurements of relative sea level rise from tide gauges conflate the climatologically relevant increase in the volume of seawater with local vertical land motion. The latter, which varies from place to place, is best measured using a Global Positioning System (GPS) station located near the tide gauge. It is driven by a range of processes that can be comparable to the climate signal and can locally exacerbate (subsidence/sinking) or mitigate (uplift) the risk of sea level rise (Wöppelmann and Marcos, 2016). Human activities triggering subsidence are often significant. They include soil drainage (*e.g.* for urban development) and subsurface extraction of groundwater or hydrocarbons.

Table 7.1 shows absolute sea-level rise (ASLR) for selected locations, determined from the sum of uncorrected sea-level rise as estimated from tide gauge time series (RSLR) (NOAA, 2025) and the vertical land motion (VLM) measurements (NAS, 2012; Letetrel *et al.*, 2015; Karegar *et al.*, 2016). The absolute sea level rise for each of these locations is significantly smaller than the measured relative sea level rise owing to local subsidence. More than half of the measured relative sea level rise is attributed to land sinking for these locations: San Francisco, Galveston, Grand Isle. For reference, the global average rate of absolute sea level rise is estimated to be 0.12 inches/year.

Location	RSLR	VLM	ASLR
San Francisco, CA	+0.08	-0.06	+0.02
Galveston, TX	+0.26	-0.19	+0.07
Grand Isle, LA	+0.36	-0.28	+0.08
St Petersburg, FL	+0.12	-0.02	+0.10
New York City, NY	+0.11	-0.05	+0.06

Table 7.1 Absolute sea level rise (inches/year) consisting of Relative Sea Level Rise (RSLR) plus Vertical Land Motion (VLM)

San Francisco Bay

Over the past 100 years, relative sea level in the San Francisco Bay area has risen by 7.8 inches, at an average rate of 0.08 inches/year (Figure 7.2). As shown in Table 7.1, San Francisco’s vertical land motion is -0.06 inches/year (sinking), producing a recent absolute rate of +0.02 inches/year. Portions of Treasure Island, San Francisco International Airport, and Foster City are sinking as fast as 0.4 inches/year (Shirzaei and Bürgmann, 2018). Problems in the San Francisco Bay area, including threats to the airport, are caused primarily by soil compaction in landfill zones that were formerly wetlands, not by the slow creep of global sea level rise.




Figure 7.2. Tide gauge measurements at San Francisco, California, obtained from NOAA - https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=9414290 (downloaded 4/22/25).

Galveston - Houston

Long-term tide measurements at Galveston Pier 21 show sea level rise of 2.18 feet in the past century, or a rate of 0.26 inches/year (Figure 7.3). Vertical land motion (subsidence) at Galveston is estimated at -0.19 inches/year, yielding an absolute rate of rise of +0.07 inches/year (Table 7.1). The U.S. Geologic Survey found that most of the land-surface subsidence in the Houston-Galveston region is a direct result of groundwater withdrawals (Kasmarek and Ramage 2017), which caused compaction of the aquifer sediments, mostly in the fine-grained silt and clay layers. By 1979, as much as 10 feet of subsidence had occurred in Houston.




Figure 7.3. Tide gauge measurements Galveston Pier, TX, obtained from NOAA - https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8771450 (downloaded 4/22/2025).

New Orleans and the Mississippi delta

Long-term tide gauge measurements at Grand Isle, Louisiana, show that sea level has risen by slightly more than 3 feet over the last 100 years at an average rate of 0.36 inches/year (Figure 7.4). Vertical land motion (subsidence) is estimated -0.28 inches/year. Table 7.1 gives the absolute sea level rise as +0.08 inches/year.




Figure 7.4. Tide gauge measurements at Grand Isle, LA, obtained from NOAA - https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8761724 (downloaded 4/22/25).

The issues of sea level rise and land loss in the Mississippi Delta region are complex, with geological subsidence and the decline in sediment transported by the Mississippi River being the dominant drivers. The construction of dams in the basin since the 1950s has decreased the Mississippi’s suspended sediment load by ~50 percent (Maloney 2018). A new subsidence map of coastal Louisiana finds the coastal region to be sinking at about one third of an inch per year (Neinhuis *et al.* 2017), associated with groundwater and resource withdrawal. As the city’s elevation averages one to two feet *below* sea level, sea level rise from anthropogenic warming is hardly the dominant driver of New Orleans’ problems.

New York City

New York City is particularly vulnerable to the effects of sea level rise because it is built primarily on islands and has 520 miles of coastline. Measurements by a tide gauge at the southern tip of Manhattan (The Battery) show that relative sea level has risen over 11 inches over the past century, at an average rate of 0.11 inches/year (Figure 7.5). But vertical land motion in the New York City area is -0.05 inches/year (roughly 5 inches per century), so that the absolute rate of sea level rise at The Battery is 0.06 inches/year, or about 55 percent of the measured relative sea level rise.




Figure 7.5. Tide gauge measurements at The Battery, New York, obtained from NOAA - https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8518750 (downloaded 4/22/25).

7.3 Projected sea level rise

The concern over sea level rise is not about the roughly eight inches of global rise since 1900. Rather, it is about projections of *accelerated* rise based upon simulations of a warming climate through the 21st century.

AR6 finds *high agreement* across published global mean sea level projections for 2050 with little sensitivity to emissions scenarios. Considering only projections incorporating ice-sheet processes in whose quantification there is at least *medium confidence*, the global sea level projections for 2050, across all emissions scenarios, fall between 3.94 and 15.75 inches (5th–95th percentile *very likely* range) relative to the 1995–2014 baseline period (Fox-Kemper *et al.*, 2021).

Conversely, AR6 states there is *low agreement* across published global mean sea level projections for 2100, particularly for higher emissions scenarios. Considering only projections representing ice sheet processes in whose quantification there is at least *medium confidence*, the AR6 global mean sea level projections for 2100 lie between 7.9 and 39.4 inches (5th–95th percentile *very likely* range) under the medium emissions scenario SSP2–4.5 (Fox-Kemper *et al.*, 2021). There is deep uncertainty surrounding projections of sea level rise to 2100 owing to uncertainties in ice sheet instabilities, particularly for the higher emissions scenarios.

In February 2022, NOAA issued its projections of sea level rise for various sites along the U.S. coast (Sweet *et al.*, 2022). They claim that by 2050, the sea will have risen one foot at The Battery in Manhattan (relative to 2020). A one-foot rise in thirty years would be more than twice the current rate and about three times the average rate over the past century. In that historical context, NOAA’s projection is remarkable—as shown in Figure 7.6, it would require a dramatic acceleration beyond anything observed since the early 20th century. But even more noteworthy is that Sweet *et al.* (2022) say this rise is “locked in”—it will happen no matter what future emissions are. We should know in a decade or so whether that prediction has legs.




Figure 7.6 Rate of sea level rise at the Battery in Manhattan. Shown is the historical thirty-year trailing trend, together with the allegedly “locked in” NOAA predicted trend for 2050. Historical data: NOAA Tides and Current.

References

- Fox-Kemper, B., Hewitt, H. T., Xiao, C., Adalgeirsdóttir, G., Drijfhout, S. S., Edwards, T. L., ... & Yu, Y. (2021). Ocean, cryosphere and sea level change. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 1211–1362). Geneva, Switzerland: Intergovernmental Panel on Climate Change.
- Karegar, M. A., Dixon, T. H., & Engelhart, S. E. (2016). Subsidence along the Atlantic Coast of North America: Insights from GPS and late Holocene relative sea level data. *Geophysical Research Letters*, 43(7), 3126–3133. <https://doi.org/10.1002/2016GL068015>

- Kasmarek, M. C., & Ramage, J. K. (2017). *Water-level altitudes 2017 and water-level changes in the Chicot, Evangeline, and Jasper aquifers and compaction 1973–2016 in the Chicot and Evangeline aquifers, Houston-Galveston Region, Texas* (Scientific Investigations Report 2017–5080). U.S. Geological Survey. <https://doi.org/10.3133/sir20175080>
- Letetrel, C., Becker, M., Llovel, W., & Cazenave, A. (2015). Estimation of vertical land movement rates along the coasts of the Gulf of Mexico over the past decades. *Continental Shelf Research*, 111(Part A), 42–51. <https://doi.org/10.1016/j.csr.2015.10.018>
- Maloney, J. M., Bentley, S. J., Xu, K., Obelcz, J., Georgiou, I. Y., & Miner, M. D. (2018). Mississippi River subaqueous delta is entering a stage of retrogradation. *Marine Geology*, 400, 12–23. <https://doi.org/10.1016/j.margeo.2018.03.001>
- National Academy of Science (NAS, 2012). *Sea-level rise for the coasts of California, Oregon and Washington: Past, present, and future*. Appendix A: Vertical land motion and sea-level data along the West Coast of the United States. National Academies Press. <https://nap.nationalacademies.org/catalog/13389/sea-level-rise-for-the-coasts-of-california-oregon-and-washington>
- NASA. (2020). *NASA sea level rise portal: 2020 edition*. <https://www.nasa.gov/specials/sea-level-rise-2020/>
- National Oceanic and Atmospheric Administration (NOAA) Center for Operational Oceanographic Products and Services. (n.d.). *Sea level trends*. <https://tidesandcurrents.noaa.gov/sltrends/>
- NOAA Tides & Currents. (n.d.). *Relative sea level trends*. <https://tidesandcurrents.noaa.gov/sltrends/>
- Nienhuis, J. H., Törnqvist, T. E., Esposito, C. R., Liang, M., & Ma, H. (2017). A new subsidence map for coastal Louisiana. *GSA Today*, 27(6), 28–29. <https://doi.org/10.1130/GSATG337GW.1>
- Shirzaei, M., & Bürgmann, R. (2018). Global climate change and local land subsidence exacerbate inundation risk to the San Francisco Bay Area. *Science Advances*, 4(3), eaap9234. <https://doi.org/10.1126/sciadv.aap9234>
- Sweet, W. V., Hamlington, B. D., Kopp, R. E., et al. (2022). *Global and regional sea level rise scenarios for the United States* (USGS Report 70229139). United States Geological Survey. <https://www.usgs.gov/publications/global-and-regional-sea-level-rise-scenarios-united-states>
- Wöppelmann, G., & Marcos, M. (2016). Vertical land motion as a key to understanding sea level change and variability. *Reviews of Geophysics*, 54, 64–92. <https://doi.org/10.1002/2015RG000502>

8 UNCERTAINTIES IN CLIMATE CHANGE ATTRIBUTION

Chapter summary

“Attribution” refers to identifying the cause of some aspect of climate change, specifically with reference to anthropogenic activity. There is an ongoing scientific debate around attribution methods, particularly regarding extreme weather events. Attribution is made difficult by high natural variability, the relatively small expected anthropogenic signal, lack of high-quality data, and reliance on deficient climate models. The IPCC has long cautioned that methods to establish causality in climate science are inherently uncertain and ultimately depend on expert judgement.

Substantive criticism of the main IPCC assessments of the role of CO₂ in recent warming focus on inadequate assessment of natural climate variability, uncertainties in measurement of solar variability and in aerosol forcing, and problems in the statistical methods used for attribution.

The IPCC does not make attribution claims for most climate impact drivers related to extreme events. Statements related to statistics of global extremes (*e.g.* event probability or return times, magnitude and frequency) are not generally considered accurate owing to data limitations and are made with low confidence. Attribution of individual extreme weather events is challenging due to their rarity. Conflicting claims about the causes of the 2021 Western North America Heatwave illustrate the perils of hasty attribution claims about individual extreme events.

8.1 Introduction

The Intergovernmental Panel on Climate Change (IPCC) distinguishes between detection of climate change and attribution of its cause. As defined by the AR6 WGI Glossary (IPCC, 2025):

Detection: Detection of change is defined as the process of demonstrating that climate or a system affected by climate has changed in some defined statistical sense, without providing a reason for that change. An identified change is detected in observations if its likelihood of occurrence by chance due to internal variability alone is determined to be small, for example, <10%.

Attribution: Attribution is defined as the process of evaluating the relative contributions of multiple causal factors to a change or event with a formal assessment of confidence.

Both detection and attribution rely on statistical analysis. Detection focuses on whether changes are significant enough to stand out from random variation regardless of cause. Attribution involves comparison of observed events to model-generated counterfactuals. Since experimentation on the climate is not possible, attribution requires statistical inference to assess how much human activities (such as GHG emissions) versus natural factors (like volcanic eruptions) contribute to observed changes. Attribution methods assume all external and internal drivers of the system are known and represented.

There are ongoing scientific debates around attribution methods, especially those for attributing extreme weather events to climate change. The IPCC has long cautioned that methods to establish causality in climate science are inherently uncertain and ultimately depend on expert judgement. AR4 Working Group I (Hegerl *et al.*, 2007) explained it as follows:

‘Attribution’ of causes of climate change is the process of establishing the most likely causes for the detected change with some defined level of confidence... unequivocal attribution would require controlled experimentation with the climate system. Since that is not possible, in practice attribution of anthropogenic climate change is understood to mean demonstration that a detected change is ‘consistent with the estimated responses to the given combination of anthropogenic and natural forcing’ and ‘not consistent with alternative, physically plausible explanations of recent climate change that exclude important elements of the given combination of forcings’ (IPCC, 2001)... The approaches used in detection and attribution research described above cannot fully account for all uncertainties, and thus ultimately expert judgement is required to give a calibrated assessment of whether a specific cause is responsible for a given climate change.

AR5 Working Group II (Cramer *et al.*, 2014) makes the following statement:

Two broad challenges to the detection and attribution of climate change impacts relate to observations and process understanding. On the observational side, high-quality, long-term data relating to natural and human systems and the multiple factors affecting them are rare. In addition, the detection and attribution of climate change impacts requires an understanding of the processes by which climate change, in conjunction with other factors, may affect the system in question.

Because of the complexity of the causal chains in the climate system, investigation of these relationships is exceptionally challenging.

8.2 Attribution methods

The IPCC employs several attribution methods to assess the causes of observed climate changes, distinguishing between natural and human-induced factors. Below is a concise description of the key IPCC attribution methods.

Optimal Fingerprinting uses linear regression to explain variations in observed climate data as a weighted sum of climate model simulations run with and without anthropogenic forcings. The data used in the regression model is weighted to try to minimize the influence of random noise and the estimation method is chosen to account for climate model error.

Time Series Analysis exploits statistical differences between anthropogenic forcing and natural variability to see which dominates observed temperatures and also uses variations in the timing of changes to determine if causal dependence across data series can be inferred.

Process-Based Attribution focuses on understanding the physical mechanisms behind observed changes. This approach combines observations, climate models, and theoretical understanding to attribute changes in specific processes to forcings. This approach is often used for regional climate phenomena, such as monsoon changes or polar amplification

Extreme Event Attribution assesses the role of human influence in the likelihood of occurrence or intensity of extreme weather events (*e.g.* heat waves or droughts). Methods include:

- Probabilistic Event Attribution uses large ensembles of climate model simulations to compare observed events to model-generated counterfactuals
- Storyline Approach examines the physical processes driving an event and evaluates how anthropogenic forcings might have modified those processes

8.3 Attribution of global warming

Attribution statements for global warming in the three most recent IPCC Assessment reports are:

AR4: **Most of the observed increase** in global average temperatures since the mid-20th century is *very likely* due to the observed increase in anthropogenic greenhouse gas concentrations. (IPCC 2007)

AR5: It is *extremely likely* that **more than half of the observed increase** in global average surface temperature from 1951 to 2010 was caused by the anthropogenic increase in greenhouse gas concentrations and other anthropogenic forcings together. The best estimate of the human-induced contribution to warming is similar to the observed warming over this period. (IPCC 2013)

AR6: The *likely* range of total human-caused global surface temperature increase from 1850–1900 to 2010–2019 is 0.8°C to 1.3°C, with a best estimate of 1.07°C. It is *likely* that well-mixed GHGs contributed a warming of 1.0°C to 2.0°C, other human drivers (principally aerosols) contributed a cooling of 0.0°C to 0.8°C, natural drivers changed global surface temperature by –0.1°C to +0.1°C, and internal variability changed it by –0.2°C to +0.2°C. It is *very likely* that **well-mixed GHGs were the main driver** of tropospheric warming since 1979. (IPCC 2021)

The AR4 and AR5 attribution statements reference the warming since the mid-20th century, the period when greenhouse gas emissions began increasing rapidly. The words “most” and “more than half” are deliberately imprecise, potentially ranging from 51 to 99 percent of the warming – presumably this imprecision is to account for structural uncertainties such as natural internal variability. The confidence level increases from *very likely* to *extremely likely* from AR4 to AR5. The structure of the attribution statement in the AR6 is fundamentally different, referencing the warming to the period 1850–1900. The AR6 attribution statement is more precise numerically, but with a lower level of confidence at “likely” – AR6 attributes essentially all the warming to increases in greenhouse gases. The most confident statement from AR6 relates to the tropospheric warming since 1979, using the words “main driver” and “*very likely*.”

There are three areas of substantive criticism of the IPCC’s assessment of the causes of the recent warming: inadequate assessment of natural climate variability, inappropriate statistical methods, and substantial discrepancies between models and observations. The last is discussed in Chapter 5, while this chapter discusses the first two factors. All of these criticisms are relevant to the IPCC’s attribution of the recent warming, which also underpins extreme event attribution

8.3.1 Natural climate variability

AR6 states that natural external drivers since 1850–1900 have changed global surface temperature by –0.1°C to +0.1°C, and internal variability has changed it by –0.2°C to +0.2°C – on average having essentially no net impact on the warming since 1850–1900. As discussed below, this minimal contribution of natural variability has been disputed by several publications that question the magnitudes of solar variability and internal variability from large-scale ocean circulations.

Solar variability

AR5 concluded that the best estimate of radiative forcing due to Total Solar Irradiance (TSI) changes over the period 1750–2011 was very small (0.05 W/m², Myrhe *et al.*, 2014). AR6 acknowledges substantially higher values and a much larger range of estimates of changes in TSI over the last several centuries, stating that the TSI between the Maunder Minimum (1645–1715) and the second half of the twentieth century increased by 0.7–2.7 W/m², a range that includes both low and high variability TSI data

sets (Gulev, 2021). However, the recommended forcing dataset for the CMIP6 climate model simulations used in AR6 for attribution studies averages two data sets with low solar variability (Matthes, 2017).

While AR6 shows a substantially greater solar impact than does AR5, the overall impact of solar forcing on the climate was still assessed to be small compared to anthropogenic forcing. However, the impact of solar variations on the climate is uncertain and subject to substantial debate (Lockwood, 2012; Connolly *et al.*, 2021) - something that is not evident in the IPCC assessment reports.

The variations of TSI over time remains a challenging problem. Since 1978, there have been direct measurements of TSI from satellites. However, the data exhibits non-negligible inconsistencies, and interpreting any multi-decadal trends in TSI requires comparisons of observations from overlapping satellites. There are several rival composite TSI datasets that disagree as to whether TSI increased or decreased during the period 1986–96 (the ACRIM gap; see Chapter 4). Further, the satellite record of TSI is used to calibrate proxy models that infer past solar variations from sunspots and cosmogenic isotope measurements (Velasco Herrera *et al.*, 2015).

There is substantial evidence for high solar activity in the second half of the 20th century (starting in 1959) and extending into the 1990's, before a decline in the early 21st century; this period is often termed the “Modern Maximum.” (Chatzistergos *et al.*, 2023; Solanki *et al.*, 2004; Usoskin *et al.*, 2007). However, some scientists have concluded that it is not possible to be confident of any multi-decadal trend in TSI (Schmutz, 2021).

This uncertainty causes some reconstructions of TSI from 1750 to have low variability (implying a very low impact of solar variations on global mean surface temperature) whereas datasets with high TSI variability can explain more than 70 percent of the temperature variability since preindustrial times (Scafetta, 2013; Stefani, 2021). The choice of TSI satellite record used in an analysis can therefore substantially influence how much climate change is attributed to human versus natural forcings.

There is growing evidence that other aspects of solar variability, which are referred to as solar indirect effects, either amplify TSI forcing or are independent of TSI forcing. Scafetta *et al.* (2023) suggests that ~80 percent of solar influence on climate might stem from non-TSI mechanisms. There are numerous candidate processes, including solar ultraviolet changes; energetic particle precipitation; atmospheric-electric-field effect on cloud cover; cloud changes produced by solar-modulated galactic cosmic rays; large relative changes in the magnetic field; and the strength of the solar wind. Such solar indirect effects are not included in climate models, although indirect methods of estimating their impacts suggest they are significant. However, claims of non-TSI mechanisms influencing climate are uncertain and debated.

Natural variability of large-scale ocean circulations

Variations in global mean surface temperature are linked to recurrent large-scale variations in ocean circulation patterns, including the Atlantic Multidecadal Oscillation (AMO), the Pacific Decadal Oscillation (PDO) and the El Niño-Southern Oscillation (ENSO). These circulations influence ocean heat uptake and heat distribution and also influence atmospheric circulation patterns and cloud distributions. There is some debate as to whether these variations are strictly internal to the climate system, or whether this variability can have a solar/astronomical origin or can be influenced by large volcanic eruptions.

While climate models simulate the large-scale ocean circulations and internal climate variability, most models have too little amplitude compared to observations at multi-decadal frequencies and phasing out-of-sync with the observed climate (Kravtsov *et al.* 2024). Averaging multiple simulations effectively averages out the internal variations, leaving only the forced climate variability (*e.g.* CO₂ forcing). With most of the modern warming beginning in the late 1970's, the recent 50-year warming is on the same time scale as the multi-decadal oscillations of the AMO and PDO.

Here are summary statements from the AR5 and AR6 reports:

AR5: Decadal variability in the Pacific, associated with the PDO or IPO [Interdecadal Pacific Oscillation], contributes significantly to regional and global temperature trends, but the relative contributions of internal variability and external forcing are difficult to disentangle in CMIP5 simulations.

AR6: Since AR5, there has been increased understanding of the role of internal variability, such as ENSO, PDO, and AMO, in modulating regional climate trends. However, limitations in simulating the exact timing and amplitude of these modes in CMIP6 models contribute to uncertainties in attributing observed changes to anthropogenic forcing (*high confidence*).

The amplitude of the peak-to-trough impact of the multi-decadal oscillations on global temperatures has been assessed by the IPCC AR6: “The likely range of change due to internal variability is -0.2°C to $+0.2^{\circ}\text{C}$ (IPCC, 2021).” This implies a trough-to-peak change of 0.4°C . Over many centuries, any global temperature changes from the troughs and peaks will cancel out, with little to no net impact.

However, with a nominal timescale of 60-80 years for oscillations such as the AMO and PDO, the timing of the peaks and troughs can be confused with the secular trend. This becomes very relevant for attributing the warming for the past 50 years, when most of the recent warming has occurred. Even if climate models have the correct amplitude of the multidecadal oscillations, the timing of the peaks and troughs is not adequately simulated, with each model and individual ensemble simulating a different phasing. Once the simulations of the ensemble members and different models are averaged, the natural internal variability contribution is averaged out owing to differences in phasing, effectively cancelling the role of natural internal variability in the attribution process.

The time series of global surface temperature anomalies since 1850 (Figure 8.1) shows irregular variations of significant amplitude against the background of an overall warming trend and, after 1976, a pronounced difference in trends between the Northern Hemisphere and the Southern Hemisphere. The period between 1905-1945 shows a strong trend of warming. The following 30 years from 1945-1976 showed slightly declining temperatures. The most recent warming period started in 1977.




Figure 8.1 Global average surface temperature anomalies 1850—2025. Top: Northern hemisphere. Middle: Southern hemisphere. Bottom: Global average. Source: UK Hadley Centre <https://crudata.uea.ac.uk/cru/data/temperature/HadCRUT5.0Analysis.pdf>

The causes of the early 20th century warming are discussed by Hegerl *et al.* (2017; 2019). The atmospheric CO₂ concentration increased from 298 ppm in 1905 to 310 ppm in 1941, implying that CO₂ had little impact. Volcanic activity during this period was very low, and solar forcing is uncertain. Yet Hegerl *et al.* (2017) somehow inferred that 40-54 percent of this warming could be attributed to external forcing, with the rest associated with internal variability. Bronniman *et al.* (2024) focused on the causes of the cooling in the first decade of the 20th century in the Southern Hemisphere. They found that the cooling was related to a La-Niña-like pattern in the Pacific, a cold tropical and subtropical South Atlantic, a cold extratropical South Pacific, and cool southern midlatitude land areas. The Southern Annular Mode was positive, with a strengthened Amundsen–Bellingshausen seas low, although the spread of the data products is considerable.

The warming in the 1930's and subsequent cooling during mid-century was particularly pronounced in the Arctic. Bokuchava and Semenov (2021) find that these variations were most likely caused by a combined effect of long-term natural climate variations in the North Atlantic and North Pacific with a contribution of the natural radiative forcing related to the reduced volcanic activity and variations of solar activity, as well as growing greenhouse gases concentrations. Tokinaga *et al.* (2017) showed that the combined effect of internally generated Pacific and Atlantic interdecadal variability intensified Arctic

warming in the early 20th century. The synchronized Pacific-Atlantic warming drastically alters planetary-scale circulations over the Northern Hemisphere; these same circulation patterns have a global influence.

The cooling period between 1945 and 1976 has been referred to as the “grand hiatus.” Numerous causes have been hypothesized: natural internal variability from fluctuations associated with the Pacific Decadal Oscillation and Atlantic Multidecadal Oscillation; cooling from increased emissions of aerosols from industrial activities; increased heat uptake in the Atlantic, Southern and Equatorial Pacific Oceans. Thompson *et al.* (2010) find that the hemispheric differences in temperature trends in the middle of the twentieth century stem largely from a rapid drop in Northern Hemisphere sea surface temperatures of about 0.3 °C between about 1968 and 1972.

The Great Pacific Climate Shift of 1976-1977 was a notable climatic event characterized by an abrupt change in the North Pacific Ocean's atmosphere-ocean system that interacted with global climate patterns. This shift is closely associated with the Pacific Decadal Oscillation (PDO) that oscillates between warm and cool phases over decades. The 1976 shift marked a transition from a predominantly negative (cool) PDO phase (1947–1976) to a positive (warm) phase (1977 through 2000), with significant implications for global and regional climate patterns, including the frequencies of El Nino and La Nina events. The Great Pacific Climate Shift coincided with the beginning of a period of accelerated global warming. When the Great Pacific Climate Shift is accounted for in climate attribution analyses since 1950, 40 percent or more of the warming in the second half of the 20th century is attributed to natural internal variability (McLean *et al.*, 2009; Tung and Zhou, 2013; Chylek *et al.*, 2016; Scafetta, 2021).

8.3.2 Optimal fingerprinting

Optimal fingerprinting is a statistical technique introduced by Allen and Tett (1999) that compares observed climate data to climate model simulations to identify patterns (or “fingerprints”) associated with human or natural forcing. It involves taking a vector of observed climate changes, such as warming rates in locations around the world, and decomposing them into a weighted sum of “signals”, which are analogues to the observations generated by climate models with different types of forcings. The weights are chosen using a regression method called Total Least Squares (TLS). The analysis commonly uses just two signals, one generated by models using only anthropogenic forcings and one using only natural forcings. If the estimated weighting coefficient attached to a signal is significantly different from zero, then that signal is said to be “detected”. If it is close to 1.0, then the model generating the signal is said to be consistent with observations. If it is less than 1.0, then the model signal for that forcing is too strong and needs to be scaled down, and vice-versa. Both the observed data and the signals are weighted using a model-generated estimate of patterns of randomness in the climate system so as, in principle, to put maximum weight on regions where natural variability is minimized.

Optimal fingerprinting is the primary tool for attribution in the research literature. While it has been widely used and prominently featured by the IPCC since 2001, there is very little literature examining the statistical properties of the results it generates. One of its inherent weaknesses is that results depend on assumptions about the accuracy of climate models, especially regarding their representation of natural variability (IPCC AR5 Ch 10.2.3).

A series of papers by McKittrick (McKittrick 2021, 2022, 2023, 2025) has challenged the optimal fingerprinting method, arguing that it is inherently unreliable and that practices adopted from econometrics can provide more valid results. Statistical theory requires that for regression models to yield unbiased coefficients, a set of assumptions called the Gauss-Markov conditions must hold. McKittrick (2021) argued that the Allen and Tett (1999) methodology violates the Gauss-Markov conditions, leading to potentially biased and inconsistent fingerprinting coefficients, undermining the reliability of the method. Chen *et al.* (2023) confirmed this analysis, although they argued that the method could yield valid results under very stringent assumptions. McKittrick (2022) and (2023) raised a further concern that climate scientists—virtually alone among

scientific disciplines—have used TLS to estimate anthropogenic greenhouse gas signal coefficients, despite its tendency to be unstable unless some strong assumptions hold that in practice are unlikely to be true. Under conditions that easily arise in optimal fingerprinting, TLS estimates have a large positive bias. Thus, any study that used TLS for optimal fingerprinting without verifying its applicability in the specific data context has likely overstated the attribution.

McKittrick (2025) presented an empirical example comparing the results of conventional optimal fingerprinting against methods drawn from mainstream econometrics that are known to be valid for the specific application of signal detection. While the IPCC optimal fingerprinting method yields an anthropogenic signal coefficient close to 1.0 on a global temperature data set spanning 1900 to 2010, the consistent method yields a coefficient around 0.4, which rises to about 0.65 on data spanning 1980 to 2010, implying the model response to greenhouse gases needs to be scaled down by about half to optimally match observations. The natural forcing signal coefficient, by contrast, is between 2.0 and 4.0, implying the climate model signals of natural forcing need to be scaled up two-to four-fold to match observed climate change. The fingerprinting coefficients estimated in McKittrick (2025), when used to scale the average sensitivity of the climate models used to generate the forcing signals in his data set, imply a Transient Climate Sensitivity of 1.4°C, which is consistent with the estimate by Lewis (2023) using a different estimation method and multiple independent data sets.

These findings indicate that the basis on which the optimal fingerprinting method has long been viewed as reliable is not valid. Re-examining previous results individually would be required to determine which findings are statistically robust.

8.3.3 Time series methods

The IPCC (AR5 WGI 10.2.2) drew attention to alternative approaches of assessing causality that emerged from the time series econometrics literature. These have the advantage of not depending on assumptions about the accuracy of climate models, but the disadvantage that they depend on difficult-to-test assumptions about the data generating process underlying climate and forcing data. The methods, now referred to as climate econometrics, use the tools of unit root testing, Granger causality and cointegration analysis, all of which are familiar in economics and finance and which are slowly being adopted in climate science. Time series analysis methods hold out the possibility of determining whether anthropogenic or natural forcings are the primary drivers of climate change without requiring the use of climate models, although key questions remain unsettled (e.g. Dergiades *et al.*, 2016, Balcombe *et al.*, 2019, Dagsvik *et al.*, 2020, Razzak, 2022, Dagsvik and Moen, 2023).

“Granger causality” modeling is a tool for determining directions of influence in data series that move together. It is a statistical, not a physical, concept. If knowing the current value of one variable significantly improves the forecast accuracy of another variable, we can infer a causal connection exists; this is called Granger causality. The modeling tools, called vector autoregression, measure direct impulse-response patterns and feedbacks. An application to the well-known Vostok ice core data revealed an error in Al Gore’s documentary *An Inconvenient Truth*. Gore showed the data and drew attention to the coherence of temperature and CO₂ changes over a 440,000 year span, which he asserted was due to CO₂ driving temperature changes. But temperature changes can also affect atmospheric CO₂ levels. Davidson *et al.* (2015) examined the series and found that temperature Granger causes CO₂ but not the reverse. In other words on the time scales represented in the Vostok data, the coherence in the series is primarily due to the influence of temperature on CO₂ levels, not the feedback of CO₂ levels on temperature.

In summary, the primary statistical methods for attributing causation in climate data are optimal fingerprinting and time series analysis. Applications of the Allen and Tett (1999) optimal fingerprinting method dominate the attribution literature and have underpinned past IPCC conclusions, but results depend

on the accuracy of climate models and the method has recently been criticized as inherently biased. Time series methods do not depend on climate models but require assumptions of their own and have generated results that have thus far not converged on a consensus.

8.4 Declining planetary albedo and recent record warmth

A sharp recent increase in global average temperatures has raised the question of short-term drivers of climate. One such candidate is the fraction of absorbed solar radiation which has also increased abruptly in recent years. The question is whether the change is an internal feedback to warming caused by greenhouse gases, or whether something else increased the fraction of absorbed radiation which then caused the recent warming.

The planetary albedo is the fraction of incoming solar radiation that is reflected back into space rather than being absorbed by the planet. Highly reflective surfaces like cloud tops and snow and ice are most important in this regard. The Earth's albedo is approximately 30 percent, meaning almost a third of the sunlight that reaches Earth is directly reflected back to space. A lower albedo implies more solar energy is absorbed by the planet to be then re-radiated as heat. Hence, other things being equal, a decline in planetary albedo is associated with a warming of the Earth.

Arguably the most striking change in the Earth's climate system during the 21st century is a significant reduction in planetary albedo since 2015, which has coincided with at least two years of record global warmth. Figure 8.2 shows the planetary albedo variations since 2000, when there are good satellite observations. The 0.5 percent reduction in planetary albedo since 2015 corresponds to an increase of 1.7 W/m² in absorbed solar radiation averaged over the planet (Hansen and Karecha, 2025). For comparison, Forster et al. (2024) estimate the current forcing from the increase in atmospheric CO₂ compared to preindustrial times to be 2.33 W/m².

Looking back prior to 2000 with less adequate data, Goessling *et al.* (2025) assessed that the planetary albedo was relatively low around the 1940's and 50's before rising industrial aerosol precursor emissions increased the albedo until the 1980's. The strongest planetary albedo excursions were high-albedo episodes caused by volcanic eruptions, such as after the Mount Pinatubo eruption in 1991. Although uncertain, the 2023 planetary albedo minimum might have been the lowest since at least 1940.

Changes in surface characteristics cannot explain this decrease in planetary albedo since 2015:

- Arctic sea ice extent has declined by about 5 percent since 1980 (https://nsidc.org/data/seaice_index/images/s_plot_hires.png), although following 2007 there has been a pause in the Arctic sea ice decline (England *et al.*, 2025)
- Regarding Antarctic sea ice, the IPCC AR6 concludes that “There has been no significant trend in Antarctic sea ice area from 1979 to 2020 due to regionally opposing trends and large internal variability.” (Summary for Policymakers, A.1.5)
- Northern hemispheric annual snow cover has been slowly declining since 1967, with barely significant trends. The data show the Northern Hemisphere has snowier winters, accompanied by more rapid melt in spring and summer, see <http://climate.rutgers.edu/snowcover/> and Section 5.6
- Global greening (Chapter 2) is contributing to the decrease in planetary albedo, as forests have a lower albedo than open lands or snow. However, there is some evidence that forests increase cloud cover (high reflectivity), which counteracts the direct albedo decrease associated with increasing forested area.




Figure 8.2. Earth's albedo (reflectivity, in percent), with seasonality removed. From Hansen and Karecha (2025)

Changes in surface reflectivity are contributing to a small, slow decline in planetary albedo. However, these changes cannot explain the sharp decline in planetary albedo beginning in 2015. Further, any changes in surface albedo are effectively damped by about a factor of three on average, primarily due to cloud masking (Loeb *et al.* 2019). Surface albedo changes have thus contributed only weakly to the recent planetary albedo decline, particularly when averaged annually and globally. This leaves the most likely explanation for the sharp albedo decline as changes in atmospheric aerosols and/or clouds.

Aerosols are small particles in the atmosphere that reflect sunlight and can interact with cloud processes to make the clouds more reflective. Aerosols have both natural origins (*e.g.* dust from soils, wildfires) as well as human origins from combustion (including fossil fuels). Regions that are greening and/or have an increase in rainfall might produce less dust. For anthropogenic aerosols, new ship fuel regulations aimed at reducing sulfur emissions were implemented in three phases, in 2010, 2015 and 2020. Sulfate particles reflect solar radiation. Cleaner air means less solar radiation is reflected, which contributes to surface warming. Indirect effects of sulfate aerosols include increasing the reflectivity of low-level clouds in the subtropics. There is controversy surrounding the importance of this change in sulfate emissions to the Earth's radiation balance (Hodnebrog *et al.*, 2024; Yuan *et al.*, 2024), but the restriction of this effect to major shipping routes suggests a small global impact (Schmidt, 2024).

Changes in clouds are the primary candidate to explain the decline in global albedo since 2015. Two recent papers (Loeb *et al.*, 2024; Goessling *et al.* 2024) have addressed recent variations in cloud properties. By considering satellite and reanalysis data, Loeb *et al.* found that decreases in low- and mid-level clouds since 2015 are the primary reason for decreasing planetary albedo in the Northern hemisphere, whereas in the Southern hemisphere the decrease in planetary albedo is primarily due to decreases in mid-level clouds across all latitude zones. Goessling *et al.* (2024) found that cloud anomalies were mainly due to reduced low-level clouds. Regions with coherent low-level cloud reductions over the past decade include the warm pool region around the Maritime Continent and the northern extratropical western Pacific, as well as large parts of the Atlantic and adjacent land regions. The reduction of global cloud cover identified in these analyses since 2015 is 1-2 percent.

The issue then becomes the cause of the change in cloud cover. Two explanations have been posited for the declining cloud cover over the past decade:

- Natural climate variability

- Changes in low cloud cover associated with warming sea surface temperatures, implying an emerging positive feedback to climate change (Hansen and Karecha, 2025)

It is not easy to justify a new positive low cloud feedback that began emerging in 2015 since there is no obvious feedback trigger starting at that time. However, there are numerous natural climate signals during this period that are associated with atmospheric circulation changes that can influence the distribution of clouds:

- The 2014-2016 was one of the strongest El Niño events on record
- A cold anomaly beginning in 2015 in the subpolar gyre of the North Atlantic reflects a shift in the ocean circulation pattern associated with decadal variability in the Atlantic (Frajka-Williams *et al.*, 2017; Arthun *et al.* 2021).
- The Pacific Decadal Oscillation positive index peaked in 2016, then declined and has been in negative territory since late 2019
- Eruption of the submarine Hunga-Tonga volcano in 2022

Interannual cloud anomalies associated with the El Niño Southern Oscillation (ENSO) have a significant global signal and strong regional signals, especially over the tropical Indian and Pacific Oceans.

The Hunga Tonga eruption is remarkable in its coincidence with the exceptionally low value of planetary albedo in 2023. Goessling *et al.* (2024) found that the cloud perturbations in 2023 showed a different pattern from the overall pattern since 2015, with reduced cloud cover being most pronounced in the northern hemisphere and tropics. Regional reductions in cloud cover were most pronounced over the eastern Indian Ocean, over South America and extending over the eastern Pacific, as well as over northern North America, in the Southern Ocean around 60S, in the subtropical and eastern North Atlantic, and in parts of the North Pacific.

The Hunga Tonga eruption was unusual in that it injected large amounts of water vapor into the stratosphere in addition to sulfate particles. Early publications focused on the impacts to stratospheric circulations and the direct radiative impact. In 2025, papers are emerging that examine the indirect effects of Hunga Tonga on surface climate using ensemble simulations from earth system models (Bednarz *et al.*, 2025; Kuchar *et al.*, 2025). These papers found statistically significant impacts on regional climates that were driven by circulation changes from couplings between the stratosphere and troposphere. These papers indicate the complex interactions between volcanic activity and climate dynamics. More research is needed to untangle any impacts of the Hunga Tonga eruption on the planetary albedo.

In summary, the decline in planetary albedo and the concurrent decline in cloudiness have emphasized the importance of clouds and their variations to global climate variability and change. A change of 1- 2 percent in global cloud cover has a greater radiative impact on the climate than the direct radiative effect of doubling CO₂. While it is difficult to untangle causes of the recent trend, the competing explanations for the cause of the declining cloud cover have substantial implications for assessing the Equilibrium Climate Sensitivity and for the attribution of the recent warming. An additional 10 years of data should help clarify whether this is a strong positive cloud feedback associated with warming or a temporary fluctuation driven by natural variability

8.5 Attribution of climate impact drivers

The IPCC (Ranasinghe *et al.* 2021) defines “climate impact drivers” or CIDs as “physical climate system conditions (e.g., means, events, extremes) that affect an element of society or ecosystems.” Hence CIDs are those features of the weather and climate system of primary interest in assessing the impacts of climate change since they potentially affect humans and the natural world. For instance, under the heading “Heat and Cold,” CIDs are identified as mean air temperature, extreme heat, cold spells and frost. The IPCC

also points out that CIDs are not necessarily harm-related: depending on the system in question they can be detrimental, neutral, beneficial or a combination.

The first columns Table 12.12 in Ranasinghe *et al.* (2021, p. 1856), reproduced here as Table 8.1, summarize the AR6 assessment of whether a signal of attributable anthropogenic influence has emerged across all major CIDs. One of the themes of this chapter is that attribution methods used by the IPCC tend to overstate the anthropogenic influence and understate the role of natural variability. Nonetheless a striking feature of that summary table is how few CIDs exhibit an anthropogenic signal sufficient to distinguish them from natural variability.

Out of the 33 weather impact categories listed, an anthropogenic signal is asserted with *high confidence* in only five, and with *medium confidence* in a further four. (Note that one of the CIDs is an increase in CO₂ levels, and since it is a tautology to attribute this to increased CO₂ levels this CID can be ignored.) For the rest the IPCC does not claim to have detected anthropogenic drivers. Of the five *high confidence assertions*, two are for changes in average temperatures (air and ocean) hence are not measures of extreme weather. Further, two of the four *medium confidence* assertions are related to ocean chemistry and thus are likewise not related to extreme weather. The IPCC does not assert a human influence on many non-temperature weather features such as wind, precipitation, flooding, or drought.

Other columns of IPCC Table 12.12 report on whether anthropogenic signals are expected to emerge this century under RCP8.5, the most extreme forcing scenario. We have omitted these columns for several reasons. First, because they refer to whether climate models project detectable signals in the observations, which is a very different question than our concern here: whether a signal has been detected in historical data. Second, as we discuss in Chapter 4, the RCP8.5 scenario is a misleading and implausible high-end storyline, it is not a “base case” or business-as-usual projection. Third, the ensemble of detailed models is acknowledged (Palmer and Stevens 2019) to be “not fit for purpose” in describing regional changes and “unable to represent future conditions at the degree of spatial, temporal, and probabilistic precision with which projections are often provided.” (Nissan et al, 2019). Even with these caveats, we note that the omitted columns in AR6 Table 12.12 show most weather impacts are not expected to exhibit an anthropogenic signal through the end of this century.

As discussed in Chapter 6 natural variability dominates patterns of extreme weather systems and simplistic assertions of trend detection are frequently undermined by regional heterogeneity and trend reversals over time. Table 8.1 makes the related point that it is not currently possible to attribute changes in most extreme weather types to human influences. Taking wind as an example, the IPCC claims that an anthropogenic signal has not emerged in average wind speeds, severe windstorms, tropical cyclones or sand and dust storms, nor is one expected to emerge this century even under an extreme emissions scenario. The same applies to drought and fire weather.

Climatic Impact-driver Type	Climatic Impact-driver Category	Already Emerged in Historical Period
Heat and Cold	Mean air temperature	1
	Extreme heat	2
	Cold spell	4
	Frost	
Wet and Dry	Mean precipitation	
	River flood	
	Heavy precipitation and pluvial flood	
	Landslide	
	Aridity	
	Hydrological drought	
	Agricultural and ecological drought	
	Few weather	
Wind	Mean wind speed	
	Severe wind storm	
	Tropical cyclones	
	Sand and dust storm	
Snow and Ice	Snow, glacier and ice sheet	
	Permafrost	
	Lake, river and sea ice	11
	Heavy snowfall and ice storm	
	Hail	
	Snow avalanche	
Coastal	Relative sea level	
	Coastal flood	
	Coastal erosion	
Open Ocean	Mean ocean temperature	
	Marine heatwaves	
	Ocean acidity	
	Ocean salinity	13
	Dissolved oxygen	14
Other	Air pollution weather	
	Atmospheric CO ₂ at surface	
	Radiation at surface	

High confidence of decrease

Medium confidence of decrease

Low confidence in direction of change

Medium confidence of increase

High confidence of increase

Table 8.1: Reproduction of column 1 of Table 12.12, IPCC AR6 Working Group I report. Emergence of anthropogenic signal in historical period for CIDs shown. White: no signal detected. Blue and orange: change detected (decrease or increase) and confidence level as indicated in color legend. Numbers refer to specific regions and confidence levels: see original Table for notes.

8.6 Extreme event attribution (EEA)

Examining the IPCC’s treatment of extreme weather and climate event attribution more broadly, AR6 provides an ambiguous assessment of the role of anthropogenic warming that differs between its chapters. Chapter 11 of WG1 states (Seneviratne *et al.*, 2021):

Evidence of observed changes in extremes and their attribution to human influence (including greenhouse gas and aerosol emissions and land-use changes) has strengthened since AR5, in particular for extreme precipitation, droughts, tropical cyclones and compound extremes (including dry/hot events and fire weather). Some recent hot extreme events would have been extremely unlikely to occur without human influence on the climate system.

By contrast, as noted in Section 8.5, Chapter 12 of WG1 (Table12.12) paints a different picture – presumably, the expert judgment of different groups of authors for the two chapters came to different conclusions (Ranasinghe, 2021):

- High confidence in an increase in extreme heat events in tropical regions where observations allow trend estimation and in most regions in the mid-latitudes, medium confidence elsewhere
- Medium confidence in a decrease in extreme cold events in Australia, Africa and most of northern South America where observations allow trend estimation
- No evidence of emergence in the historical period of a change in river floods, heavy precipitation, drought, fire weather, severe wind storms, and tropical cyclones

While the overall issue of detecting changes in extreme weather events and their attribution remains ambiguous, most of the activity in this area relates to the attribution of particular extreme weather events. The most prominent effort is World Weather Attribution (WWA; worldweatherattribution.org), an international research initiative for extreme event attribution that purports to analyze how climate change influences the likelihood and intensity of extreme weather events. Their approach is to use large ensembles of regional climate models to compare an event in today’s climate with one in a counterfactual pre-industrial climate without human influences.

WWA has a prominent public presence in linking extreme weather to climate change, with its press releases attracting considerable attention in public and policy discussions. However, the WWA’s extensive promotion of non-peer-reviewed findings, its open admission to shaping analyses to serve litigation, and its methodological challenges have sparked controversies, with critics questioning the robustness and impartiality of their conclusions (Pielke Jr. 2024). Despite these issues, WWA’s work continues to influence climate science and media narratives. Technical criticisms of the approach include a lack of a formal detection process; an implicit assumption that 100 percent of the post-industrial warming is caused by greenhouse gases; and a failure to adequately account for internal climate variability.

Because EEA is relatively new, many basic methodological issues have yet to be settled in the expert literature. An important challenge is the lack of data. Extreme events are by definition rare. Many analyses of extreme event types (including the U.S. National Assessment Reports) only evaluate data since 1950 or 1970. However, as emphasized by Chapter 6, many of the worst extreme weather and climate events in U.S. history occurred in the first half of the 20th century, including in the early 19th century. And if paleoclimate reconstructions are considered, it becomes very difficult for an event to pass thresholds of what is expected from natural variability, particularly if a reasonably sized geographic region is considered.

Another challenge is defining the event under study. There is a longstanding literature in statistics and econometrics on the challenge of analyzing data with outliers. The issue arises because a data series establishes a probability distribution defining the expected range of observations. If an outlier is observed it might indicate that the underlying process giving rise to the data distribution has changed (which in the weather context would mean that a climate change has been detected) or that the underlying process has multiple regimes each with a different probability distribution, in which case observing an outlier simply means we were temporarily in a different regime, but the system itself was unchanged. If a time series contains only a single outlier event at the end of the series it is not possible to determine which model is the correct one (Chen and Liu 1993). For instance, there might be an “ordinary” weather regime that yields a distribution of summer daytime highs in a particular coastal region, and a second “heatwave” regime that kicks in when an inland blocking event occurs, which yields a temperature distribution centered 15°C higher than the first one. A day with temperatures 13°C above normal would either be an extreme heat anomaly under the first regime or a somewhat cool event under the second, and we have no way in this case of knowing on statistical grounds which view is correct.

Visser and Petersen (2012) and Sardeshmukh *et al.* (2015) both point out that different distributions might fit observed data equally well but yield very different implications about the likelihood of a specific weather event. Visser and Petersen argue that, in view of the deep uncertainties of extreme weather analysis, drawing a connection between individual events and global climate change should be avoided. Furthermore, the existence of an outlier at the end of a data series poses the problem that estimates of the event probabilities will be biased whether the outlier is included or excluded (Barlow *et al.*, 2020). Methods to eliminate the bias have not yet been established, leading some experts (*e.g.* Miralles and Davison 2023) to argue that in settings in which a data series contains a single extreme event at the end, estimation of a return period for the extreme event will be so biased and uncertain that it should be avoided altogether.

We provide a case study of a recent high impact extreme event in the U.S. to illustrate the challenges and ambiguities in attributing the frequency and intensity of extreme weather events to human-caused warming.

8.6.1 Case study – 2021 Western North America heat wave

The 2021 Western North America heat wave was an extreme event that affected much of Western North America in late June 2021. Surface temperature records were set in Portland, OR (116°F; previous record 107°F) and Seattle, WA (108 °F; previous record 103°F) (Mass *et al.*, 2024).

The WWA team generated international headlines with their analysis, which provided the following attribution statements (WWA, 2021; Philip *et al.*, 2022):

- Based on observations and modeling, the occurrence of this heatwave was virtually impossible without human-caused climate change.
- The event is estimated to be about a one in 1000-year event in today’s climate.
- The event would have been at least 150 times rarer without human-induced climate change.
- This heatwave was about 2°C hotter than it would have been if it had occurred at the beginning of the industrial revolution (when global mean temperatures were 1.2°C cooler than today).

But an important counter to the first claim is that other researchers concluded from historical weather data that while a heat wave of the magnitude observed was indeed virtually impossible without anthropogenic climate change, it was also virtually impossible *with* climate change. Bercos-Hickey (2022) noted “these temperatures were virtually impossible under any previously experienced meteorological conditions, with or without global warming.” McKinnon and Simpson (2022) stated “the most likely explanation remains that the weather event itself was ‘bad luck.’” The 2023 Oregon Climate Assessment (Fleischman, 2023) concluded that the heat dome would have formed even without climate change and “There is no evidence that the highly unusual combination of weather features that drove the heat dome

were made more likely by climate change, and climate models do not project an increase in the frequency of high-pressure ridges over the Pacific Northwest” (Fleischman, 2023, p. 49).

Mass *et al.* (2024) summarizes the proximate sequence of compound events leading to the heat wave. There was a record-breaking mid-tropospheric ridge over the Pacific Northwest, forced by a tropical disturbance in the western Pacific. This produced record-breaking mid-tropospheric temperatures, strong subsidence in the lower atmosphere, low-level easterly flow that produced downslope warming on regional terrain and the removal of cooler marine air, and an approaching low-level trough that enhanced downslope flow. The event occurred at a time of maximum solar insolation, and drier-than-normal soil moisture. Using a storyline approach, Mass *et al.* assessed that there was no trend in drought and dry soils in the Pacific Northwest; there is no evidence of global warming producing stronger ridges of high pressure, and no observed trend in heat waves or record temperatures in the region. They concluded that although anthropogenic warming might have contributed as much as 2°F to the magnitude of the event, there is little evidence of further amplification of the event from increasing greenhouse gases.

Bercos-Hickey *et al.* (2022) conducted a statistical analysis of a model-based attribution study of this heat wave. Because the event is a far outlier and far above the bounds of Generalized Extreme Value distributions fitted from historical data, they concluded that estimates of return times, quantitative changes in event magnitude and frequency, and probability of the extreme temperatures such as provided by the WWA are not accurate and should be interpreted as having low confidence. They found that hindcast attribution methods using an ensemble of regional climate models, combined with Pearl (2009) causal inferences, can provide limited and conditional information about the magnitude of the human influence on this heatwave - they provided a more highly constrained estimate that human activities caused a ~1.4°F–1.8°F increase in the daily maximum temperatures.

Zeder *et al.* (2023) also concluded that the methods employed by Philip *et al.* (2022, the WWA analysis) tend to overstate the rarity of extreme heat waves, leading to a biased perception of the effect of climate change on the heatwave event: “The tendency to overestimate the return period of observed extreme heatwave events may fuel the impression that seemingly impossible heatwave extremes are currently clustering at an unprecedented rate.”

The 2021 Western North America heatwave was a rare and unprecedented compound weather event that broke century-old temperature records by as much as 9°F. While the WWA team received worldwide publicity for their rapid attribution claims blaming anthropogenic climate change, subsequent peer-reviewed analyses showed that the event was caused by rare meteorological conditions that were not made more probable by global warming.

References

- Allen, M. R., & Tett, S. F. B. (1999). Checking for model consistency in optimal fingerprinting. *Climate Dynamics*, 15(6), 419–434. <https://doi.org/10.1007/s003820050291>
- AR4: *Intergovernmental Panel on Climate Change Fourth Assessment Report* (2021) Working Group I Contribution. www.ipcc.ch.
- AR5: *Intergovernmental Panel on Climate Change Fifth Assessment Report* (2021) Working Group I Contribution. www.ipcc.ch.
- AR6: *Intergovernmental Panel on Climate Change Sixth Assessment Report* (2021) Working Group I Contribution. www.ipcc.ch.
- Arthun, M, RCJ Willis, HL Johnson, L Chafik, HR Langehaug (2021) Mechanisms of Decadal North Atlantic Climate Variability and Implications for the Recent Cold Anomaly. *Journal of Climate*, Vol 34

- Balcombe, Kevin, Iain Fraser and Abhijit Sharma (2019) Is radiative forcing cointegrated with temperature? A further examination using a structural time series approach.” *Management of Environmental Quality* 30(5) DOI 10.1108/MEQ-12-2018-021.
- Barlow, Anna Maria, Chris Sherlock and Jonathan Tawn (2020) Inference for Extreme Values Under Threshold-Based Stopping Rules. *Journal of the Royal Statistical Society Series C* August 2020, Pages 765–789, <https://doi.org/10.1111/rssc.12420>
- Bednarz, EW, AH Butler, X Wang, Z Zhou, W Yu, G Stenchikov, M Toohey, Y Zhu (2025) Indirect climate impacts of the Hunga eruption. Preprint, <https://doi.org/10.5194/egusphere-2025-1970>
- Bercos-Hickey, E., *et al.* (2022). Anthropogenic contributions to the 2021 Pacific Northwest heatwave. *Geophysical Research Letters*, 49(23), e2022GL099396. <https://doi.org/10.1029/2022GL099396>
- Bokuchava, DD and VV Semenov (2021) Mechanisms of the early 20th century warming in the Arctic. *Earth-Science Reviews*, 222, 103820
- Chatzistergos, T., Krivova, N. A., & Yeo, K. L. (2023). Long-term changes in solar activity and irradiance. *Journal of Atmospheric and Solar-Terrestrial Physics*, 252, 106150. <https://doi.org/10.1016/j.jastp.2023.106150>
- Chen, C., & Liu, L.-M. (1993). Joint estimation of model parameters and outlier effects in time series. *Journal of the American Statistical Association*, 88(421), 284–297. <http://www.jstor.org/stable/2290724>
- Chen, H, SX Chen and M Mu (2023) A statistical review on the optimal fingerprinting approach in climate change studies. *Climate Dynamics* 62:1439-1446 <https://link.springer.com/article/10.1007/s00382-023-06975-5>
- Chylek, P., *et al.* (2016). The role of Atlantic Multi-Decadal Oscillation in the global mean temperature variability. *Climate Dynamics*, 47(9–10), 3271–3279. <https://doi.org/10.1007/s00382-016-3025-7>
- Connolly, R., *et al.* (2021). How much has the sun influenced Northern Hemisphere temperature trends? An ongoing debate. *Research in Astronomy and Astrophysics*, 21(6), 131. <https://doi.org/10.1088/1674-4527/21/6/131>
- Cramer, W., *et al.* (2014). Detection and attribution of observed impacts. In C. B. Field *et al.* (Eds.), *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects* (pp. 979–1037). Cambridge University Press.
- Dagsvik, JK SH Moen (2023) To what extent are temperature levels changing due to greenhouse gas emissions? Statistics Norway Discussion Paper 1007 September 2023
- Dagsvik, JK, M Fortuna and SH Moen (2020) How Does Temperature Vary Over Time?: Evidence on the Stationary and Fractal Nature of Temperature Fluctuations. *Journal of the Royal Statistical Society Series A: Statistics in Society*, Volume 183, Issue 3, June 2020, Pages 883–908, <https://doi.org/10.1111/rssa.12557>
- Davidson, James E.H., David B. Stephenson and Alemtsehai A. Turasie (2015) Time series modeling of paleoclimate data *Environmetrics* 27(1) <https://onlinelibrary.wiley.com/doi/abs/10.1002/env.2373>
- Dergiades, T., R.K. Kaufmann and T. Panagiotidis (2016). Long-run changes in radiative forcing and surface temperature: The effect of human activity over the last five centuries. *Journal of Environmental Economics and Management* 76:67—85. <https://www.sciencedirect.com/science/article/abs/pii/S0095069615000911>
- England, Mark, Lorenzo M Polvani, James A Screen, *et al* (2025). Surprising, but not unexpected, multi-decadal pause in Arctic sea ice loss. *ESS Open Archive*. March 29, 2025.
- Fleishman, E., editor. (2023). Sixth Oregon Climate Assessment. Oregon Climate Change Research Institute, Oregon State University, Corvallis, Oregon. <https://blogs.oregonstate.edu/occric/oregon-climate-assessments>

- Forster et al. (2024) “Indicators of Global Climate Change 2024: annual update of key indicators of the state of the climate system and human influence” *Earth System Science Data* 17(6) <https://doi.org/10.5194/essd-17-2641-2025>
- Frajka-Williams, E, C Beaulieu, Aurelie Duchez (2017) Emerging negative Atlantic Multidecadal Oscillation index in spite of warm subtropics. *Scientific Reports*, 7:11224
- Goessling, Helge F. Thomas Rackow and Thomas Jung (2024) Recent global temperature surge intensified by record-low planetary albedo. *Science* Vol 387, Issue 6729 pp. 68-73 <https://www.science.org/doi/abs/10.1126/science.adq7280>
- Gulev, S. K., *et al.* (2021). Changing state of the climate system. In V. Masson-Delmotte *et al.* (Eds.), *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 287–422). Geneva: Intergovernmental Panel on Climate Change.
- Hansen J and P Karecha (2025) Large cloud feedback confirms high sensitivity. <https://www.columbia.edu/~jeh1/mailings/2025/CloudFeedback.13May2025.pdf>
- Hegerl, G. C., *et al.* (2007). Understanding and attributing climate change. In S. Solomon *et al.* (Eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 663–745). Cambridge University Press.
- Hegerl, GC, S Bronniman, A Schurer *et al.* (2017) Early 20th century warming: Anomalies, causes and consequences. *WIREs Climate Change*, DOI: 10.1002/wcc.522
- Hegerl, GC, S Bronniman, T Cowan (2019) Causes of climate change over the historical record. *Environmental Research Letters*, 14, 123006
- Hodnebrog, Ø *et al.* (2024), Recent reductions in aerosol emissions have increased Earth’s energy imbalance. *Communications Earth and Environment* 5, 166 (2024). doi:10.1038/s43247-024-01324-8
- Houghton, J. T., *et al.* (Eds.). (2001). *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 1–881). Cambridge University Press. <https://mailchi.mp/caa/large-cloud-feedback-confirms-high-climate-sensitivity>
- Kravtsov, S., Westgate, A., & Gavrilov, A. (2024). Global-scale multidecadal variability in climate models and observations, part II: The stadium wave. *Climate Dynamics*, 62, 10281–10306. <https://doi.org/10.1007/s00382-024-07451-4>
- Kuchar, A., Sukhodolov, T., Chiodo, G., Jörmann, A., Kult-Herdin, J., Rozanov, E., and Rieder, H. H. (2025): Modulation of the northern polar vortex by the Hunga Tonga–Hunga Ha’apai eruption and the associated surface response, *Atmospheric Chemistry and Physics*, 25, 3623–3634, <https://doi.org/10.5194/acp-25-3623-2025>.
- Lewis, Nicholas (2023) “Objectively Combining Climate Sensitivity Evidence,” *Climate Dynamics* 61, no. 9–10 (2023): 3155–3163, <https://doi.org/10.1007/s00382-022-06398-8> .
- Lockwood, M. (2012). Solar influence on global and regional climates. *Surveys in Geophysics*, 33(3–4), 503–534. <https://doi.org/10.1007/s10712-012-9181-3>
- Loeb, N. G. *et al.* (2019) Decomposing Shortwave Top-of-Atmosphere and Surface Radiative Flux Variations in Terms of Surface and Atmospheric Contributions. *Journal of Climate* 32(16), 5003-5019 (2019). doi:10.1175/JCLI-D-18-0826.1

- Loeb, N., S-H Ham, RP Allen, TJ Thorsen, B Meyssignac, S Kato, GC Johnson, JL Lyman (2024) Observational Assessment of Changes in Earth’s Energy Budget since 2000. *Surveys in Geophysics*, 45, 1757-1783
- Mass, C., *et al.* (2024). The Pacific Northwest heat wave of 25–30 June 2021: Synoptic/mesoscale conditions and climate perspective. *Weather and Forecasting*, 39(2), 275–291.
<https://doi.org/10.1175/WAF-D-23-0154.1>
- Matthes, K., *et al.* (2017). Solar forcing for CMIP6 (v3.2). *Geoscientific Model Development*, 10(6), 2247–2302. <https://doi.org/10.5194/gmd-10-2247-2017>
- McKinnon, K. and I. Simpson (2022) How Unexpected Was the 2021 Pacific Northwest Heatwave? *Geophysical Research Letters*, 49, e2022GL100380. <https://doi.org/10.1029/2022GL100380>
- McKinnon, K. and I. Simpson (2022) How Unexpected Was the 2021 Pacific Northwest Heatwave? *Geophysical Research Letters*, 49, e2022GL100380. <https://doi.org/10.1029/2022GL100380>
- McKittrick, R. (2021). Checking for model consistency in optimal fingerprinting: A comment. *Climate Dynamics*, 58(1–2), 405–411. <https://doi.org/10.1007/s00382-021-05913-7>
- McKittrick, R. (2023). Total least squares bias in climate fingerprinting regressions with heterogeneous noise variances and correlated explanatory variables. *Environmetrics*, 35(2), e2835.
<https://doi.org/10.1002/env.2835>
- McKittrick, Ross R (2022) On the choice of TLS versus OLS in climate signal detection regression. *Climate Dynamics* 10.1007/s00382-022-06315-z
- McKittrick, Ross R (2025) Consistent Climate Fingerprinting. *Climate Dynamics* forthcoming preprint available at www.rossmckittrick.com
- McLean, J.D., C. R. de Freitas, and R. M. Carter, (2009) “Influence of the Southern Oscillation on Tropospheric Temperature,” *Journal of Geophysical Research: Atmospheres* 114, no. D14 (July 23, 2009): D14104, <https://doi.org/10.1029/2008JD011637>.
- Miralles, Ophélie and Anthony Davison (2023) Timing and spatial selection bias in rapid extreme event attribution. *Weather and Climate Extremes* 41 100584 <https://doi.org/10.1016/j.wace.2023.100584>
- Myhre, G., D. Shindell, F.-M. Bréon, *et al* (2013): Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., *et al.* (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Nissan, Hannah, Lisa Goddard, Erin Coughlan de Perez, *et al.* (2019) On the use and misuse of climate change projections in international development. *Wires Climate Change* 14 March 2019
<https://doi.org/10.1002/wcc.579>
- Palmer, Tim N and Bjorn Stevens (2019) The scientific challenge of understanding and estimating climate change. *Proceedings of the National Academy of Sciences* 116 (49) 24390-24395
<https://doi.org/10.1073/pnas.1906691116>
- Pearl, J. (2009). *Causality*. Cambridge University Press.
- Philip, S. Y., *et al.* (2022). Rapid attribution analysis of the extraordinary heatwave on the Pacific coast of the U.S. and Canada June 2021. *Earth System Dynamics*, 13(4), 1689–1713.
<https://doi.org/10.5194/esd-13-1689-2022>
- Pielke Jr., Roger (2024) Weather Attribution Alchemy. Substack essay
<https://rogerpielkejr.substack.com/p/weather-attribution-alchemy>
- Ranasinghe, R., Ruane, A. C., Vautard, R., Arnell, N., Coppola, E., Cruz, F. A., ... Zaaboul, R. (2021). Chapter 12: Climate change information for regional impact and for risk assessment. In V. Masson-Delmotte *et al.* (Eds.), *Climate Change 2021: The Physical Science Basis. Contribution of Working*

- Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (pp. 1767–1926). Cambridge University Press. <https://doi.org/10.1017/9781009157896.014>
- Razzak, Weshah A. (2022). "The Econometrics of Global Warming," *Journal of Economics and Econometrics*, vol. 65(2), pages 13-47.
- Sardeshmukh, P. D., Compo, G. P., & Penland, C. (2015). Need for caution in interpreting extreme weather statistics. *Journal of Climate*, 28(23), 9166–9187. <https://doi.org/10.1175/JCLI-D-15-0020.1>
- Scafetta, N. (2013). Discussion on climate oscillations: CMIP5 general circulation models versus a semi-empirical harmonic model based on astronomical cycles. *Earth-Science Reviews*, 126, 321
- Scaffeta, Nicola (2021) "Reconstruction of the Interannual to Millennial Scale Patterns of the Global Surface Temperature," *Atmosphere* 12, no. 2: 147, <https://doi.org/10.3390/atmos12020147>
- Schmidt, G. (2024) Climate models can't explain 2023's huge heat anomaly — we could be in uncharted territory. *Nature* 627, 467. <https://www.nature.com/articles/d41586-024-00816-z>
- Schmutz, Werner K. (2021) "Changes in the Total Solar Irradiance and Climatic Effects," *Journal of Space Weather and Space Climate* 11: 40, <https://doi.org/10.1051/swsc/2021016>.
- Seneviratne, S. *et al.* (2021) "Chapter 11: Weather and Climate Extreme Events in a Changing Climate." In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by V. Masson-Delmotte, P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, *et al.*, 1513–1766. Cambridge: Cambridge University Press, 2021. <https://doi.org/10.1017/9781009157896.013>.
- Solanki, S.K. *et al.* (2004), "Unusual Activity of the Sun during Recent Decades Compared to the Previous 11,000 Years," *Nature* 431, no. 7012 (October 28, 2004): 1084–1087, <https://doi.org/10.1038/nature02995>.
- Stefani, Frank (2021) "Solar and Anthropogenic Influences on Climate: Regression Analysis and Tentative Predictions," *Climate* 9, no. 11 (November 3, 2021): 163, <https://doi.org/10.3390/cli9110163>.
- Thompson, DJW, JM Wallace, JJ Kennedy, PD Jones (2010) An abrupt drop in Northern Hemisphere Sea Surface Temperature around 1970. *Nature*, 467, 444-447
- Tokinaga, H, SP Xie, and H Mukougawa (2017) Early 20th century Arctic warming intensified by Pacific and Atlantic multidecadal variability. *Proceedings of the National Academy of Sciences*, 114, 6227-6233 <https://www.pnas.org/doi/abs/10.1073/pnas.1615880114>
- Tung, Ka-Kit and Jiansong Zhou (2013) "Using Data to Attribute Episodes of Warming and Cooling in Instrumental Records," *Proceedings of the National Academy of Sciences* 110, no. 6 (January 23, 2013): 2058–2063, <https://doi.org/10.1073/pnas.1212471110>.
- Usoskin, I.G., S. K. Solanki, and G. A. Kovaltsov (2007) "Grand Minima and Maxima of Solar Activity: New Observational Constraints," *Astronomy & Astrophysics* 471, no. 1 (June 6, 2007): 301–309, <https://doi.org/10.1051/0004-6361:20077704>.
- Velasco Herrera, V.M., B. Mendoza, and G. Velasco Herrera (2015) "Reconstruction and Prediction of the Total Solar Irradiance: From the Medieval Warm Period to the 21st Century," *New Astronomy* 34 (January 2015): 221–233, <https://doi.org/10.1016/j.newast.2014.07.009>.
- Visser, H. and A.C. Petersen (2012) Inferences on weather extremes and weather-related disasters: a review of statistical methods *Climates of the Past* 8, 265–286 <https://cp.copernicus.org/articles/8/265/2012/cp-8-265-2012.pdf>

- World Weather Attribution (2021) “Western North American Extreme Heat Virtually Impossible without Human-Caused Climate Change.” July 7, 2021. <https://www.worldweatherattribution.org/western-north-american-extreme-heat-virtually-impossible-without-human-caused-climate-change/>.
- Yuan, *et al.* (2024) Abrupt reduction in shipping emission as an inadvertent geoengineering termination shock produces substantial radiative warming. *Communications Earth and Environment* 5, 281 (2024). doi:10.1038/s43247-024-01442-3
- Zeder, J., Sippel, S., Pasche, O. C., Engelke, S., & Fischer, E. M. (2023). The effect of a short observational record on the statistics of temperature extremes. *Geophysical Research Letters*, 50, e2023GL104090. <https://doi.org/10.1029/2023GL104090>

PART III: IMPACTS ON ECOSYSTEMS AND SOCIETY

9 CLIMATE CHANGE AND U.S. AGRICULTURE

Chapter summary

There has been abundant evidence going back decades that rising CO₂ levels benefit plants, including agricultural crops, and that CO₂-induced warming will be a net benefit to U.S. agriculture. The increase in ambient CO₂ has also boosted productivity of all major U.S. crop types. There is reason to conclude that on balance climate change has been and will continue to be neutral or beneficial for most U.S. agriculture.

9.1 Econometric analyses

Econometric analyses of the effects of climate change on agriculture have sought to integrate information about long term crop yield changes in response to temperature and precipitation changes under assumptions about adaptive behavior by farmers. One method focuses on variations in agricultural land values. The rationale is that if climate change is a long-term net benefit for agriculture it should be capitalized into higher market values for agricultural land, and vice versa. While individual crops might benefit from, or be harmed by, climate change, once the adaptative responses of farmers are considered the value of farmland represents an index of whether the changes are expected to be beneficial or not. Mendelsohn *et al.* (1994) examined the relationship between historical climatic variations on agricultural land values and concluded global warming would be slightly beneficial to American agriculture.

The Mendelsohn *et al.* method, called Ricardian analysis after David Ricardo, the 19th century British economist who pioneered the study of land values, attracted subsequent criticism from authors who argued it failed to account for differences in land values attributable to fixed locational characteristics like soil quality, and non-climatic changes including nearby urbanization. Deschênes and Greenstone (2007) looked at agricultural profits instead of land values and reached conclusions similar to Mendelsohn *et al.* (1994), namely that past climatic variations had relatively little effect on farm profitability and that warming would likely yield small overall benefits for the U.S. agricultural sector. However, a subsequent exchange with critics led them (Deschênes and Greenstone 2012) to revise their conclusions and project potentially large losses in U.S. agriculture due to climate warming.

Burke and Emerick (2016) looked at temperature variations over 1980 to 2000 and argued that farmers were not as able to adapt to temperature changes as the Ricardian method assumes and that climate change would have large negative impacts on corn and soy yields. Schlenker and Roberts (2009) similarly argued that yield gains to past warming would not carry over to the future and corn and soy yields would sharply decrease this century due to climate change.

Two recent studies have argued that pessimistic findings such as these are not robust. Ortiz-Bobea (2019) argued that land values aggregate farm and non-farm influences and the latter need to be filtered out. He developed a data set using cash rents for agricultural activity as a measure of land value specifically for farming activity. Whereas the land value model implied future losses under climate warming, the same model estimated using cash rents did not, leading the author to conclude the pessimistic results were due to using an inaccurate measure of the returns to farming activity. Bareille and Chakir (2023) assembled a large data base on farm sale prices in France for properties that sold twice between 1996 and 2019. They could replicate pessimistic results showing negative effects of warming on agricultural land values using conventional econometric modeling. But by taking advantage of the repeat sales data, which provides information on site-specific changes in land prices, they found the results reversed and implied that climate change will be very beneficial for French agriculture. The authors concluded that, taking adaptation into account, a warming climate would yield positive benefits for French agriculture that were between two and

20 times larger than had previously been estimated. On average, with full adaptation, they concluded that climate changes under the medium RCP4.5 scenario could double the value of French farmland by 2100.

A major deficiency of all these studies, however, is that they omit the role of CO₂ fertilization. Climate change as it relates to this report is caused by GHG emissions, chiefly CO₂. The econometric analyses referenced above focus only on temperature and precipitation changes and do not take account of the beneficial growth effect of the additional CO₂ that drives them. As explained in Chapter 2, CO₂ is a major driver of plant growth, so this omission biases the analysis towards underestimation of the benefits of climate change to agriculture.

9.2 Field and laboratory studies of CO₂ enrichment

One of the ways the effect of CO₂ on crop growth has been studied is through “free air enrichment experiments” or FACE plots, in which small sources of CO₂ are placed in fields surrounding plants and the growth response to elevated CO₂ under varying weather conditions are recorded. Ainsworth *et al.* (2020) summarizes results from about 250 such studies. They found that elevation of CO₂ by 200 ppm caused an average 18 percent increase in crop yield in C3 plants. C4 plants exhibited benefits mainly under drought conditions.

In addition to FACE plot experiments there have been thousands of laboratory experiments on the effects of CO₂ on all kinds of plant growth. Here we review some of the results on key U.S. agricultural crops.

Soybean

Studies on the impact of elevated CO₂ on Soybean (*Glycine max* (L.) Merr.) plants in the water-deficient region of Huang-Huai-Hai Plain, China, showed that elevated CO₂ concentrations improved photosynthesis rate, water use efficiency, and growth Li (2013) under both normal conditions and drought conditions. The CO2Science.org website reports on 108 published experiments between 1985 and 2019 exposing soybean to enriched CO₂ levels. Converted to a +300ppm common scale the average growth benefit was +50.9 percent. There were also ten studies reporting on +600 ppm CO₂ enrichment, which increased photosynthesis by an average of 90.3%.

Maize (corn)

The CO2Science.org website reports on 28 published experiments between 1983 and 2018 exposing corn (*Zea mays* L.) to enriched CO₂ levels. Converted to a +300ppm common scale the average growth benefit was +23.7 percent. Corn also benefits from increased drought tolerance under elevated CO₂. An experimental study (Allen Jr., 2011) exposed plants to water stress conditions in sunlit controlled-environment chambers at 360 ppm (ambient) and 720 ppm (elevated) CO₂. The drought stress caused a 41 percent loss of growth under ambient CO₂ but only a 13 percent loss under elevated CO₂.

Wheat

The CO2Science.org website reports on 92 published experiments between 1983 and 2020 exposing common wheat (*Triticum aestivum* L.) to enriched CO₂ levels. Converted to a +300ppm common scale the average growth benefit was +67.6 percent. Blandino (2020) measured both the yield and nutritional quality of an “improver” hybrid wheat variety and its parents under elevated CO₂ levels (+166 ppm). They reported a grain yield increase of +16 percent but a 7 percent decrease in grain protein levels. But they also found that the food quality of different wheat varieties responded differently to elevated CO₂ levels, so that with proper varietal selection, growers could select wheat types that best take advantage of the elevated CO₂.




Figure 9.1: U.S. average CO₂ levels and yields of corn, soy and wheat all normalized so 1940=100. Source: Taylor and Schlenker (2021)

Further evidence

A 2021 report from the U.S. National Bureau of Economic Research (Taylor and Schlenker 2021) used satellite-measured observations of outdoor CO₂ levels across the United States, matched to county-level agricultural output data and other economic variables. After controlling for the effects of weather, pollution and technology the authors concluded that CO₂ emissions had boosted U.S. crop production since 1940 by 50 to 80 percent, attributing much larger gains than had previously been estimated using FACE experiments. They found that every ppm of increase in CO₂ concentration boosts corn yields by 0.5 percent, soybeans by 0.6 percent, and wheat by 0.8 percent.

Beyond growth benefits, extra CO₂ boosts plant resilience to dryness. Deryng *et al.* (2016) surveyed evidence on crop water productivity (CWP, the yield per unit of water used), drawing attention to the potential for CO₂ both to enhance photosynthesis and reduce leaf-level transpiration (water loss during leaf respiration). They surveyed all available FACE data on crop yield changes for maize (corn), wheat, rice and soybean, and combined it with crop model data simulating yield responses as of 2080 under the extreme RCP8.5 scenario in four growing regions (Tropics, Arid, Temperate and Cold) each of which were split into rainfed and irrigated sub-regions. In every region they reported that models without CO₂ fertilization predicted CWP losses but CO₂ fertilization more than offset them; once fertilization was taken into account, all regions showed a net CWP gain. Deryng *et al.* also reported that negative yield impacts on wheat and soybean were fully offset by CWP gains and mitigated by up to 90 percent for rice and 60 percent for maize. Similarly, Cheng *et al.* (2017) noted that increased global plant growth from 1982 to 2011 due to rising CO₂ uptake was accompanied by such large gains in CWP that despite the extra biomass global water use by plants had not increased.

Deryng *et al.* (2016) assumed that climate change would “exacerbate water scarcity”. It is noteworthy that while models predict drylands will expand under climate warming, observations show the opposite: greening is happening even in arid areas. Zhang *et al.* (2024) report that due to increased CO₂ levels “increasing aridity in drylands won’t lead to a general loss of vegetation productivity”; increased desertification will happen in at most 4 percent of currently arid areas.

9.3 Crop modeling meta-analyses

Notwithstanding the abundant evidence for the direct benefits of CO₂ and of CO₂-induced warming on crop growth, in 2023 the U.S. Environmental Protection Agency (EPA 2023) boosted its estimate of the Social Cost of Carbon (SCC) about five-fold based largely on a very pessimistic 2017 estimate of global agricultural damages from climate warming (Moore *et al.*, 2017). One of the two damage models used by the EPA attributed nearly half of the 2030 SCC to projected global agricultural damages based on the Moore *et al.* (2017) analysis. This study was a meta-analysis of crop model studies simulating yield changes for agricultural crops under various climate warming scenarios. Moore *et al.* projected declining global crop yields for all crop types in all regions due to warming.




Figure 9.2 Crop yields under CO₂-induced climate warming. Blue: as published in Moore *et al.* (2017). Green: after including omitted data. Source: McKittrick (2025).

McKittrick (2025) re-examined the Moore *et al.* database and found that, while it claimed to cover 1,722 studies, only half the entries ($N=862$) had complete records, so that the sample available for regression analysis was much smaller than both studies indicated. McKittrick noted that the records most commonly missing were the changes in ambient CO₂ and found that in many cases these could be recovered from the underlying studies or the original climate scenario tables, thereby increasing the usable sample size by 40 percent. The crop yield projections incorporating the newly available data changed considerably. As shown in Figure 9.2, whereas the partial data set implied warming would decrease yield (blue lines), the complete data set implied constant or increase global yields, even out to 5°C warming (green lines).

9.4 CO₂ fertilization and nutrient loss

Evidence has shown that CO₂-induced biomass gains are sometimes accompanied by reductions in the concentrations of protein and other key nutrients such as iron and zinc (Ebi *et al.* 2021). Some experimental evidence has shown that the rising temperatures expected to accompany higher CO₂ levels will offset this loss (Köhler *et al.* 2019) although the evidence for this is mixed, as is the evidence that nutrient dilution observed to date is entirely attributable to higher CO₂ (Ziska 2022). If nutrient dilution does occur under rising CO₂ levels there are several adaptive strategies that could be pursued.

First, selective breeding to raise micronutrient content is already long established (Saltzman *et al.* 2017) and has proven to be a cost-effective agronomic strategy (Ebi *et al.* 2021). Strategies can include both conventional breeding and genetically-modified organisms. An example of the latter is Golden Rice, which contains elevated levels of beta-carotene to boost biosynthesis of vitamin A in the human body. Optimal strategies will be location-specific because they vary by crop, climate and soil type (Ebi *et al.* 2021). Second, fortification of food products with micronutrients is already routine. Folic acid (a B vitamin) is added to flour and many other foods; iodine is added to table salt, most commercial breakfast cereals are fortified with iron and numerous vitamins, *etc.* Third, dietary supplements in the form of multivitamin tablets are inexpensive, widely-available and routinely consumed.

One concern about reliance on adaptive strategies is whether they are feasible in low-income countries. Micronutrient deficiency is already a problem in the developing world and dietary supplements have proven to be an effective low-cost response (Ebi *et al.* 2021). It should also be noted the IPCC emission scenarios that generate high levels of warming do so on the basis of high levels of income growth. The SSP scenarios³ assume that, compared to 2005 levels, global per capita income will double by 2100 in the lowest growth case (SSP3), and in the highest emission case (SSP5) global per-capita income will grow nearly 16-fold. In that scenario even the poorest regions (Africa and the Middle East) end up with a per capita income of about US\$126,000, 70 percent higher than current U.S. per capita income (about US\$75,000). Consequently the same scenarios in which CO₂ levels increase the most are also those in which global poverty is largely eliminated, in which case all countries would be able to afford dietary supplements as necessary to address micronutrient deficiencies, if they arise and cannot be addressed using on-farm agricultural strategies.

In summary, there is abundant evidence going back decades that rising CO₂ levels benefit plants, including agricultural crops, and that CO₂-induced warming will be a net benefit to U.S. agriculture. To the extent nutrient dilution occurs there are mitigating strategies available that will need to be researched and adapted to local conditions.

References

- Ainsworth, Elizabeth and Stephen P. Long (2020) 30 years of free-air carbon dioxide enrichment (FACE): What have we learned about future crop productivity and its potential for adaptation? *Global Change Biology* <https://doi.org/10.1111/gcb.15375>
- Allen Jr, L. K. (2011). Elevated CO₂ increases water use efficiency by sustaining photosynthesis of water-limited maize and sorghum. *Journal of Plant physiology*, 168(16), 1909-1918.
- Bareille, F. and R. Chakir (2023) The impact of climate change on agriculture: A repeat-Ricardian analysis. *Journal of Environmental Economics and Management* 119
- Blandino, M. B. (2020). Elevated CO₂ impact on common wheat (*Triticum aestivum* L.) yield, wholemeal quality, and sanitary risk. *Journal of agricultural and food chemistry*, 68(39), 10574-10585.

³ See compilation at [Our World in Data](#).

- Burke, Marshall and Kyle Emerick (2016) Adaptation to Climate Change: Evidence from US Agriculture. *American Economic Journal: Economic Policy* 8(3) August 2016 106–140 <https://www.aeaweb.org/articles?id=10.1257/pol.20130025>
- Cheng, L., Zhang, L., Wang, YP. *et al.* Recent increases in terrestrial carbon uptake at little cost to the water cycle. *Nature Communications* 8, 110 (2017). <https://doi.org/10.1038/s41467-017-00114-5>
- CO2Science.org (undated) Plant growth archive http://www.co2science.org/data/plant_growth/dry/dry_subject_b.php
- Deryng, D., Elliott, J., Folberth, C. *et al.* (2021) Regional disparities in the beneficial effects of rising CO₂ concentrations on crop water productivity. *Nature Climate Change* 6, 786–790 (2016). <https://doi.org/10.1038/nclimate2995>
- Deschênes, Olivier and Michael Greenstone (2012) The Economic Impacts of Climate Change: Evidence from Agricultural Output and Random Fluctuations in Weather: Reply to Comment *American Economic Review*. 102, no. 7: 3761–3773. <http://hdl.handle.net/1721.1/82641>
- Ebi, Kristie L, C Leigh Anderson, Jeremy J Hess, *et al.* (2021) Nutritional quality of crops in a high CO₂ world: an agenda for research and technology development. *Environmental Research Letters* 16(6) 064045 <https://iopscience.iop.org/article/10.1088/1748-9326/abfcfa>
- Köhler, Iris H., Steven C. Huber, Carl J. Bernacchi, Ivan R. Baxter (2019) Increased temperatures may safeguard the nutritional quality of crops under future elevated CO₂ concentrations. *The Plant Journal* 97(5) 872 – 886 <https://onlinelibrary.wiley.com/doi/epdf/10.1111/tpj.14166>
- Li, D. L. (2013). Effects of elevated CO₂ on the growth, seed yield, and water use efficiency of soybean (*Glycine max* (L.) Merr.) under drought stress. *Agricultural Water Management*, 129, 105–112.
- McKittrick, Ross R. (2025) Extended Crop Yield Meta-analysis Data do not Support Upward SCC Revision. *Scientific Reports* 15 article 5575 <https://www.nature.com/articles/s41598-025-90254-2>
- Mendelsohn, R., Nordhaus, W. D., & Shaw, D. (1994). The impact of global warming on agriculture: a Ricardian analysis. *The American Economic Review*, Sept 1994 753–771.
- Moore FC, Baldos U, Hertel T, Diaz D (2017) New science of climate change impacts on agriculture implies higher social cost of carbon. *Nature Communications* 8:1607. <https://doi.org/10.1038/s41467-017-01792-x>
- Ortiz-Bobea, Ariel (2019) The role of nonfarm influences in ricardian estimates of climate change impacts on US agriculture. *American Journal of Agricultural Economics*. 102 (3), 934–959.
- Saltzman, Amy, Ekin Birol, Adewale Oparinde *et al.* (2017) Availability, production, and consumption of crops biofortified by plant breeding: current evidence and future potential. *Annals of the New York Academy of Sciences* Volume 1390, Issue1 <https://doi.org/10.1111/nyas.13314>
- Schlenker, Wolfram and Michael Roberts (2009) Nonlinear temperature effects indicate severe damages to U.S. crop yields under climate change. *Proceedings of the National Academy of Sciences* 106 (37) 15594–15598 <https://doi.org/10.1073/pnas.0906865106>
- Taylor, Charles and Wolfram Schlenker (2023) Environmental drivers of agricultural productivity growth: CO₂ fertilization of US field crops. National Bureau of Economic Research Working paper 29320, <https://www.nber.org/papers/w29320>
- US Environmental Protection Agency (2023) Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, “Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review”, EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances, Nov. 2023, https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf
- Zhang *et al.* (2024) Less than 4% of dryland areas are projected to desertify despite increased aridity under climate change. *Nature Communications Earth and Environment* 5 June 2024 <https://www.nature.com/articles/s43247-024-01463-y>
- Ziska, Lewis H (2022) Rising Carbon Dioxide and Global Nutrition: Evidence and Action Needed. *Plants* 1(7), 1000; <https://doi.org/10.3390/plants11071000>

10 MANAGING RISKS OF EXTREME WEATHER

Chapter summary

Trends in losses from extreme weather and climate events are dominated by population increases and economic growth. Technological advances such as improved weather forecasting and early warning systems have substantially reduced losses from extreme weather events. Better building codes, flood defenses, and disaster response mechanisms have lowered economic losses relative to GDP. The U.S. economy's expansion has diluted the relative impact of disaster costs, as seen in the comparison of historical and modern GDP percentages. Heat-related mortality risk has dropped substantially due to adaptive measures including the adoption of air conditioning, which relies on the availability of affordable energy. U.S. mortality risks even under extreme warming scenarios are not projected to increase if people are able to undertake adaptive responses.

10.1 Socioeconomic context

Risks from human-caused climate change are affected by natural weather and climate variability and are dominated by the exposure of wealth in coastal and other disaster-prone regions and vulnerabilities of poorer populations. The evolution of climate risk in the U.S. has been dominated by societal factors, rather than by changes to the actual weather and climate hazards. Deaths from weather disasters have decreased substantially since 1900, even as U.S. population grew from 76 million in 1900 to over 331 million in 2020 (Goklany 2011). For example, the Galveston hurricane killed over 8,000 people in 1900 (0.01 percent of the U.S. population), whereas the worst recent disaster, Hurricane Katrina in 2005, killed 1800 people (or 0.0006 percent of the U.S. population) (NOAA National Hurricane Center 2025; U.S. Census Bureau 2025).

Technological advances have substantially reduced losses from extreme weather events. Early warning systems, satellite monitoring, and improved weather forecasting have reduced deaths, although exact numbers are hard to quantify (Deryugina and Hsiang 2023). U.S. weather forecasting has been estimated to reduce losses from weather events with an annualized benefit of \$31.5B, protecting lives, property, and supporting agriculture and transportation (NRC 2010). Improved hurricane forecasts have reduced pre- and post-landfall spending, with annual per hurricane cost reductions estimated at \$5B (Molina and Rudik 2024).

Infrastructure improvements have contributed to substantial reductions in losses from extreme weather events. Building codes, such as those implemented in Florida after Hurricane Andrew (1992), have reduced losses by ensuring structures can withstand high winds and floods. Homes built after 2002 showed minimal damage during Hurricane Michael (2018), unlike older homes (FEMA 2020). Sea walls, like the Galveston Seawall, protect against wave action and storm surge. The New Orleans Hurricane and Storm Damage Risk Reduction System successfully mitigated storm surge during Hurricane Isaac (2012) (Battelle Memorial Institute 2013). Inland dams in the U.S. help control flooding by storing excess water during heavy rains. It is estimated that the Tennessee Valley Authority (TVA) dams prevent about \$309M in annual flood damage in the TVA region and along the Ohio and Mississippi Rivers (TVA, 2025). During Hurricane Helene (2024), TVA's strategies prevented approximately \$406M in potential damages (APPA 2024).

10.2 Data challenges

Since 1980, NOAA has provided a count of annual U.S. weather-related disasters that it estimates to have exceeded \$1 Billion (inflation adjusted), showing a substantial increase starting in 2008. NOAA and other government officials have cited the upward trend in the Billion Dollar Disaster series as evidence that climate change is making extreme weather worse (Pielke Jr., 2024). But over time, population and wealth have increased dramatically in the U.S., so when an extreme weather or climate event occurs, there is more damage even if there is no underlying trend in the frequency or intensity of extreme weather. Pielke Jr. (2024) demonstrates that losses per weather disaster as a proportion of GDP have decreased by about 80 percent since 1980, as shown in Figure 10.1. Pielke Jr. (2024) also argued that in addition to relying on opaque data sources and unreported adjustments, NOAA failed to normalize its Billion Dollar Disaster data series properly for changes in population exposure and wealth. In May 2025 NOAA announced it has withdrawn the Billion Dollar Disaster product from publication (Pielke Jr., 2025).




Figure 10.1: Losses per disaster as a % of Gross Domestic Product in NOAA's billion-dollar disaster dataset (the version downloaded in July 2023), 1980 to 2022. Source: (Pielke, Jr. 2024)

In summary, trends in losses from extreme weather and climate events are dominated by population increases and economic growth. Technological advances such as improved weather forecasting and early warning systems have substantially reduced losses from extreme weather events. Better building codes, flood defenses, and disaster response mechanisms have lowered economic losses relative to GDP. Finally, the U.S. economy's expansion has diluted the relative impact of disaster costs, as seen in the comparison of historical and modern GDP percentages.

10.3 Mortality from temperature extremes

10.3.1 Heat and cold risks

Changes in temperature extremes are among the most certain impacts expected in a warming world. It stands to reason that extreme heat events would likely become more frequent while extreme cold events would become less frequent. This pattern is evident in the historical period, though not in the continental U.S. (Chapter 6), and is expected to continue with further warming.

Mortality during heat extremes is typically caused by heat stroke and heat exhaustion, while mortality during cold extremes typically stems from hypothermia and heart strain. Global mortality is substantially greater for cold conditions than for hot conditions (Zhao et al 2021, Ritchie 2024). Unlike with heat-related mortality, cold-weather risks set in even at moderately cold conditions (Gasparini *et al.* 2015, Lee and Dessler 2023). The U.S. EPA (citing data from the Centers for Disease Control) reports that on average over 1999 to 2015 there were 2.2 deaths per million Americans for which cold was listed as the main underlying cause and an additional 2.4 deaths per million for which cold was listed as a contributing factor (EPA 2025). By contrast there were 1.3 deaths per million for which heat was the main cause and an additional 0.8 per million for which it was a contributor. By this metric cold accounts for approximately double the weather-related deaths as does heat.

Epidemiological methods that consider correlational evidence, not just death certificate reports, indicate the cold/warm ratio might be much higher. A 13-country study of 74 million deaths from 1985 to 2012 estimated that, on average, 7.7 percent of deaths were attributable to sub-optimal temperatures, of which 7.4 percent were attributable to cold and only 0.4 percent were attributable to heat (Gasparini *et al.* 2015). In other words, cold weather killed 18.5 times as many people as did hot weather.

Figure 10.2 shows the distribution of results from Gasparini *et al.* (2015) by country. For the United States the fraction of deaths attributable to temperature was 5.9 percent, of which 5.5 percent was due to cold, thus cold weather killed 14 times as many people as hot weather.




Figure 10.2. Mortality attributable to extreme and moderate cold and heat by country. Source: reproduced from Gasparini *et al.* (2015).

There is strong evidence that people adapt to weather risks. Lee and Dessler (2023) reported that 86 percent of temperature-related deaths across 40 cities in the U.S. were due to cold-related mortality, and that due to adaptation the relative risk of death declined in hot and cold cities alike as seasonal temperatures increased. Allen and Sheridan (2018) found that short, early-season cold events were 2 to 5 times deadlier than hot events, but the mortality risk of both cold and hot extremes drops to nearly zero if the events occur late in the season.

Davis *et al.* (2003) examined heat-related mortality in 28 U.S. cities from the 1960s to the end of the 1990s and found that heat-related mortality declined by three-quarters over the sample period. Bobb *et al.* (2014) examined mortality data for 106 million people in over 100 cities U.S. cities and found a 60 percent decline in average heat-related mortality over the period 1987—2005, from 51 per thousand deaths to 19. They furthermore found that the greatest drop was among seniors over the age of 75. In a study of 42 million

deaths in 211 U.S. cities from 1962 to 2006. Nordio *et al.* (2015) found a more than 90 percent decline in the risk of mortality from excess heat.

In the context of large declines in heat-related mortality, rising temperatures are associated with a net saving of lives since they reduce mortality from cold events. The IPCC AR6 Working Group 2 Chapter 16.2.3.5 (O'Neill *et al.* 2022) acknowledges that heat-related mortality risk is declining over time:

Heat-attributable mortality fractions have declined over time in most countries owing to general improvements in health care systems, increasing prevalence of residential air conditioning, and behavioral changes. These factors, which determine the susceptibility of the population to heat, have predominated over the influence of temperature change.

Yet the IPCC misrepresents the overall situation in its AR6 Synthesis report. Section A.2.5 of that document states: “In all regions increases in extreme heat events have resulted in human mortality and morbidity (*very high confidence*).” But it is silent on the larger decline of deaths during extreme cold events.

The observed decline in U.S. heat-related mortality has been specifically attributed to adaptation. Wang *et al.* (2018) exploited the spatial variability of heatwave-related mortality across 209 U.S. cities from 1962 to 2006. While simple correlation appeared to imply an increase in mortality risk during heatwaves, accounting for adaptation to heatwave intensity caused the effect to fall to near zero and become statistically insignificant. They used the results of their epidemiological model to project heat-related mortality out to 2050 under four RCP warming scenarios (including RCP8.5) with and without adaptive behavior. Assuming people continue to adapt to the heatwave risks in the regions in which they live. Wang *et al.* (2018) project not only no increase in heat-related mortality, but an overall mortality *decrease* for the U.S. as a whole. They conclude that

Ignoring adaptation would result in a substantial overestimate of future mortality related to heat waves... Accounting for adaptation, the overall heat-related mortality by 2050 would not change substantially over time compared to 2006.

10.3.2 Mortality risks and energy costs

A 2016 study of U.S. long term mortality risks associated with temperature variations (Barreca *et al.* 2016) showed that increases in mortality across the U.S. are associated with both cold weather and hot weather. But over time, the introduction of electricity and the adoption of central heating and air conditioning (AC) dramatically reduced both risks, especially those associated with hot weather. Prior to 1960, a day above 90°F (32°C) added 2.2 percent to the average mortality risk rate, but after 1960 the same weather added only 0.3 percent to mortality risk, an 85 percent reduction. Prior to 1960 temperatures below 39°F (4°C) added about 1 percent to mortality risk but after 1960 the same weather only added about half that amount. Adaptation through conventional household improvements dramatically reduced public vulnerability to weather extremes. The entire reduction in hot weather mortality was attributable to widespread adoption of indoor AC, which depended on the availability of reliable and affordable electricity.

The corollary of this finding is that the use of heating and cooling systems depends on energy being affordable. Doremus *et al.* (2022) showed that wealthy and poor households in the U.S. adjust their energy expenditures at similar rates in response to moderate temperature swings, but not in response to extreme temperature swings. When temperatures swing to very cold levels (< 5°C) energy spending in high-income

households rises by 1.2 percent but in low-income households by only 0.5 percent. On very hot days (>30°C) electricity spending in high-income households rises by 0.5 percent but does not change at all in low-income households. The latter result is observed even in subsamples where all households have AC. Cong et al. (2022) report similar findings for a sample of households in Arizona. The implication is that even with widespread adoption of home heating and cooling systems, inability to afford energy leaves low-income households exposed to weather extremes.

References

- Allen, M.J. and S.C. Sheridan (2018) Mortality risks during extreme temperature events (ETEs) using a distributed lag non-linear model. (2018) *International Journal of Biometeorology* 62, 57–67 (2018). <https://doi.org/10.1007/s00484-015-1117-4>
- American Public Power Association. (APPA, 2024, March 7). TVA flood mitigation strategies prevented approximately \$406 million in potential damages. Public Power Current. <https://www.publicpower.org/periodical/article/tva-flood-mitigation-strategies-prevented-approximately-406-million-potential-damages>
- Barreca, Alan, Karen Clay, Olivier Deschenes, Michael Greenstone, and Joseph S Shapiro (2016), Adapting to climate change: The remarkable decline in the US temperature mortality relationship over the twentieth century, *Journal of Political Economy*, 2016, 124 (1), 105–159.
- Battelle Memorial Institute. (2013, December). Final independent external peer review report: Hurricane Isaac (2012) assessment (Prepared for U.S. Army Corps of Engineers, New Orleans District). <https://www.mvn.usace.army.mil/Portals/56/docs/PD/PeerReview/TCN13-003HurrlsaacAssessIEPRFinal.pdf>
- Bobb JF, RD Peng, ML Bell and F Dominici (2014) Heat-related mortality and adaptation to heat in the United States. *Environmental Health Perspectives* 122:811–816; <https://ehp.niehs.nih.gov/doi/full/10.1289/ehp.1307392>
- Cong, S., D. Nock, Y.L. Qiu, Y.L. et al. Unveiling hidden energy poverty using the energy equity gap. *Nature Communications* 13, 2456 (2022). <https://doi.org/10.1038/s41467-022-30146-5>
- Davis, Robert E., Paul C. Knappenberger, Patrick J. Michaels and Wendy M. Novicoff (2003) “Changing Heat-Related Mortality in the United States” *Environmental Health Perspectives* 111(14) November 2003 pp. 1712—1718 <https://ehp.niehs.nih.gov/doi/pdf/10.1289/ehp.6336>
- Deryugina, T., & Hsiang, S. M. (2023). Does the environment still matter? Daily temperature and income in the United States (NBER Working Paper No. 31361). National Bureau of Economic Research. <https://www.nber.org/papers/w31361>
- Doremus, JM, I. Jacqz and S. Johnston (2022), “Sweating the energy bill: Extreme weather, poor households, and the energy spending gap.” *Journal of Environmental Economics and Management*, doi: <https://doi.org/10.1016/j.jeem.2022.102609> .
- Environmental Protection Agency (2025) Climate Change Indicators. <https://www.epa.gov/climate-indicators/view-indicators>
- Federal Emergency Management Agency (FEMA). (2020, February 13). The role of Florida’s building codes in Hurricane Michael recovery. <https://www.fema.gov/case-study/role-floridas-building-codes-2018-hurricane-michael>
- Gasparini, A., Y. Guo, M. Hashizume et al. (2015) Mortality risk attributable to high and low ambient temperature: a multicountry observational study. *The Lancet* 386: 369-385 July 25, 2015 [http://dx.doi.org/10.1016/S0140-6736\(14\)62114-0](http://dx.doi.org/10.1016/S0140-6736(14)62114-0)

- Goklany, I. M. (2011). Deaths from extreme weather events, 1900–2010. Reason Foundation. https://a8d50b36.delivery.rocketcdn.me/wp-content/uploads/2011/09/deaths_from_extreme_weather_1900_2010.pdf
- Lee, Jangho and Andrew Dessler (2023) Future Temperature-Related Deaths in the U.S.: The Impact of Climate Change, Demographics, and Adaptation. *GeoHealth* 7(8) August 2023 <https://doi.org/10.1029/2023GH000799>
- Molina, R and I Rudik (2024) The value of improving hurricane forecasts. *National Bureau of Economic Research Working Paper* 32548 June 2024. <https://www.nber.org/digest/202409/value-improving-hurricane-forecasts>
- National Hurricane Center. (n.d.). Costliest U.S. tropical cyclones tables updated. NOAA National Hurricane Center. Retrieved May 8, 2025, from <https://www.nhc.noaa.gov/outreach/history/>
- National Research Council. (2010). Adapting to the impacts of climate change. The National Academies Press. <https://nap.nationalacademies.org/read/12888/chapter/2>
- Nordio, Francesco, Antonella Zanobetti, Elena Colicino, Itai Kloog and Joel Schwartz (2015). “Changing patterns of the temperature–mortality association by time and location in the US, and implications for climate change” *Environment International* Volume 81, pp 80-86 <https://doi.org/10.1016/j.envint.2015.04.009>.
- O'Neill, B., *et al.* (2022). Key risks across sectors and regions. In AR6 Working Group II *Climate Change 2022: Impacts, Adaptation and Vulnerability* (pp. 2411–2538). Intergovernmental Panel on Climate Change. www.ipcc.ch
- Pielke Jr., R. (2024). Scientific integrity and U.S. 'billion dollar disasters'. *npj Natural Hazards*, 1, Article 12. <https://www.nature.com/articles/s44304-024-00011-0>
- Pielke Jr., R. (2025, May 8). RIP NOAA's billion dollar disasters. The Honest Broker [Substack]. <https://rogerpielkejr.substack.com/p/rip-noaas-billion-dollar-disasters>
- Ritchie, H. (2024). How many people die from extreme temperatures, and how this could change in the future: Part one. Our World in Data. <https://ourworldindata.org/part-one-how-many-people-die-from-extreme-temperatures-and-how-could-this-change-in-the-future>
- Tennessee Valley Authority (TVA). (n.d.). Flood management. Retrieved May 8, 2025, from <https://www.tva.com/environment/managing-the-river/flood-management>
- U.S. Census Bureau. (n.d.). Decennial census by decade. Retrieved May 8, 2025, from <https://www.census.gov/programs-surveys/decennial-census/decade.html>
- Wang, Yan, Francesco Nordio, John Nairn, Antonella Zanobetti and Joel D. Schwartz (2018) “Accounting for adaptation and intensity in projecting heat wave-related mortality” *Environmental Research* 161 pp. 464—471. <https://doi.org/10.1016/j.envres.2017.11.049>
- Zhao, Q., *et al.* (2021). Global, regional, and national burden of mortality associated with non-optimal ambient temperatures from 2000 to 2019: A three-stage modelling study. *The Lancet Planetary Health*, 5(7). [https://doi.org/10.1016/s2542-5196\(21\)00081-4](https://doi.org/10.1016/s2542-5196(21)00081-4)

11 CLIMATE CHANGE, THE ECONOMY, AND THE SOCIAL COST OF CARBON

Chapter summary

Economists have long considered climate a relatively unimportant factor in economic growth, a view echoed by the IPCC itself in AR5. Mainstream climate economics has recognized that CO₂-induced warming might have some negative economic effects but they are too small to justify aggressive abatement policy and that trying to “stop” or cap global warming even at levels well above the Paris target would be worse than doing nothing. An influential study in 2012 suggested that global warming would harm growth in poor countries, but the finding has subsequently been found not to be robust. Studies that take full account of modeling uncertainties either find no evidence of a negative effect on global growth from CO₂ emissions or find poor countries as likely to benefit as rich countries.

Social Cost of Carbon (SCC) estimates are highly uncertain due to unknowns in future economic growth, socioeconomic pathways, discount rates, climate damages, and system responses. The SCC is not intrinsically informative as to the economic or societal impacts of climate change. It provides an index connecting large networks of assumptions about the climate and the economy to a dollar value. Some assumptions yield a high SCC and others yield a low or negative SCC (i.e. a social benefit of emissions). The evidence for or against the underlying assumptions needs to be established independently; the resulting SCC adds no additional information about the validity of those assumptions. Consideration of potential tipping points does not justify major revisions to SCC estimates.

11.1 Climate change and economic growth

11.1.1 Overview

It has long been noted that economies tend to do poorly in very cold and very hot regions, with the optimum somewhere in between (Nordhaus, 2006). This implies that warming will tend to be harmful in hot regions but beneficial in cool ones. Temperature-sensitive economic activity migrates, whenever possible, to where it is best suited, and society adapts to the local climate. Based in part on these observations, in 1992 Thomas Schelling, then President of the American Economic Association, argued that any effects of climate change on U.S. economic activity would be small relative to the many other changes that would happen (Schelling 1992).

... Manufacturing rarely depends on climate, and where temperature and humidity used to make a difference, air conditioning has intervened. When Toyota chooses among Ohio, Alabama, and Southern California for locating an automobile assembly, geographical considerations are important, but not because of climate... Finance is little affected by climate; similarly for health care, or education, or broadcasting. Transportation can be affected, but improvements in all-weather landing and take-off in the last 30 years are greater than any differences that climate makes. If the average effect is a warming, iced waterways and snow removal may decline in importance.

Construction is affected, mainly by cold, and if the average effect is in the direction of warming, construction may benefit slightly.

It is really agriculture that is affected. But even if agricultural productivity declined by a third over the next half century, the per capita GNP we might have achieved by 2050 we would achieve only in 2051. ...

I conclude that in the United States, and probably Japan, Western Europe, and other developed countries, the impact on economic output will be negligible and unlikely to be noticed.

Thirty years later virtually the identical point was made by the IPCC itself in the Fifth Assessment Report (Arent *et al.* 2014, emphasis added)

For most economic sectors, the impact of climate change will be small relative to the impacts of other drivers. Changes in population, age, income, technology, relative prices, lifestyle, regulation, governance, and many other aspects of socioeconomic development will have an impact on the supply and demand of economic goods and services that is large relative to the impact of climate change.

Evidence since AR5 does not change this assessment. Mohaddes *et al.* (2023) found that warm weather shocks have small negative effects on U.S. state-level output but not income, while cold weather shocks negatively affect both and the impacts are about four times larger, implying a shift to warmer conditions would, if anything, yield a net economic benefit for the U.S.

These statements are validated by experience. Since 1900, the average global surface temperature anomaly warmed 1.3°C, about as much as the IPCC predicts will occur in the next century under a moderate emissions scenario. But even as the globe warmed and the population quintupled, humanity prospered as never before. For example, global average lifespan went from thirty-two years to seventy-two years, economic activity per capita grew by a factor of seven, and the death rate from extreme weather events plummeted by a factor of fifty. Climate change damage projections typically refer to reductions in how much life will improve for humanity, they don't state that it will get worse in an absolute sense (O'Neill 2023).

While extreme weather events are costly, in all modern economies they are becoming steadily less and less important (Formetta and Feyen, 2019). Since 1990 weather-related disaster losses have declined as a proportion of global GDP (Pielke Jr , 2018, 2020) as have mortality risks (Formetta and Feyen, 2019). While economic weather-related insurance payouts are rising, this is fully explained by the growth in the size of the economy and the value of the insured asset stock. Past increases in episodes extreme weather have not had a significant effect (positive or negative) on the market value of insurance firms (Hu and McKittrick, 2015). Nor have past extreme weather events had a significant effect on U.S. banks' performance (Blickle *et al.*, 2021); warming has even been shown to be beneficial for the finance and insurance sector (Mohaddes *et al.* 2023). Figure 11.1 below is illustrative. For these reasons, economists have long been reluctant to endorse attempts to “stop” climate change or even aggressively reduce GHG emissions because the costs would not be worth it. As one critic of the economics of climate policy put it (Storm 2017):

Mainstream climate economics takes global warming seriously, but perplexingly concludes that the optimal economic policy is to almost do nothing about it... The contrast is striking. While climate science is sending out loud-and-clear messages that fossil-fuel disinvestment must start now, letting

go of coal and oil and diverting resources into renewable energy technology systems, to keep warming below the 2°C limit (IPCC 2014), mainstream climate economics claims that overly ambitious climate targets will unnecessarily hurt the economy and immediate de-carbonization is too expensive. Most climate economists thus recommend humanity to just wait-and-see.




Figure 11.1 Global weather losses as a fraction of GDP. Source: Pielke Jr. (2023)

The mainstream economics position on the climate policy question is best represented by the findings over the past three decades from Integrated Assessment Models (IAMs) of climate change policy, for which Yale economist William Nordhaus was awarded the 2018 Nobel Memorial Prize in economics. IAMs combine economic, climate, and social data into a unified framework for simulating climate damages and evaluating the optimal policy response (Resources for the Future, 2025). Nordhaus' work has generally supported modest global climate policy, with aggressive measures largely deferred to later in the century. Nordhaus developed the so-called Dynamic Integrated Model of the Climate and Economy model, or DICE, in the early 1990s (Nordhaus, 1993) to study the interaction between climate change, climate policy, and global economic growth over long times. DICE assumes global emission control can be coordinated at no cost and asks what the optimal policy target should be. The climate component of the DICE model is based on simplified climatological modeling. The version of DICE at the time of Nordhaus' Nobel Prize award was described in Nordhaus (2018); it had a climate sensitivity parameter of 3.1°C for CO₂ doubling, consistent with the best estimate in IPCC reports (3.0°C).

The baseline (no policy) scenario incurs only \$0.4T in global abatement costs and leads to \$134.2T (trillion dollars) in global climate damages for a total cost of \$134.6T. The climate model component of DICE projects 4.1°C warming by 2100 relative to preindustrial temperatures. Note that this is a higher estimate of warming than many IPCC climate models.

The Optimal Policy scenario barely deviates from the business-as-usual path. It aims for +3.5°C warming, in other words we modestly scale back fossil fuel use and otherwise just live with almost all the warming. This suggests that most CO₂ emissions are less harmful than the policies that would be necessary to abate them. Trying to prevent warming causes costs quickly to outstrip the benefits. Pursuing a goal of capping warming at 2.5°C creates total costs of \$177.8T, which is \$43.2T worse than doing nothing at all. Nordhaus didn't evaluate trying to reach a Paris-type target of 1.5°C warming but it would be even more costly.

A subsequent edition of DICE includes higher assumed damages from warming which, not surprisingly leads to more aggressive policy recommendations, as explained in the section below on the Social Cost of Carbon.

11.1.2 Empirical analysis of climate change and economic growth

Many other studies have used econometric methods applied to historical data, instead of IAMs, to study the potential impact of climate change on economic growth. Dell *et al.* (2012, herein DJO12) was an influential study that used a multi-country panel of national-level data spanning 1950 to 2005, in which they matched climate and economic data by averaging temperature from the local grid cell level up to the national level using population weights. They found that warming yields an insignificant positive effect on income growth in rich countries but a significant negative effect in poor countries. Moore and Diaz (2015) modified the DICE model to take that finding into account and concluded that it implied a dramatically higher Social Cost of Carbon due to the compounding effects of income loss over time.

A large subsequent literature has debated the robustness of the DJO12 findings. Burke *et al.* (2015) analyzed a global panel with temperatures averaged up to the national level and found a negative effect on growth from warming in rich and poor countries alike when the national average temperature is above 13°C. Zhao *et al.* (2018) used the G-Econ data set of Nordhaus (2006), which breaks economic activity down to the grid cell level, replicated DJO12-type results on the same subset of countries as used in DJO12 but then showed that on the full global sample warming increases growth in rich countries and poor countries alike, though the positive effect in the latter group is confined to where local temperatures are below about 16°C. Greßer *et al.* (2021) developed a regional economic data set for 1,542 sub-national regions around the world between 2005 and 2015 and found temperature had no effect on growth. Yang *et al.* (2023) updated the DJO12 data set and applied an estimator robust to mixed sample frequencies, finding that while temperature shocks exerted a temporary effect on income levels, they did not have a significant lasting effect on growth rates.

Newell *et al.* (2021) noted that there is no underlying theory to guide econometric model specification in this literature. Taking into account the arbitrary choices of which explanatory variables to include they identified over 800 possible model specifications. Using the Burke *et al.* (2015) data they used an estimation method that accounts for model uncertainty and found that the model form preferred by Burke *et al.* (2015), which implied negative effects of warming on growth even in rich countries, is explicitly excluded by the optimal model selection algorithm. Overall they could not detect a temperature effect on GDP or GDP growth, and they estimated the 95 percent confidence interval for the impact on global growth as of 2100 even under the exaggerated RCP8.5 warming scenario spans –86 percent to +388 percent. In other words the net effect is likely positive but too uncertain to distinguish from zero.

Barker (2023) criticized the DJO12 assumption that countries can be grouped into fixed “poor” and “rich” categories based on their incomes many decades ago. He noted that many countries were once poor but became rich over time, and if this is considered the original temperature effects reported by Dell *et al.* became small and insignificant.

Berg *et al.* (2023) argued that countries shouldn’t be grouped into large Rich/Poor categories because they are too heterogeneous. They instead estimated country-specific temperature response coefficients then grouped countries with similar response coefficients into small panels. They separately estimated responses to global and idiosyncratic local temperature shocks to better identify the climate signal in weather data. Overall, they found countries experiencing negative effects of warming on growth outnumbered those experiencing positive effects, but only temporarily: eventually the effects reverse such that about twice as many countries experience a net positive growth effect. They also found global (as opposed to local) temperature changes are much more likely to benefit growth in poor countries than rich ones. In a simulation to 2100, even using the extreme RCP8.5 scenario, they computed the average global GDP loss would be

only 1.9 percent compared to a scenario with no warming. That is, instead of the economy growing 400 percent it would grow 392%. The implication of Nordhaus' earlier analysis is that trying to prevent the warming would lead to far less than 392 percent growth.

To summarize, economists consider climate a relatively unimportant factor in economic growth, a view echoed by the IPCC itself in the Fifth Assessment Report. Mainstream climate economics has recognized that CO₂-induced warming might have some negative economic effects but they are too small to justify aggressive abatement policy, and trying to “stop” or cap global warming even at levels well above the Paris target would be worse than doing nothing. An influential study in 2012 suggested that global warming would harm growth in poor countries but the finding has subsequently been found not to be robust (Tol 2024). Studies that take full account of modeling uncertainties either find no evidence of a negative effect on global growth from CO₂ emissions or find poor countries as likely to benefit from it as rich countries.




Figure 11.2: Decline in U.S. GDP per degree of warming. Source: CEA-OMB (2023)

The expectation that significant global warming would have a small impact on the U.S. economy was acknowledged quietly by the Biden Administration, even as the President was proclaiming a climate emergency. Figure 11.2, from a 2023 CEA-OMB report, shows the expected decrement in U.S. GDP as a function of temperature rise. The colored lines show the results of a dozen peer-reviewed published studies while the solid black line is their average. The figure could be summarized as “a few percent impact for a few degrees of warming”. Given that the economy’s annual growth rate is expected to be 1-2 percent, the impact of a warming globe on the U.S. GDP is indeed negligible.

11.2 Models of the Social Cost of Carbon

The Social Cost of Carbon (SCC) is a tool for quantifying the economic impact of carbon dioxide emissions, helping policymakers weigh the costs and benefits of climate policies. It estimates the damage caused by emitting one additional ton of CO₂, expressed in dollars. More formally, the SCC is the discounted present value of the current and future marginal loss of economic welfare due to an additional ton of CO₂ entering the atmosphere.

11.2.1 *Estimating the SCC*

Although the literature refers to “estimates” of the SCC, it is not estimated in the way other economic statistics are estimated. For instance, data on market transactions including prices and quantities can be used to estimate the current inflation rate or the growth rate of per capita real Gross Domestic Product, and there are well-understood uncertainties associated with these quantities. But there are no market data available to measure many, if not most, of the marginal damages or benefits believed to be associated with CO₂ emissions, so these need to be imputed using economic models.

For example, an influential component of some SCC calculations is the perceived social cost associated with a changed risk of future mortality due to extreme weather. There is no market in which people can directly attach a price to that risk. At best economists can try to infer such values by looking at transactions in related markets such as real estate or insurance, but isolating the component of price changes attributable to atmospheric CO₂ levels is very difficult.

Economists use IAMs to compute the SCC. Two of the best-known are the Climate Framework for Uncertainty, Negotiation and Distribution (“FUND”, Tol 1997) and Nordhaus’ DICE. EPA (2023) introduced new ones for its recent work. IAMs embed a “damage function” or set of functions relating ambient temperature to local economic conditions. The assumptions embedded in the damage function will largely determine the resulting SCC. IAMs also assume a long-term discount rate or, as in DICE, compute the optimal internal discount rate as part of the solution.

One approach to developing a damage function is to begin with estimates of the costs (or benefits) of warming in specific sectors in countries around the world and aggregate up to a global amount. This was the approach used in FUND. An alternative approach is to develop a simple equation that penalizes global income according to a simple quadratic function of the average global temperature. This approach was used in DICE. In the case of the FUND model many hundreds of parameters had to be selected, whereas in DICE only three were needed and were originally chosen to assign a predetermined penalty (1.2%) to global output as of 3°C warming, with a quadratic term implying that damages grow with the square of the global average temperature anomaly. Barrage and Nordhaus (2024) recently changed the parameters to increase the penalty at 3°C to 1.6%, and added a discrete additional 1.0 percent GDP penalty at 3°C warming to account for “tipping points” (discrete large-scale environmental changes triggered by crossing a warming threshold) and a “judgmental adjustment” of 0.5 percent for excluded impacts at 3°C warming. Not surprisingly the newer version of DICE generates much higher SCC estimates than before.

The concepts of estimation and uncertainty do not readily apply to SCC calculations. No amount of data collection can change the fact that many components of the SCC are unknown and rely on judgment and opinion based on knowledge of the underlying literature on the physical effects of climate change. SCC calculations are thus best thought of as “if-then” statements: *if* the following assumptions hold, *then* the SCC is \$X per tonne. The list of ‘if’ statements includes the premise that the world’s climate and economy work according to the representation in the IAM. One way this might fail to hold is in the timing of warming. Every IAM assumes a value of the Equilibrium Climate Sensitivity (ECS), which controls the amount of warming that results from CO₂ emissions and can be freely varied for the purpose of generating a distribution of SCC values associated with uncertainties over ECS. But as Roe and Baumann (2013) pointed out, time-to-equilibrium increases with the square of ECS, so an upward adjustment of the ECS parameter without an appropriate slowing down of the adjustment process can yield distorted present value damage estimates. In particular, the upper tail of warming associated with some commonly-used ECS distributions is physically impossible for even a thousand years into the future (Roe and Baumann 2013) yet in an IAM would be realized within a couple of centuries. Failure to align ECS with time-to-equilibrium will lead to an overestimate of the SCC value.

11.2.2 Variations in the SCC

Every level of the IAM calculation includes assumptions, some more influential than others. Key assumptions include the following.

- The discount rate: Climate damages accrue over a long time horizon and costs a century or two in the future need to be discounted to the present. The higher the discount rate the smaller the present value of future damages and vice versa. The discount rate represents the opportunity cost of spending money today rather than investing it and then having more to spend tomorrow. Some economists have argued for the use of very low discount rates in SCC calculations, resulting in policy recommendations that favor relatively large immediate investments in CO₂ emission reductions. The downside is that other investments could potentially earn a larger rate of return for society.
- Equilibrium Climate Sensitivity: IAMs have customarily employed a value of 3.0°C or 3.1°C following the IPCC's guidance. The most recent data-driven ECS values tend to be lower than this (see Chapter 4). Dayaratna *et al.* (2017, 2020, 2023) have shown that use of lower empirically-derived ECS values dramatically lower the resulting SCC estimate, even when low discount rates are used.
- Damage function coefficients: IAMs assume CO₂ and warming cause net harms that increase exponentially with temperature. More recently, IAMs have also incorporated effects from assumed potential climate tipping points. The FUND model took limited but explicit account of CO₂ fertilization effects in agriculture. Since the coefficients were selected prior the publication of the current evidence of global greening and the magnitude of benefits to crops from elevated CO₂ (see Chapters 2 and 9) the growth effects are likely understated. The DICE model (and others) did not explicitly include any CO₂ fertilization benefit, except to the extent it was taken into account in the literature consulted when picking the damage function coefficients. The damage function in FUND contains a region in which low warming yields net benefits in many regions, a finding which is supported by econometric models of warming and growth (Berg *et al.* 2023) and econometric simulations of agricultural changes (McKittrick 2025). The DICE damage function, by construction, rules out net benefits at any warming level.
- Emission scenarios: IAMs generate SCC estimates that increase as the pre-existing concentration of CO₂ increases. Consequently the value of damages later in the century will be higher, depending on the assumed baseline emissions over the coming decades.

Some IAMs (such as DICE) include the cost to the economy of reducing CO₂ emissions in order to identify the SCC along an optimal growth path. If CO₂ emission reductions are assumed to be inexpensive, then the model will conclude that the optimal policy should aim for deeper emission cuts and vice versa.

It is informative to ascertain whether SCC results are invariant to changes in some assumptions. But when different assumptions result in higher or lower SCC values, the change in the SCC value does not provide *prima facie* evidence about the validity of the assumptions. For example, in 2023 the U.S. Environmental Protection Agency raised its preferred SCC value about 5-fold over the estimates it had issued ten years earlier. This is not because new data had been collected or better mathematical methods had been invented, but because new assumptions had been used, and the validity of those assumptions was a separate question. Tabulations on EPA (2023) p. 81 show that if assumptions similar to earlier analyses had been applied, the results would not have materially differed from before. One new assumption was that global agricultural damages were far higher than previously believed, based on an analysis in Moore *et al.* (2017). But as discussed in Chapter 9, McKittrick (2025) showed that Moore *et al.* (2017) had used a database in which half the CO₂ change observations were missing. When as many of those observations as possible were recovered from underlying sources and the analysis was rerun, the projected crop yield losses disappeared and turned into gains at all warming levels. Hence the portion of the EPA's SCC revision attributable to agricultural yield losses was unwarranted.

11.2.3 Evidence for low SCC

Chapter 2 reviewed evidence on climate change and greening, and Chapter 9 looked at climate change and U.S. agriculture. Evidence shows that CO₂ fertilization has a stronger beneficial effect on agriculture than was known at the time that IAMs like DICE and FUND were parameterized. Haverd *et al.* (2020) found the observed CO₂ fertilization rate has been almost double what had been predicted by crop models. Dayaratna *et al.* (2023) used the updated empirical ECS distribution estimate of Lewis (2022), which assimilates modern instrumental and paleoclimatic temperature records, and allowed for a 30 percent gain in the CO₂ fertilization benefit in the FUND model, and found that, even at a low discount rate of 2.5 percent, the median SCC as of 2050 is only \$18.67, with a 24 percent probability of the true value being negative. At a five percent discount rate the median SCC value is negative until the mid-2040s and at 2050 was only \$0.37 with a 49 percent probability of being negative. Thus, under reasonable assumptions a mainstream IAM using updated scientific inputs yields evidence consistent with the SCC not being significantly greater than zero.

It should also be noted that the SCC is focused on the social costs of CO₂ emissions from fossil fuel use. It is not intended to measure the private marginal benefits to consumers and society from the availability of fossil fuels. Public willingness to pay for fuels of all types indicates the value to society of reliable, abundant fossil energy. Tol (2017) estimates that the private benefit of carbon is large relative to the social cost. This can be illustrated by noting that the price of a gallon of gas indicates the marginal value to the consumer of the fuel. Suppose we assume a relatively high Social Cost of Carbon of, say, \$75 per tonne. Deflated by a MCPF⁴ value of 1.5 that would result in a carbon tax of \$50 per tonne, which equates to about 44 cents per gallon of gas (Lavelle, 2019). A pre-tax price of \$3.00 per gallon would imply the marginal social benefit of the fuel is nearly seven times the marginal social cost.

11.2.4 Tipping points

SCC calculations typically consider gradual impacts of a warming climate, such as slowly-melting glaciers and increasing average temperature. A driver of potentially high values of the Social Cost of Carbon (SCC) is the introduction into models of discrete catastrophic outcomes associated with abrupt changes (Dietz *et al.*, 2021). They are often referred to as “tipping points.”

The term “tipping point” mingles two different physical concepts that pose different research challenges. Many physical systems are inherently stable unless acted upon with sufficient external energy. For example, an ice cap might remain intact over a wide range of temperatures but once the temperature crosses the 0°C threshold it melts. Such discontinuities are ubiquitous in nature and require an external force. Whether the force needs to be large relative to the size of the system depends on the underlying stability of the system.

A different type of tipping point is called a *bifurcation* and arises from the study of the internal dynamics of nonlinear systems (Crawford, 1991). Many systems have been observed to have more than one equilibrium point and can move between them with minimal or no external influence. For example, a weather system might have two different equilibrium states: one calm and one with a tornado. A transition from one to another can happen either with no external force or with a minuscule change, such as a flap of the proverbial butterfly’s wings (Shen *et al.* 2014). The term “tipping point” is sometimes used to mean a bifurcation of this type and implies instability inherent in the system itself, which is not necessarily dependent on outside forces. It depends, instead, on parameters of the system taking values that support the emergence of bifurcations (Crawford, 1991).

The two different concepts imply different research questions. Regarding the first we are interested in whether components of the climate system are susceptible to abrupt discontinuities in response to

⁴ Marginal Cost of Public Funds: the optimal carbon tax rate is the SCC divided by the MCPF (Sandmo 1977).

sufficiently-large anthropogenic forcing. Regarding the second we are interested in whether the Earth’s climate system has inherent bifurcations that imply the possibility of abrupt transitions with or without external forcing.

Models have been developed that imply the second type is a possibility. Kypke *et al.* (2020) presented a simple climate model in which the GHG concentration is one of the parameters that controls the emergence of bifurcations of the Arctic climate to one with both cold and warm equilibria. If sufficiently high GHG forcing combined with a sufficiently high rate of ocean heat transport are imposed a bifurcation becomes possible. More generally GCMs have been observed to contain bifurcations and multiple equilibria (Brunetti and Ragon, 2023).

The possible existence of bifurcations in the Earth’s climate system implies abrupt transitions are possible, not just in response to large forcing but also to small perturbations. This places tipping points into the category of low-likelihood and potentially catastrophic events, such as large meteor strikes. A key question to ask is whether those kinds of tipping points can be predicted. Current research has not resolved that question (Dakos *et al.*, 2024) and indeed might not be able to since one implication of the “butterfly effect” is the existence of predictability boundaries of nonlinear systems (Palmer *et al.*, 2014). It is therefore not obvious how to incorporate such possibilities into SCC calculations. Small variations in assumptions will lead to arbitrarily large variations in the resulting SCC with no grounds for choosing among them. If such tipping points are possible the most appropriate stance for economic policy is to maximize resilience to any form of external catastrophe since it is unlikely we could predict it or prevent it from happening.

AR6 (WGI, Chapter 1) focuses mainly on the first type of tipping point, namely an abrupt change in response to external forcing. This is also the meaning associated with popular usage of the “tipping point” concept in discussions of climate change (see <https://report-2023.global-tipping-points.org/what-is-a-tipping-point/> for example). As summarized by AR6, there is evidence of abrupt change in the paleoclimate record, and some of these events have been interpreted as tipping points. Some projections with Earth System Models for example have produced tipping points such as Amazon forest dieback in response to specified values of CO₂ concentration or temperature increases.

The alarm surrounding climate tipping points is reflected by *The Global Tipping Points Report* that was launched at the COP28 on 6 December 2023 (Lenton *et al.*, 2023). It identifies more than 25 parts of the climate system said to constitute tipping points. What gets classified as a climate “tipping point” is a moving target. The most common examples in the literature and assessment reports include: Greenland ice sheet disintegration, West Antarctic ice sheet disintegration, summertime disappearance of Arctic sea ice, Amazon rainforest dieback, coral reef dieoff, thawing of permafrost and methane hydrates, Atlantic Meridional Overturning Circulation collapse, boreal forest shift, West African monsoon shift, and Indian Monsoon shift.

All such tipping points require a certain amount of system instability in order to exhibit an abrupt transition in response to warming. For this reason there seems to be very little to discriminate between a tipping event and natural climate variability. Natural climate variability has in the past produced shifts in the West African and Indian monsoons, Amazon forest and coral reef dieback, and disintegration of parts of the Greenland and West Antarctic ice sheets. These impacts can reverse on the decadal and century timescales associated with natural climate variability and ecosystem responses.

Some abrupt changes are potentially more consequential, including collapse of the West Antarctic ice sheet and collapse of the Atlantic Meridional Overturning Circulation (AMOC). AR6 WG1 Summary for Policy Makers states:

The Atlantic Meridional Overturning Circulation is very likely to weaken over the 21st century for all emission scenarios. While there is *high confidence* in the 21st century decline, there is only *low*

confidence in the magnitude of the trend. There is *medium confidence* that there will not be an abrupt collapse before 2100. (C.3.4)

There is limited evidence for low-likelihood, high-impact outcomes (resulting from ice-sheet instability processes characterized by deep uncertainty and in some cases involving tipping points) that would strongly increase ice loss from the Antarctic Ice Sheet for centuries under high GHG emissions scenarios (e.g., SSP5-8.5). (B.5.2)

For the purpose of SCC calculations, the research question implied by this type of tipping point is whether such events have been observed in the past in climate conditions similar to what we currently experience or will in the near future. AR6 finds little evidence for impending collapse of the Atlantic Meridional Overturning Circulation or the West Antarctic ice sheet. It finds there is no tipping point associated with Arctic Sea ice (AR6 Technical Summary p. 76)

Dietz *et al.* incorporated several potential tipping points (abrupt changes) into an SCC model and found they added about 25 percent to the estimate, mainly associated with thawing permafrost and release of methane hydrates. However, the IPCC considers this scenario *very unlikely* (AR6 Technical Summary p. 107).

In summary, there might be unknown bifurcation tipping points that are associated with natural climate processes, but this possibility does not translate into specific guidance on the SCC. There are potential abrupt change points in the climate system in response to warming, although the IPCC assigns low probabilities to most, including the largest ones. When these have been considered, the result is only a modest increase in the SCC value in the 21st century.

11.2.5 Are there alternatives?

It is increasingly being argued that the SCC is too variable to be useful for policymakers. Cambridge Econometrics (Thoung, 2017) stated it's "time to kill it" due to uncertainties. The UK and EU no longer use SCC for policy appraisal, opting for "target-consistent" carbon pricing (UK Department for Energy Security and Net Zero 2022, Dunne 2017). However, the uncertainty of SCC estimates doesn't mean that other regulatory instruments are inherently better or more efficient. Many emissions regulations (such as electric vehicle mandates, renewable energy mandates, energy efficiency regulations and bans on certain types of home appliances) cost far more per tonne of abatement than any mainstream SCC estimate, which is sufficient to establish that they fail a cost-benefit test.

References

- AR6: *Intergovernmental Panel on Climate Change Sixth Assessment Report* (2021) Working Group I Contribution. www.ipcc.ch.
- Arent, D.J., R.S.J. Tol, E. Faust, J.P. Hella, S. Kumar, K.M. Strzepek, F.L. Tóth, and D. Yan (2014) Key economic sectors and services. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 659-708. https://www.ipcc.ch/site/assets/uploads/2018/02/WGIIAR5-Chap10_FINAL.pdf
- Barker, D. (2023) Temperature Shocks and Economic Growth: Comment on Dell, Jones, and Olken. *Econ Journal Watch* 20(2) 234-253 <https://econjwatch.org/articles/temperature-shocks-and-economic-growth-comment-on-dell-jones-and-olken>

- Barrage, Lint and William Nordhaus (2024) Policies, projections, and the social cost of carbon: Results from the DICE-2023 model. *Proceedings of the National Academy of Sciences* 121 (13) e2312030121 <https://doi.org/10.1073/pnas.2312030121>
- Berg, Kimberly, Chadwick C Curtis and Nelson Mark. (2021) GDP and temperature: Evidence on cross-country response heterogeneity. NBER Working Paper 31327 June 2023 <http://www.nber.org/papers/w31327>
- Blickle, Kristian; Sarah N. Hamerling, and Donald P. Morgan (2021) : How bad are weather disasters for banks?, Staff Reports, No. 990, Federal Reserve Bank of New York, New York, NY <https://www.econstor.eu/bitstream/10419/247913/1/sr990.pdf>
- Brunetti, Maura and Charline Ragon (2023) Attractors and bifurcation diagrams in complex climate models. *Physical Review E* 107 054214 <https://journals.aps.org/pre/abstract/10.1103/PhysRevE.107.054214>
- Burke, Marshall, Solomon M Hsiang, and Edward Miguel (2015). Global non-linear effect of temperature on economic production. *Nature*, 527(7577):235–239.
- Council of Economic Advisors-Office of Management and Budget (CEA-OMB 2023) Methodologies and considerations for integrating the physical and transition risks of climate change into macroeconomic forecasting for the president’s budget. March 13, 2023. <https://bidenwhitehouse.archives.gov/wp-content/uploads/2023/03/CEA-OMB-White-Paper.pdf>
- Crawford, John D (1991) Introduction to Bifurcation Theory. *Reviews of Modern Physics* 63(4) October 1991 991—1037.
- Dakos, Vasilis, Chris Boulton, Joshua Buxton *et al.* (2024) Tipping point detection and early warnings in climate, ecological, and human systems. *Earth System Dynamics* 15 1117—1135 <https://doi.org/10.5194/esd-15-1117-2024>
- Dayaratna, Kevin and Ross McKittrick (2023) "Reply to comment on “climate sensitivity, agricultural productivity and the social cost of carbon in fund” by Philip Meyer" *Environmental Economics and Policy Studies* 2023 link.springer.com/article/10.1007/s10018-023-00364-2
- Dayaratna, Kevin, Ross McKittrick and David Kreutzer (2017) Empirically-Constrained Climate Sensitivity and the Social Cost of Carbon. *Climate Change Economics* April 2017 DOI: <http://dx.doi.org/10.1142/S2010007817500063>
- Dayaratna, Kevin, Ross McKittrick and Patrick J. Michaels (2020) Climate Sensitivity, Agricultural Productivity and the Social Cost of Carbon in FUND. *Environmental Economics and Policy Studies* <https://doi.org/10.1007/s10018-020-00263-w>
- Dell, Melissa, Benjamin F. Jones, and Benjamin A. Olken (2012). Temperature shocks and economic growth: Evidence from the last half century. *American economic journal: Macroeconomics*, 4(3):66–95, 2012. ISSN 1945-7707.
- Dietz, S., J. Rising, T. Stoerk, G. Wagner (2021) Economic impact of tipping points in the climate system. *Proceedings of the National Academy of Science* 2021 Vol. 118 No. 34 e2103081118
- Dunne, Daisy (2017), “Q&A: The Social Cost of Carbon,” Carbon Brief, February 14, 2017, <https://www.carbonbrief.org/qa-social-cost-carbon/>
- Formetta, Giuseppe and Luc Feyen (2019) Empirical evidence of declining global vulnerability to climate-related hazards. *Global Environmental Change* 57 101920 <https://doi.org/10.1016/j.gloenvcha.2019.05.004>
- Greßer, Christina, Daniel Meierrieks, and David Stadelmann (2021) The link between regional temperature and regional incomes: econometric evidence with sub-national data. *Economic Policy*, 36(107):523–550, 2021.
- Haverd, V., Raupach, M., Briggs, P., *et al.* (2020). Higher than expected CO₂ fertilization inferred from leaf to global observations. *Global Change Biology*, 26, 2390–2402.
- Hu, Bin and Ross McKittrick (2015) Climatic Variations and the Market Value of Insurance Firms. *Journal of Insurance Issues* Vol. 39, No. 1, pp. 92-111 <https://www.jstor.org/stable/43741103>
- Kypke, K. L., Langford, W. F., and Willms, A. R.: Anthropocene climate bifurcation, *Nonlinear Processes in Geophysics*, 27, 391–409, <https://doi.org/10.5194/npg-27-391-2020>, 2020.

- Lavelle, Marianne (2019) Carbon Tax Plans: How They Compare and Why Oil Giants Support One of Them. *Inside Climate News* March 7, 2019 <https://insideclimatenews.org/news/07032019/carbon-tax-proposals-compare-baker-shultz-exxon-conocophillips-ccl-congress/>
- Lenton, T.M., D.I. Armstrong McKay, *et al.* (2023) *The Global Tipping Points Report 2023*. University of Exeter, Exeter, UK. <https://report-2023.global-tipping-points.org/>
- Lewis, N., (2013) An objective Bayesian, improved approach for applying optimal fingerprint techniques to estimate climate sensitivity. *Journal of Climate*, 26, 7414-7429. <https://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-12-00473.1>
- McKittrick, Ross R. (2025) Extended Crop Yield Meta-analysis Data do not Support Upward SCC Revision. *Scientific Reports* 15 article 5575 <https://www.nature.com/articles/s41598-025-90254-2>
- Mohaddes, Kamiar, Ryan Ng., Hashem Psaran, Mehdi Raissi and Jui-Chung Yang (2023) Climate change and economic activity: evidence from US states. *Oxford Open Economics* Vol 2 odac010 <https://doi.org/10.1093/ooec/odac010>
- Moore, Frances C. and Delavane B. Diaz (2015) Temperature impacts on economic growth warrant stringent mitigation policy. *Nature Climate Change*, 5(2):127–131, 2015. ISSN 1758-678X.
- Moore FC, Baldos U, Hertel T, Diaz D (2017) New science of climate change impacts on agriculture implies higher social cost of carbon. *Nature Communications* 8:1607. <https://doi.org/10.1038/s41467-017-01792-x>
- Newell, Richard G., Brian C. Prest, and Steven E. Sexton (2021). The GDP-temperature relationship: Implications for climate change damages. *Journal of Environmental Economics and Management*, 108 <https://doi.org/10.1016/j.jeem.2021.102445>
- Nordhaus, William (2006). Geography and macroeconomics: New data and new findings. *Proceedings of the National Academy of Sciences*, 103(10):3510–3517, 2006. ISSN 0027-8424.
- Nordhaus, William. (2018) Projections and Uncertainties About Climate Change in an Era of Minimal Climate Policies. *American Economic Journal: Economic Policy* August 2018, 10(3): 333-360. <https://pubs.aeaweb.org/doi/pdfplus/10.1257/pol.20170046>
- Nordhaus, W. (1993) Optimal Greenhouse-Gas Reductions and Tax Policy in the "DICE" Model *American Economic Review* Vol. 83, No. 2, (Papers and Proceedings) pp.313-317.
- O'Neill, Brian C. (2023) Envisioning a future with climate change. *Nature Climate Change* 13, pages 874–876 <https://www.nature.com/articles/s41558-023-01784-4>
- Palmer, TN, A Döring and G Seregin (2014) The real butterfly effect. *Nonlinearity* 27(9) 10.1088/0951-7715/27/9/R123 <https://iopscience.iop.org/article/10.1088/0951-7715/27/9/R123/meta>
- Pielke Jr., Roger (2018): Tracking progress on the economic costs of disasters under the indicators of the sustainable development goals, *Environmental Hazards*, DOI: 10.1080/17477891.2018.1540343
- Pielke Jr., Roger (2020) Economic ‘normalisation’ of disaster losses 1998–2020: a literature review and assessment,” *Environmental Hazards*, 2020, pp. 1-19
- Pielke Jr., Roger (2023) Global disaster losses: 1990-2023 Substack essay <https://rogerpielkejr.substack.com/p/global-disaster-losses1990-2023>
- Resources for the Future (2025) “Social Cost of Carbon Explorer,” Resources for the Future website, accessed April 22, 2025, <https://www.rff.org/publications/data-tools/scc-explorer/>.
- Roe, Gerard H. and Yoram Bauman (2013) Climate Sensitivity: Should the climate tail wag the policy dog? *Climatic Change* (2013) 117:647–662 DOI 10.1007/s10584-012-0582-6
- Sandmo, Agnar (1975) “Optimal Taxation in the Presence of Externalities.” *Swedish Journal of Economics* 77 (1), 86–98.
- Schelling, Thomas C (1992) Some economics of global warming. *The American Economic Review*, 82 (1):1–14.
- Shen, Bo-Wen, Roger A Pielke Sr and Xubin Zeng (2023) The 50th Anniversary of the Metaphorical Butterfly Effect since Lorenz (1972): Multistability, Multiscale Predictability, and Sensitivity in Numerical Models. *Atmosphere* 2023 14(8) <https://doi.org/10.3390/atmos14081279>
- Storm, S (2017) “How the Invisible Hand is Supposed to Adjust the Natural Thermostat: A Guide for the Perplexed.” *Science and Engineering Ethics* vol 23: 1307—1331.

- Thoung, Chris (2017) “Is It Time to Kill Off the Social Cost of Carbon?” Cambridge Econometrics Blog, September 14, 2017, <https://www.camecon.com/blog/time-kill-social-cost-carbon/>.
- Tol, Richard S.J. (1997) On the optimal control of carbon dioxide emissions: an application of FUND *Environmental Modeling and Assessment* 2: 151—163.
- Tol, Richard S.J. (2017). "The Private Benefit of Carbon and its Social Cost," Working Paper Series 0717, Department of Economics, University of Sussex Business School. <https://ideas.repec.org/p/sus/susewp/0717.html>
- Tol, Richard S.J. (2024) A meta-analysis of the total economic impact of climate change. *Energy Policy* 185 113922 <https://doi.org/10.1016/j.enpol.2023.113922>
- UK Department for Energy Security and Net Zero (2022). “Carbon Valuation.” GOV.UK, last updated April 4, 2022. <https://www.gov.uk/government/collections/carbon-valuation--2>.
- US Environmental Protection Agency (2023) Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, “Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review”, EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances, November 2023, https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf accessed February 28, 2024.
- Yang, Yimin, Fei Jia, and Haoran Li. Estimation of panel data models with mixed sampling frequencies (2023). *Oxford Bulletin of Economics and Statistics*, 85(3):514–544.
- Zhao, Xiaobing, Mason Gerety, and Nicolai V. Kuminoff (2018) Revisiting the temperature-economic growth relationship using global subnational data. *Journal of environmental management*, 223:537–544, 2018.

12 GLOBAL CLIMATE IMPACTS OF U.S. EMISSIONS POLICIES

Chapter Summary

U.S. policy actions are expected to have undetectably small direct impacts on the global climate and any effects will emerge only with long delays.

12.1 The scale problem

The emissions rates and atmospheric concentrations of criteria air contaminants are closely connected because their lifetimes are short and their concentrations are small; when local emissions are reduced the local pollution concentration drops rapidly, usually within a few days. But the global average CO₂ concentration behaves very differently, since emissions mix globally and the global carbon cycle is vast and slow. Any change in local CO₂ emissions today will have only a very small global effect, and only with a long delay.

Following the emission of a pulse (release) of CO₂ into the atmosphere only about 40± 15 percent of the extra CO₂ will have been sequestered after twenty years. That fraction rises to 75± 10 percent after a thousand years, and the remainder will be gradually removed over the ensuing tens of thousands of years (Ciais *et al.*, 2013, pp. 472-473). Consequently, any reduction in U.S. emissions would only modestly slow, but not prevent, the rise of global CO₂ concentration. And even if global emissions were to stop tomorrow, it would take decades or centuries to see a meaningful reduction in the global CO₂ concentration and hence human influences on the climate.

Reducing the atmospheric stock of CO₂ would require emissions to fall below the natural sequestration rate, assuming the entire increase is anthropogenic. Since that rate has been averaging about 50 percent of emissions in recent decades, a reduction of global emissions by 50 percent would (at least temporarily) halt the rise in atmospheric CO₂. The 1997 Kyoto Protocol proposed to cap industrial nations' CO₂ emissions at a modest five percent below 1990 levels by the year 2012. Even though this policy was too difficult for most nations to implement, full compliance would not have substantially reduced atmospheric CO₂ levels. It would only have slightly slowed CO₂ growth, reaching the projected year 2100 level in 2105 instead (Wigley, 1998). Lomborg (2016) estimated that full compliance with the initial commitments in the Paris Accord would not stop warming, it would only prevent about 0.1C warming and delay hitting the baseline year 2100 temperature levels by about a decade.

Thus, in contrast with conventional air pollution control, even drastic local actions will have negligible local effects, and only with a long delay. The practice of referring to unilateral U.S. reductions as “combatting climate change” or “taking action on climate” on the assumption we can stop climate change therefore reflects a profound misunderstanding of the scale of the issue.

12.2 Case study: U.S. motor vehicle emissions

The scale problem can be illustrated with reference to U.S. motor vehicles. The EPA's 2009 Endangerment Finding focused on CO₂ emissions from cars and light-duty trucks in the U.S. because Section 202(a) of Clean Air Act mandates the EPA to set emissions standards for motor vehicles if pollutants are found to endanger public health or welfare. The 2009 Endangerment Finding therefore obligated the EPA to regulate emissions from new motor vehicles, ostensibly to reduce or eliminate climate-related harms to the U.S. public.

Two questions that naturally arise are: (1) How large a reduction in CO₂ emissions would result from such regulation? and (2) What would be the climate impact of such regulation?

The first question can be addressed by comparing U.S. vehicle-based CO₂ emissions to the global total. The second question can be addressed by using the fact that the reduction in global warming would be, according to the models relied upon by the EPA, proportional to the reduction in global emissions, keeping in mind that the change in the CO₂ content of the atmosphere in any given year is the result of total global CO₂ emissions, not just U.S. emissions.

In 2022, the emissions from U.S. cars and light duty trucks totaled 1.05 billion metric tons of carbon dioxide (GtCO₂, EPA 2024). Meanwhile global CO₂ emissions from energy use totaled 34.6 GtCO₂ (Energy Institute 2024). Hence U.S. cars and light trucks account for only 3.0 percent of global energy-related CO₂ emissions. To a first approximation we can say that even eliminating *all* U.S. vehicle-based emissions would retard the accumulation of CO₂ in the atmosphere by a year or two over a century.

It would also reduce the overall warming trend by at most about 3 percent. For the period 1979-2023, which has the most extensive global coverage of a variety of weather data types, warming trends are determined to a precision of about ± 15 percent, so the impact of reducing the rate of global warming by eliminating U.S. vehicle CO₂ emissions would be far below the limits of measurability. Given that global-average temperature is the most direct climate change metric, impacts on any secondary climate metrics (e.g. severe weather, floods, drought, etc.) from reducing U.S. vehicle CO₂ emissions would be even less measurable.

Consequently, in contrast to the case of local air contaminants like particulates and ozone, even the most aggressive regulatory actions on GHG emissions from U.S. vehicles cannot be expected to remediate alleged climate dangers to the U.S. public on any measurable scale.

12.3 Concluding thoughts

This report supports a more nuanced and evidence-based approach for informing climate policy that explicitly acknowledges uncertainties. The risks and benefits of a climate changing under both natural and human influences must be weighed against the costs, efficacy, and collateral impacts of any “climate action”, considering the nation’s need for reliable and affordable energy with minimal local pollution. Beyond continuing precise, un-interrupted observations of the global climate system, it will be important to make realistic assumptions about future emissions, re-evaluate climate models to address biases and uncertainties, and clearly acknowledge the limitations of extreme event attribution studies. An approach that acknowledges both the potential risks and benefits of CO₂, rather than relying on flawed models and extreme scenarios, is essential for informed and effective decision-making.

References

- Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, *et al.* (2013): Carbon and Other Biogeochemical Cycles. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, *et al.* (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA
- EPA (2024) *Fast Facts: U.S. Transportation Sector Greenhouse Gas Emissions 1990-2022*. Available online at <https://nepis.epa.gov/Exe/ZyPDF.cgi?Dockey=P101AKR0.pdf>
- Energy Institute (2024) *Statistical Review of World Energy*. Available online at <https://www.energyinst.org/statistical-review>
- Lomborg, Bjorn (2016) “Impact of Current Climate Proposals” *Global Policy* 7(1) 109—118. Available at <http://onlinelibrary.wiley.com/doi/10.1111/1758-5899.12295/full>.

Wigley, T.M.L. (1998). “The Kyoto Protocol: CO₂, CH₄ and climate implications.” *Geophysical Research Letters* 25(13), 2285-2288

GLOSSARY

2XCO2: Doubling of the atmospheric carbon dioxide (CO₂), a commonly used baseline for addressing climate change. In 2025, atmospheric CO₂ concentrations were approximately 50 percent of the way to 2XCO₂ compared to 1800.

ABATEMENT: In the context of environmental policy, the reduction of emissions.

ACE (Accumulated Cyclone Energy): A statistical measure of a tropical cyclone's energy accumulated over its lifetime, calculated from a summation of the square of maximum sustained wind speeds.

ACIDIFICATION: A commonly used term for the reduction in ocean pH from more alkaline to less alkaline values.

ACRIM (Active Cavity Radiometer Irradiance Monitor) GAP: the gap in total solar irradiance data between 1989 and 1991 caused by a delay in the launch of a satellite-borne monitor in time to overlap and intercalibrate with the prior system.

AEROSOLS: Tiny solid or liquid particles suspended in the air.

ALKALINE: Having a chemical pH greater than 7.0. The “opposite” of acidic, which means a pH less than 7.0.

AMBIENT: Relating to the immediate surroundings of something.

AMO: Atlantic Multidecadal Oscillation, a 60-year cycle of sea surface temperatures in the North Atlantic.

ANTHROPOGENIC: Created by humans.

AR (Assessment Reports): Periodic assessments published by the Intergovernmental Panel on Climate Change evaluating current knowledge of the climate system, global change, and associated issues. An AR now consists of reports from three working groups, the first (WG1) concerned with physical science and the others concerned with societal impacts and mitigation strategies. The most recent was AR6.

ATTRIBUTION: Assertion of a causal relation, chiefly from anthropogenic GHG emissions to observed climate patterns.

AUTOCORRELATION: The phenomenon in which the current value of a random time series variable is correlated with its value in a previous time period.

BIOMASS: The total quantity or weight of organic matter in a given area or volume.

CARBON: The 6th element in the periodic table. Popularly used as a shorthand term for carbon dioxide.

CMIP (Coupled Model Intercomparison Project): The Intergovernmental Panel on Climate Change's organized effort to compare many different global ocean-atmosphere climate models when they are all forced with the same emissions scenario(s). The two most recent were the CMIP5 and CMIP6 for, respectively, AR5 and AR6.

CLIMATE CHANGE: A change in climate due to human influences, as opposed to climate variability which is natural in origin.

CONUS: The contiguous 48 U.S. states.

DICE (Dynamic Integrated model of the Climate and Economy): An Integrated Assessment Model developed by William Nordhaus in 1993.

DISCOUNT RATE: The interest rate used to convert future costs and benefits into their present value. Varies inversely with present value so a high discount rate reduces the present value of future costs and benefits.

DRY ICE: Frozen carbon dioxide.

ECONOMETRICS: The branch of economics concerned with the use of statistical methods (chiefly multiple regression) to analyze economic systems.

ECS (Equilibrium Climate Sensitivity): The total amount of eventual global-average surface warming in response to a hypothetical doubling of the atmospheric CO₂ concentration compared to pre-Industrial levels.

EL NIÑO: The warm phase of the El Niño – Southern Oscillation, involving weaker upwelling of cold water off the South American coast and weaker Pacific trade winds.

EMISSIONS SCENARIO: An assumed scenario of future greenhouse gas emissions (or resulting atmospheric concentrations) based upon assumptions regarding global economic activity, the prevalence of fossil fuel use, and (sometimes) global carbon cycle model estimates of the rate of CO₂ uptake by land and ocean.

ENDANGERMENT FINDING (EF): The 2009 finding by the Administrator of the Environmental Protection Agency that emissions of well-mixed greenhouse gases (primarily CO₂) endanger human health and welfare.

ENSO: El Niño – Southern Oscillation, a prominent natural climate fluctuation involving year-to-year variations in the upwelling of cold water off the equatorial coast of South America and associated strengthening or weakening of trade winds across the Pacific Ocean.

FACE: (Free-Air CO₂ Enrichment) Large-scale open-air studies that expose plants to elevated CO₂ levels, simulating future climate scenarios.

FOSSIL FUEL: A natural hydrocarbon fuel such as coal, oil, or natural gas formed in the geological past from the remains of living organisms.

FUND (Framework for Uncertainty, Negotiation, and Distribution): An Integrated Assessment Model developed by Richard Tol in 1997.

GBR (Great Barrier Reef): The world's largest coral reef ecosystem located off the northeast coast of Australia.

GDP (Gross Domestic Product): Total market value of all the final goods and services produced within a country's border, usually over one year.

GHCN (Global Historical Climatology Network): A time-varying number of globally distributed weather stations making hourly, daily, or monthly measurements of precipitation, temperature and sometimes snowfall and snow depth.

GLOBAL GREENING: The satellite-observed increase in greenness in most vegetated land areas observed since the early 1980s.

GNP (Gross National Product): Total market value of all the final goods and services produced by the citizens of a country, including exports and imports, usually over one year.

GDP (Gross National Product): Total market value of all the domestically-produced final goods and services produced by the citizens of a country, usually over one year.

GREENHOUSE EFFECT (GHE): The tendency for any planetary atmosphere containing greenhouse gases to be warmer in the lowest layers than if those gases did not exist.

GREENHOUSE GAS (GHG): An atmospheric gas that absorbs and emits infrared radiation, especially water vapor, CO₂, and methane.

GROSS PRIMARY PRODUCTION: The total chemical energy used by plants during photosynthesis over a specified period of time. Net Primary Production is the residual after using the energy required for respiration and represents the contribution to biomass growth.

HURST PHENOMENON: See long term persistence.

HYDROLOGY: The study of the movement of water, especially on land and in the atmosphere.

IAM (Integrated Assessment Model): A computerized tool that combines economics, climate science, and social sciences to provide quantitative relationships between human and Earth systems, helping to inform policy decisions.

IEA (International Energy Agency): A Paris-based intergovernmental organization, established in 1974, that provides policy recommendations, analysis and data on the global energy sector.

IPCC (Intergovernmental Panel on Climate Change): A panel of international experts and governmental representatives established by the United Nations and World Meteorological Organization in 1988 to provide governments at all levels with scientific information that they can use to develop climate policies.

IR (Infrared): Heat radiation that all solid objects and greenhouse gases emit by virtue of their temperature.

LA NIÑA: The cool phase of ENSO, involving stronger upwelling of cold water off the South American coast and stronger Pacific trade winds.

MAIZE: Corn.

MODEL: A computer code or collection of codes which quantify our knowledge of various processes and the interactions between those processes.

MORTALITY: Death or number of deaths, usually expressed for a specific population and period.

NCA (National Climate Assessment): A periodic report mandated by the Global Change Research Act of 1990 summarizing the impact of climate change on the United States. There have been five NCA reports: NCA1 (2000), NCA2 (2009), NCA3 (2014), NCA4 (2017-18), and NCA5 (2023).

OECD (Organization for Economic Cooperation and Development): A group of 38 member countries that have a democratic system of government and free economic systems.

PALEOCLIMATE: Climate of the distant past, before modern weather instruments were widely used (typically pre-1850), necessitating the use of “proxy” measurements such as tree rings, ice core, pollen records, etc.

PDI (Power Dissipation Index): A statistical estimate of the accumulated destructive potential of a tropical cyclone over time, calculated as a summation of the cube of the maximum sustained wind speeds.

PDO: Pacific Decadal Oscillation, a recurring ocean-atmosphere interaction centered over the Pacific basin associated with predominantly warm or cool patterns that influence the global climate.

PHOTOSYNTHESIS: The process by which plants grow, requiring carbon dioxide, water, and light at sufficiently warm temperatures (10 to 35 deg. C for optimal function).

PPM: Parts per million, a measure of the concentration of a substance in (for example) the atmosphere

PRE-INDUSTRIAL: The near-term historical period before which human greenhouse gas emissions were considered significant, usually assumed to be the mid-1700s.

RADIATIVE FEEDBACK: A change in the balance between absorbed solar radiation and emitted infrared (IR) radiation, referenced to the top of the atmosphere, caused by changes in surface temperature.

RADIATIVE FORCING: The change in the balance between absorbed solar radiation and emitted infrared (IR) radiation, referenced to the top of the atmosphere, caused by changes in greenhouse gases, anthropogenic aerosols, volcanoes, etc.

RCP (Representative Concentration Pathway) SCENARIOS: Different scenarios for future greenhouse gas emissions and their impact on the atmosphere, introduced in the IPCC's Fifth Assessment Report (AR5). RCP scenarios have a number label (e.g. RCP6.0) indicating how much radiative forcing they assume in 2100 relative to pre-Industrial times, in Watts per square meter. RCP4.5 and RCP6.0 are intermediate scenarios, whereas RCP8.5 is an extreme scenario with very large future emissions of greenhouse gases.

REGRESSION: A statistical method for selecting the coefficients of a linear equation to explain the behaviour of a dependent variable in terms of variations in one or more explanatory variables.

RICARDIAN ANALYSIS: A method used in environmental economics named after David Ricardo to estimate the economic impacts of climate change, particularly on agriculture, by making use of the hypothesis that expected changes in the return to future agricultural activity will be capitalized into current land values.

SCC (Social Cost of Carbon): An estimate of the net economic damages, measured in present-value dollars, caused by emitting one additional ton of CO₂ into the atmosphere.

SSP (Shared Socioeconomic Pathway) SCENARIOS: Different scenarios for future greenhouse gas emissions and their impact on the atmosphere, introduced in the IPCC's Sixth Assessment Report (AR6). Some of the SSP scenarios are analogous to the older RCP scenarios, for purposes of continuity and comparison to earlier climate model assessments.

TC: Tropical cyclone.

TMAX: The daily maximum surface air temperature.

TMIN: The daily minimum surface air temperature.

TOA: Top of atmosphere.

TONNE: Metric ton.

TOXICOLOGY: The study of the adverse effects of chemical substances on living organisms.

TREND: A coefficient, usually estimated using linear regression, representing the slope of a line of best fit through a time series of data, indicating any tendency for the series mean to move up or down over time.

TSI (Total Solar Irradiance): The measure of the total amount of incident solar radiation per unit area, including all wavelengths of electromagnetic radiation, that reaches the Earth's atmosphere. Typically shown in units of watts per square meter (W/m²).

UHI (Urban Heat Island): The tendency for inhabited areas to be warmer than their rural surroundings, due to the replacement of natural land and vegetation with roads, parking lots, buildings, and waste heat sources.

USHCN: The U.S. Historical Climatology Network, consisting of quality-controlled temperature and precipitation data from 1,218 surface weather stations.

METADATA FOR FIGURES AND TABLES

Figure 2.1 Screen shot from Figure 3 of Zhu *et al.* 2016 <https://www.nature.com/articles/nclimate3004>

Figure 2.2 Screen shot from Gerhart and Ward (2010) Figure 2
<https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-8137.2010.03441.x>

Figure 2.3 Screenshot from Copernicus Marine Services
https://data.marine.copernicus.eu/product/GLOBAL_OMI_HEALTH_carbon_ph_area_averaged/description

Figure 2.4 Screen shot from AIMS (2023) Figures 2-4
https://www.aims.gov.au/sites/default/files/2022-08/AIMS_LTMP_Report_on%20GBR_coral_status_2021_2022_040822F3.pdf

Figure 3.1.1 Screen shot of AR6 WG1 Ch2 Fig 10

Figure 3.1.2 Screen shot of AR6 WG1 Ch7 Fig 7-6

Figure 3.1.3 Author created figure. CO2 data source: <https://gml.noaa.gov/ccgg/trends/index.html>

Figure 3.2.1 Screen shot from Hausfather *et al.* (2019) Figure S4 <https://doi.org/10.1029/2019GL085378>

Figure 3.2.2 Author created figure. Data source: Freidlingstein *et al.* (2024)
<https://essd.copernicus.org/preprints/essd-2024-519>
Data source: <https://globalcarbonbudget.org/download/1442/>

Figure 3.2.3 Author-created using OLS trends, Data source:
<https://globalcarbonbudget.org/download/1442/>

Figure 3.2.4 Author-created using OLS trends, Data source:
<https://globalcarbonbudget.org/download/1442/>

Figure 4.1 Screen shot from Scafetta (2021) Figure 1 <https://doi.org/10.3390/cli9110161>

Figure 5.1 Data source: <https://climexp.knmi.nl/start.cgi>. Author-created figure. One ensemble member per model, yearly average temperatures are computed from the global monthly raw temperatures; ‘range’ is the warmest model temperature minus the coolest model temperature in each year; standard deviation computed across all 33 models’ raw temperatures in each year.

Figure 5.2 Screen shot from Scafetta (2023) Figure 2
<https://doi.org/10.1007/s00382-022-06493-w>

Figure 5.3 Author created figure. Data source: <https://climexp.knmi.nl/start.cgi>. One SSP370 ensemble member per model. Linear OLS trends computed from global monthly temperature anomalies relative to 1981-2010.

Figure 5.4 Author created figure, based on Figure 3 of McKittrick and Christy (2020)
<https://doi.org/10.1029/2020EA001281>. Updated using same methods and data extended from 2014 to 2024, see McKittrick and Christy (2025) in chapter references.

Figure 5.5 Screen shot of IPCC AR5 Figure 10.SM1, with author annotations

Figure 5.6 Author created figure following Christy and McNider (2017) <https://doi.org/10.1007/s13143-017-0070-z> with updated data available at

https://www.nsstc.uah.edu/data/cmip6/ERA5_REANALYSES/ and
https://www.nsstc.uah.edu/data/cmip6/JRA3Q_REANALYSES/ compiled here
https://www.nsstc.uah.edu/data/cmip6/CWG25_Fig_5.7_Table_250627.xlsx

Figure 5.7 Screen shot from

https://climate.rutgers.edu/snowcover/chart_seasonal.php?ui_set=nhland&ui_season=1 (accessed May 27, 2025)

Figure 5.8 Screen shot from Rugenstein and Hakuba (2023) Figure 1

<https://doi.org/10.1029/2022GL101802>

Figure 5.9 Author created figure. Data source: <https://climexp.knmi.nl/start.cgi>. OLS linear trends. One SSP370 ensemble member per model, averaged surface air temperatures for June through August over 31N to 49N latitude, 84W to 102W longitude. Observed temperature data for same months from NOAA/NCEI at <https://www.ncei.noaa.gov/access/monitoring/climate-at-a-glance/statewide/time-series> for the 12 Corn Belt states averaged together.

Figure 6.1.1 Author replot of Figure 10 from Koutsoyiannis (2013)

<http://dx.doi.org/10.1080/02626667.2013.804626>

Figure 6.2.1 Screen shot of figure from Maue (2025) <https://climatlas.com/tropical/>

Figure 6.2.2 Author created figure. Data from National Hurricane Center (2024)

<https://www.nhc.noaa.gov/climo/>

Figure 6.2.3 Author created figure. Data from NOAA HRD

https://www.aoml.noaa.gov/hrd/hurdat/All_U.S._Hurricanes.html

Table 6.2.1 Author created table. Data from NOAA HRD

https://www.aoml.noaa.gov/hrd/hurdat/most_intense.html

Figure 6.3.1 Screen shot from NCA4 Figure 6.4

Figure 6.3.2 Author created figure. Data source: https://www.nsstc.uah.edu/data/ushcn_jrc/

Figure 6.3.3 Author created figure. Data source: https://www.nsstc.uah.edu/data/ushcn_jrc/

Figure 6.3.4 Author created figure. Data source: https://www.nsstc.uah.edu/data/ushcn_jrc/

Figure 6.3.5 Author created figure Data source: https://www.nsstc.uah.edu/data/ushcn_jrc/

Figure 6.3.6 Author created figure. Data source: https://www.nsstc.uah.edu/data/ushcn_jrc/

Figure 6.3.7 Screen shot from <https://www.globalchange.gov/indicators/heat-waves>

Box tables. Author created table. Data from McKittrick and Christy (2025)

Figure 6.4.1 Author created figure. Data from McKittrick and Christy (2025)

Figure 6.4.2 Author created figure. Data from McKittrick and Christy (2025)

Figure 6.4.3 Author created figure. Data from McKittrick and Christy (2025)

Figure 6.4.4 Author created figure. Data from McKittrick and Christy (2025)

Figure 6.4.5 Author created figure. Data from McKittrick and Christy (2025)

Figure 6.5.1 Author created figure. Data source: https://www.spc.noaa.gov/wcm/data/1950-2024_actual_tornadoes.csv

Figure 6.7.1 Author created figure. Data source: <https://www.ncei.noaa.gov/access/monitoring/uspa/wet-dry/0> accessed June 16, 2025. OLS trend line added.

Figure 6.8.1 Screen shot from Lizundia-Loiola *et al.* (2021) Figure 12.

Figure 6.8.2 Screenshot from Marlon *et al.* (2012) Figure 2 panel C.
<https://www.pnas.org/doi/pdf/10.1073/pnas.1112839109>

Figure 6.8.3 Author created Figure. Data 1926 to 2016:
https://web.archive.org/web/20200212033452/https://www.nifc.gov/fireInfo/fireInfo_stats_totalFires.html
. Post-2017 <https://www.nifc.gov/fire-information/statistics/wildfires> Accessed June 16, 2025.

Figure 7.1 Screenshot from <https://tidesandcurrents.noaa.gov/sltrends/>

Table 7.1 Author created table. From Section 6.2 of
<https://judithcurry.com/wp-content/uploads/2018/11/special-report-sea-level-rise-3.pdf>

Figure 7.2 Screenshot from https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=9414290
(downloaded 4/22/25)

Figure 7.3 Screenshot from
https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8771450 (downloaded 4/22/2025)

Figure 7.4 Screenshot from
https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8761724 (downloaded 4/22/25)

Figure 7.5 Screenshot from
https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8518750 (downloaded 4/22/25)

Figure 7.6 Author created figure. Data from
https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?id=8518750

Figure 8.1 Screenshot from
<https://crudata.uea.ac.uk/cru/data/temperature/HadCRUT5.0Analysis.pdf>

Figure 8.2 Screenshot from Hansen and Karecha (2025) Figure 1
<https://www.columbia.edu/~jeh1/mailings/2025/CloudFeedback.13May2025.pdf>

Table 8.1 Screenshot of column 1 of Table 12.12 IPCC AR6 Working Group I report

Figure 9.1 Screenshot From Taylor and Schlenker (2021) Figure 1
<https://www.nber.org/papers/w29320>

Figure 9.2 Screenshot from McKittrick (2025) Figure 1
<https://www.nature.com/articles/s41598-025-90254-2>

Figure 10.1 Screenshot from Pielke (2024) <https://www.nature.com/articles/s44304-024-00011-0> Figure 3.

Figure 10.2 Screenshot from Gasparini *et al.* (2015) Figure 2
[http://dx.doi.org/10.1016/S0140-6736\(14\)62114-0](http://dx.doi.org/10.1016/S0140-6736(14)62114-0)

Figure 11.1 Screen shot from Pielke Jr (2023)
<https://rogerpielkejr.substack.com/p/global-disaster-losses1990-2023>

Figure 11.2 Screenshot from CEA-OMB (2023) <https://bidenwhitehouse.archives.gov/wp-content/uploads/2023/03/CEA-OMB-White-Paper.pdf> Figure 1.

ABOUT THE AUTHORS

John Christy, Ph.D. is the Distinguished Professor of Atmospheric and Earth Sciences and Alabama's State Climatologist at the University of Alabama in Huntsville. He has a B.A. (Mathematics) from CSU Fresno and an M.S. and Ph.D. (Atmospheric Sciences) from the University of Illinois in Urbana-Champaign. He is a Fellow of the American Meteorological Society and recipient of The NASA Medal for Exceptional Scientific Achievement for constructing, with Dr. Roy Spencer, the first global atmospheric temperature dataset from polar orbiting satellites. He served as Lead Author of the 3rd IPCC Assessment Report (2001), he coauthored the National Academy of Science expert report on Surface Temperature Reconstructions (2006), and he has served on the NOAA Climate Change Science Program, NASA's Earth Science Subcommittee, the National Research Council Space Studies Board and the EPA Science Advisory Board. In addition, Dr. Christy has given testimony in 20 Congressional Hearings.

Judith Curry, Ph.D. is Professor Emerita at the Georgia Institute of Technology, where she served as Chair of Earth and Atmospheric Sciences for 13 years. She is President and co-founder of Climate Forecast Applications Network (CFAN). Curry has a PhD in Geophysical Sciences from the University of Chicago. She has authored or coauthored 192 peer-reviewed papers in atmospheric and climate sciences, two textbooks, and most recently the book *Climate Uncertainty and Risk*. Curry is a Fellow of the American Meteorological Society, the American Association for the Advancement of Science, American Geophysical Union, and a Member of the American Academy of Sciences and Letters. She has served on several Science Steering Committees of the World Climate Research Program and also the DOE Biological & Environmental Research Advisory Committee, Earth Science Subcommittee of the NASA Advisory Council, and the NRC Space Studies Board and Climate Research Committee. Dr. Curry has given testimony in 13 Congressional Hearings..

The Honorable Steven E. Koonin, Ph.D. is the Edward Teller Senior Fellow at Stanford's Hoover Institution. He has served as Under Secretary for Science at the Department of Energy (2009-2011), as Chief Scientist for British Petroleum (2004-2009), and as a professor at Caltech (1975 – 2004, the last nine years as Vice President and Provost). Koonin is a member of the National Academy of Sciences and the JASON group of government advisors, as well as a Governor of Lawrence Livermore National Laboratory, having served in similar capacities for the Los Alamos, Sandia, Brookhaven, and Argonne National Laboratories. Other prior roles include a University Professor at New York University, a non-resident Senior Fellow at the American Enterprise Institute, and a Trustee of the Institute for Defense Analyses. Koonin has a BS in Physics from Caltech and a PhD in Theoretical Physics from Massachusetts Institute of Technology. He authored the 2021 bestseller *Unsettled: What Climate Science Tells Us, What It Doesn't, and Why It Matters*, the 1985 textbook *Computational Physics*, and some 200 peer-reviewed papers in physics and astrophysics, scientific computation, energy technology and policy, and climate science.

Ross McKittrick, Ph.D., is a Professor of Environmental Economics at the University of Guelph in Ontario, Canada. He holds a BA in economics from Queen's University, and an MA and Ph.D. in economics from the University of British Columbia. He has published widely in both economics and physical science journals on topics related to climate change, pollution and public policy. His book *Economic Analysis of Environmental Policy* was published by the University of Toronto Press in 2010. His background in applied statistics has led him to author or coauthor peer-reviewed publications across a wide range of topics in the physical sciences including paleoclimate reconstruction, malaria transmission, surface temperature measurement, climate attribution methodology and climate model evaluation. Professor McKittrick has made many invited academic presentations around the world, was an invited contributor to the 2006

National Academy of Sciences report on Surface Temperature Reconstructions, has served as an expert reviewer for the last three IPCC Assessment Reports (Working Groups I and II) and has testified before the U.S. Congress and committees of the Canadian House of Commons and Senate.

Roy W. Spencer, Ph.D., is a Principal Research Scientist at the University of Alabama in Huntsville. He holds a BS in Atmospheric and Oceanic Science from the University of Michigan, and MS and PhD in Meteorology from the University of Wisconsin -Madison. Spencer's published research has focused on monitoring of weather processes and climate change from Earth-orbiting satellites, as well as using simple physical models for diagnosing climate sensitivity and climate feedbacks. As a NASA Senior Scientist for Climate Studies he served as the U.S. Science Team Leader for the Advanced Microwave Scanning Radiometer (AMSR-E) flying on the Aqua satellite, providing our most accurate global measurements of sea surface temperatures, sea ice, and several other features of the climate system. Along with Dr. John Christy, Spencer developed the first satellite-based technique for global temperature monitoring, for which they received NASA's Exceptional Scientific Achievement Medal and the American Meteorological Society's Special Award.

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Monday, May 19, 2025 6:36 PM
To: Roy W. Spencer; Judith Curry; John Christy; Travis Fisher; Steven Koonin
Subject: post-meeting drafts

I have opened a new folder called Post-meeting versions which will hold the drafts as they stand after we have gone through them.

You should all have access to them but if not let me know.

With regards to the remaining minor changes in Ch 5 in order to control the process when you change a draft if it's just a revised figure and caption, or a new paragraph, you can send it to me via email and I'll insert it, but if you change the draft in the Dropbox folder add [AB] initials to the file name so we keep track of the version change.

I didn't upload extreme weather yet since we have work to do on it tomorrow.

From: Ross McKittrick [ross.mckittrick@gmail.com]
Sent: 7/4/2025 8:05:30 PM
To: Judith Curry [curry.judith@gmail.com]; Steven Koonin [steven.koonin@gmail.com]; John Christy [climatemanager60@gmail.com]; Roy W. Spencer [roywspencer@hotmail.com]; Travis Fisher [travis.scott.fisher@gmail.com]; Josh Loucks [loucksj14@gmail.com]; Cohen, Seth [seth.cohen@hq.doe.gov]
Subject: Post-meeting edition
Attachments: NCAS.ReviewJuly4.docx

I applied the agreed-upon changes (Sct 3.3 & 4.1) and cleaned things up. I added some language about trend analysis (green font, Sct 4.1); pls check. See also proposed name for Ch 4.

Happy Independence Day everyone

Critique of the Fifth National Climate Assessment

EXECUTIVE SUMMARY	3
1 INTRODUCTION	4
1.1 ABOUT THIS REPORT	4
1.2 ABOUT THE NATIONAL CLIMATE ASSESSMENTS	4
1.3 STANDARDS FOR ASSESSING NCA PRODUCTS	5
1.4 WHEN PEER REVIEW FAILED	6
2 NCA5 OVERVIEW CHAPTER: HYPERBOLIC ADVOCACY LANGUAGE AND UNSCIENTIFIC TONE.....	9
3 OVERCONFIDENCE WHEN STATING CONCLUSIONS	11
3.1 INFLATED CERTAINTY LEVELS	11
3.2 OVERSTATING FINDINGS IN UNDERLYING SOURCES	12
3.3 TECHNICAL EXAMPLE: EMERGENT CONSTRAINTS	13
4 ARBITRARY AND AMBIGUOUS TREND ANALYSIS.....	15
4.1 AVERAGE AND EXTREME PRECIPITATION.....	15
4.2 HEATWAVES.....	16
5 INADEQUATE TRACEABLE ACCOUNTS.....	19
6 OVEREMPHASIS ON EXTREME SCENARIOS	21
7 CONFLATION OF NATURAL AND ANTHROPOGENIC CLIMATE CHANGES	23
8 SUMMARY EVALUATION OF NCA5 AND RECOMMENDATIONS FOR NCA6.....	24
8.1 COMPLIANCE WITH USGCRP'S STATUTORY OBLIGATION	24
8.2 ADHERENCE TO THE TENETS OF GOLD STANDARD SCIENCE	25
8.3 SUITABILITY FOR INFORMING FEDERAL CLIMATE AND ENERGY POLICY	26
8.4 RECOMMENDATIONS FOR NCA6.....	26
9 REFERENCES	28
APPENDICES.....	30
APPENDIX 1: CHARGE TO THE REVIEW	30
ABOUT THE AUTHORS.....	31

EXECUTIVE SUMMARY

This report critically evaluates the Fifth National Climate Assessment (NCA5), finding it an inadequate basis for climate and energy policies. A detailed review of the foundational chapters on climate science raises several key concerns:

Scientific Integrity

- An advocacy tone, particularly in the Overview chapter, that is inconsistent with scientific objectivity
- Endemic overconfidence when stating conclusions, including inflated certainty levels and overstatement of findings from underlying sources

Flawed methodology

- Use of short time periods for trend analyses; more generally lack of a consistent rule for choosing time scales
- Overemphasis on extreme scenarios for future emissions and sea level rise
- Inadequate traceable accounts to support key conclusions

Deficient Review Process

- A review by the National Academy of Sciences, Engineering, and Medicine (NASEM) failed to challenge some fundamental flaws in NCA5
- The NCA5 authors failed to adopt some improvements recommended by NASEM

The report shows NCA5 to be a biased and incomplete picture of climate science that skews toward excessive climate risk. Consequently, NCA5 should not be used as basis for climate and energy policies.

1 INTRODUCTION

1.1 About this Report

This report responds to a request from the Office of Science and Technology Policy (OSTP) Director Michael Kratsios (reproduced in Appendix 1) that we assess the suitability of the Fifth National Climate Assessment (NCA5 2023) for informing climate and energy policies.

To conduct this assessment, we reviewed NCA5 itself, the National Academies of Science, Engineering and Medicine (NASEM) review of the Third Order NCA5 Draft, and the USGCRP responses to that review. We have drawn upon our long experience working on advisory reports, including for the NASEM, the Intergovernmental Panel on Climate Change (IPCC) and other organizations, as writers, reviewers and study leaders.

We find that NCA5 should not be relied on for policy making as it is an incomplete and misleading picture of our understanding of the changing climate and the risks it poses to the nation. NCA5 was selective and incomplete in its survey of climate science. Further, we have reviewed recent scientific developments in a separate report to Energy Secretary Wright (Climate Working Group 2025, herein denoted “CWG25”). The overall effect of the biases and omissions of NCA5 is an excessively dire picture of climate risk.

To support our finding, the following sections of this chapter provide some background on the National Climate Assessments and the standards by which we have assessed NCA5. Chapters 2 through 7 detail important ways in which NCA5 fails to meet those standards. Finally, Chapter 8 summarizes our evaluation of NCA5 and makes recommendations for NCA6.

1.2 About the National Climate Assessments

The U.S. Global Change Research Program (USGCRP) was established by Presidential Initiative in 1989 and mandated by Congress in the Global Change Research Act (GCRA) of 1990 to develop and coordinate “a comprehensive and integrated United States research program which will assist the Nation and the world to understand, assess, predict, and respond to human-induced and natural processes of global change.”

The National Climate Assessment (NCA) Reports evolved from the GCRA requirement to:

Prepare and submit to the President and the Congress an assessment which (1) integrates, evaluates, and interprets the findings of the Program and discusses the scientific uncertainties associated with such findings; (2) analyzes the effects of global change on the natural environment, agriculture, energy production and use, land and water resources, transportation, human health and welfare, human social systems, and biological diversity; and (3) analyzes current trends in global change, both human-induced and natural, and projects major trends for the subsequent 25 to 100 years.

The five NCA Reports to date have interpreted this mandate in different ways:

- NCA1 (2000) was a multidisciplinary effort to study and portray in regional detail the potential effects of human-induced global warming on the United States.

- NCA2 (2009) was a relatively short report that not only focused on the potential consequences of climate change but also sought to identify potential adaptation measures and identify the highest priorities for future research.
- NCA3 (2014) was an expanded effort that continued along the lines of NCA2, and was distinguished by a comprehensive review process, including a review by the National Academy of Sciences. NCA3 introduced a traceability analysis for each key finding, including a description of uncertainties.
- NCA4 (2017/2018) was a further expanded effort that was published in two volumes: I Climate Science Special Report (CSSR), and II Impacts, Risk, and Adaptation in the United States. This was the first NCA report to devote significant effort to assessing climate science.
- NCA5 (2023) returned to a single volume format, reducing discussion of climate science to a cursory two chapters, with continuing focus on regional and sectoral impacts of climate change. NCA5 expanded its remit to topics that are arguably outside the purview of the USGCRP, focusing on “Climate Action” and policy recommendations with a social justice emphasis.

These successive interpretations of the NCA are poorly aligned with the GCRA’s statutory requirements. Only NCA4 gives serious consideration to climate science, which receives the great majority of USGCRP funding, and even then NCA4 misrepresents some important points. Natural climate variability is essentially dismissed as unimportant in NCA4 and barely mentioned in NCA5 in most contexts, despite the mandate to include both human-caused and natural climate change. Most significantly, NCA5 has redirected the focus away from climate science and even the hazard elements of impacts, towards evaluation of societal vulnerabilities and advocacy for particular policy options, especially deployment of wind and solar generation. The cover image for the NCA5 Report is the installation of a solar panel.

1.3 Standards for Assessing NCA Products

Beyond the statutory requirements stated above, another pertinent standard is President Trump’s executive order of May 23, 2025, Restoring Gold Standard Science. It enumerates some of the characteristics of good science, namely that it should be:

- i. reproducible;
- ii. transparent;
- iii. communicative of error and uncertainty;
- iv. collaborative and interdisciplinary;
- v. skeptical of its findings and assumptions;
- vi. structured for falsifiability of hypotheses;
- vii. subject to unbiased peer review;
- viii. accepting of negative results as positive outcomes; and
- ix. without conflicts of interest.

Although these should be self-evident to those trained in science, it is surprising that some scientists have found them controversial (Dellawalla et al. 2025).

====debated text:

When science is assessed for the purpose of climate policymaking there are further requirements to bear in mind. Good climate and energy policies should balance the risks and benefits of a changing

climate against the world's growing demand for reliable and affordable energy. To strike that balance, policymakers must understand at least two complex and nuanced topics: the certainties and uncertainties of climate science and the technoeconomic realities of energy systems. Experts in those subjects have a responsibility to communicate their understanding to non-experts clearly, completely, and without bias.

Much expertise in energy systems resides in the private sector, where it is essential to providing the reliable and affordable energy services that are the lifeblood of society. Understanding is therefore continually refined and tested by operational demands and the discipline of the marketplace. In contrast, climate science expertise is found largely in the non-profit realm of universities and government laboratories and so has a more academic character, with greater uncertainties and hence need for expert judgement.

=====

For those reasons assessments of climate science that will be influential for policy making must include statements about the strength of knowledge. The IPCC's approach to assessing strength of knowledge is based on judgment of the available evidence and agreement among experts. But there are more sophisticated knowledge characterizations for risk management. Here are the elements according to Aven (2017):

- The degree to which the assumptions made are reasonable/realistic based on background knowledge
- The degree to which data/information exists and are reliable and relevant
- The degree to which there is disagreement among experts (including those from different environments or academic fields)
- The degree to which the phenomena involved are understood and accurate models exist
- The degree to which the knowledge has been thoroughly examined with respect to unknown knowns (issues that are known to matter but are ignored, such as, in this context, natural variability).

1.4 When Peer Review Failed

Peer review of reports like IPCC and NCA products is not a silver bullet that guarantees quality. The peer review process for these reports differs in some important respects from traditional peer review in academic journals. First, during the open review phases of early NCA drafts reviewers are volunteers who self-select what parts of the report to read, whereas the editors of academic journals editors recruit reviewers with relevant expertise and assign them to specific papers. As a result, there is no guarantee at that stage that any particular section of an NCA report was reviewed by anyone, let alone by someone with relevant expertise and a perspective that differs from the authors. In fact, many potential reviewers have given up over the years because their input has been systematically ignored. The introduction of a NASEM review stage is at best a partial remedy for this. However, NCA authors have near complete authority whether to accept or reject reviewer comments. By contrast, the Editor of an academic journal mandates the author to respond, and the reviewer and Editor assess its adequacy. Finally, the work of an NCA chapter author team is guaranteed to be published, whereas academic journals routinely reject papers that do not pass review.

These points are important to bear in mind because some readers might dismiss any criticism of NCA products out of hand on the grounds that they were "peer reviewed". But the test of a report is whether its contents are a valid representation of the available evidence. Here we give an example in

which NCA4 gave prominence to a figure that completely misrepresented the underlying data and which evaded peer review to do so. Further details can be found in Koonin (2024).

NCA4's Part I says (prominently and with *Very High Confidence*) on Page 19 of the Executive Summary:

There have been marked changes in temperature extremes across the contiguous United States. The number of high temperature records set in the past two decades far exceeds the number of low temperature records.

and offers this figure in support:




FIGURE 1: Reproduction of Figure ES.5 from NCA4

The figure appears to validate the claim that record warm daily temperatures are occurring more often. But the underlying data do not. Rather, they show that extreme heat in the U.S. peaked in the 1920s and 1930s and declined thereafter, as supported by Figure 6.4 deep within the CSSR itself (reproduced in Section 4.2 of this report).

Figure ES.5 does not even show what its title implies, as it displays not simple counts of cold and warm extreme events, but rather ratios of the two counts. As defined in the analysis, both types of extremes declined over the sample interval, but cold events declined more than warm events, so the ratio of warm extremes to cold extremes increased. By graphing the ratio instead of counts, NCA4 created a visual impression opposite to what is in the data.

Figure 6.3.3 of CWG25 (reproduced below) shows the results of a proper analysis and presentation of the data – defining each record as the singular extreme in the sample interval, displaying numbers

of warm and cold records separately, and beginning the analysis in 1899 rather than 1930. It makes evident the exceptional heat of the 1920s and especially the 1930s, peaking in 1936. (The increased heat of the past five years is too brief to be deemed climatically significant and, in any event, occurred after the CSSR was published in 2017.) The frequency of cold records has declined, especially over the last quarter of the period in which only 13 percent of the extreme cold events were measured. In contrast, 25 percent of the heat records were achieved in the last quarter, which is consistent with statistical expectations. As a result, the ratio of hot to cold extremes shown in Figure ES.5 of the CSSR explodes in recent decades. The overall reduction in numbers of both cold and hot extremes over the past century indicates a climate *less* prone to extremes, in sharp contrast to the message of NCA4 Figure ES.5.




FIGURE 2: Number of daily record High (red) and Low (blue) temperatures for warm and cold seasons in the CONUS. The lines represent the 15-year running, centered average. Source: 1,211 USHCN stations supplemented as needed to achieve a minimum of 92 percent of observations in the 126-year period since Dec 1898. US48: contiguous U.S. states. Tmax: maximum temperature. Tmin: minimum temperature. Source: CWG25 Figure 6.3.3.

How did the flawed figure ES.5 pass the NCA4 NASEM Review? The NASEM Review panel criticized the discussion of temperature extremes in a late draft of the CSSR. Key Finding 2 of Chapter 6 in that draft stated:

Accompanying the rise is [sic] average temperatures, there have been – as is to be expected – increases in extreme temperature events in most parts of the United States. Since the early 1900s, the temperature of extremely cold days has increased throughout the contiguous

United States, and the temperature of extremely warm days has increased across much of the West. In recent decades, intense cold waves have become less common while intense heat waves have become more common. (*Extremely likely, Very high confidence*)

In reference to this NASEM said:

Further, it is difficult to understand how a statement that includes increases in extreme warmth can be associated with a high confidence or extremely likely statement, given that most of the graphics in this chapter show a decrease in extreme warmth in the historical record.

The Federal official in charge of the CSSR (Michael Kuperberg, Executive Director of the USGCRP) responded to the Academies' review by noting that

Almost every recommendation from the NAS was incorporated [in the final version]... and a new figure was added on changes in record high and low temperatures

The "new figure" referred to is the misleading Figure ES.5, which did not appear in the draft reviewed by the Academies' panel. It seems unlikely that the Academies panel ever saw that figure because they surely would have commented if they had and it wouldn't have survived to publication. And the figure's problems were evidently not flagged (or were ignored) in the subsequent internal government reviews.

With this background from NCA4, we now present our critique of NCA5.

2 NCA5 OVERVIEW CHAPTER: HYPERBOLIC ADVOCACY LANGUAGE AND UNSCIENTIFIC TONE

NCA5 Chapter 1, "Overview", holds a prominent place in the Report since it summarizes the entire report and is the only section that many will read. Although some material in the later chapters of NCA5 is not without merit, Chapter 1 is marred by an activist tone and language that aggressively promotes mitigation policies. As such, this chapter adds an imprimatur of scientific authority to naïve and simplistic declarations of opinion about public policy. Many examples can be cited; here we note a few that are particularly egregious.

Chapter 1 opens as follows:

The more the planet warms, the greater the impacts. Without rapid and deep reductions in global greenhouse gas emissions from human activities, the risks of accelerating sea level rise, intensifying extreme weather, and other harmful climate impacts will continue to grow. Each additional increment of warming is expected to lead to more damage and greater economic losses compared to previous increments of warming, while the risk of catastrophic or unforeseen consequences also increases.

This statement presupposes that all warming is anthropogenic, that all warming is bad and causes economic damage, and that warming intensifies extreme weather. Such sweeping generalizations go well beyond what science has or can establish. The statement also refers to growing risks without saying whether they were large to begin with – a small risk that grows may still be a small risk, and a large risk may be only incrementally worsened.

On the next page the Overview states:

Each additional increment of warming leads to greater risks [here it lists 7 key components of society including food security and the economy]... Without deeper cuts in global net emissions, climate risks to the US will continue to grow... Action to limit future warming and reduce risks can have near-term benefits and opportunities: low-carbon energy jobs, Improved air quality, Health benefits, Economic benefits...”

The authors write as if climate policy does not involve trade-offs and all measures to reduce future warming will be effective and beneficial. Neither the vast body of scholarship on climate science and economics, nor historical experience, support such simplistic claims (CWG25 ch.XX). Not every additional increment of warming leads to economic harms; in fact, NCA5 Chapter 19 acknowledges that there will also be benefits. Nor is climate policy guaranteed to benefit the economy; it could impose net costs. Such misleading statements are a disservice to policymakers.

Pages 8 and 9 of the Overview promote renewable energy options by noting, correctly, that the levelized costs of wind/solar generation have fallen sharply since 2010, and so offers this rationale for pursuing deep emission reductions:

Recent growth in the capacities of wind, solar, and battery storage technologies is supported by rapidly falling costs of zero- and low-carbon energy technologies, which can support even deeper emissions reductions. For example, wind and solar energy costs dropped 70% and 90%, respectively, over the last decade, while 80% of new generation capacity in 2020 came from renewable sources.

Yet it has long been known that intermittent renewables generation must be supplemented by dispatchable generation to meet grid reliability standards, and that doing so (e.g., through natural gas, fission, or storage) significantly increases the cost of electricity. Further, there are yet unmet synchronization challenges in electrical grids with heavy penetration of wind and solar generation. Here again it is advisory malpractice not to acknowledge these facts. But more importantly, we see no reason for NCA5 to be discussing renewable energy technologies and economics in the context of its mandate.

Page 14 of the Overview presents future emission scenarios with the following description: “Climate modeling experts develop global climate projections for a range of plausible futures.” The word “plausible” is applied to all the scenarios. The accompanying figure includes both SSP3-7.0 and SSP5-8.5, the latter reaching the same radiative forcing as RCP8.5. The figure does not compare observed emissions with previous projections, nor does it caution readers that SSP3-7.0 and SSP5-8.5 are extreme scenarios that, like RCP8.5, have been heavily criticized in the scientific literature as being implausible (CWG25 3.2.1). By describing the entire envelope of emission projections as “plausible” the report opens the door to later presenting impacts associated with implausible worst case emission scenarios as part of a baseline “business-as-usual” future.

Page 17 of the Overview introduces the discussion of risks from extreme weather events by saying “Harmful impacts from more frequent and severe extremes are increasing across the country.” This creates the impression that extreme weather events are worsening and that disaster losses are driven by trends in extreme weather. While the section goes on to acknowledge that some of the increase is due to increases in the value of assets at risk, it then invokes the NOAA Billion Dollar Dataset as evidence of extreme weather getting worse. Scientists have shown that this is an invalid metric (CWG 10.2) since it measures the growth of the economy not the frequency of extreme weather, and indeed NOAA has recently withdrawn the data product (Pielke Jr. 2024, 2025). Even though the dataset was still distributed by NOAA at the time of the NCA5, the report authors failed to assess it skeptically and see that it was unfit for the purpose of measuring climate trends.

Page 23 of the Overview states, in large font:

Climate changes are making it harder to maintain safe homes and healthy families; reliable public services; a sustainable economy; thriving ecosystems, cultures, and traditions; and strong communities.

It is pointless to try to evaluate this statement on scholarly grounds; it is self-evidently hyperbolic. This kind of language doesn’t belong in a scientific assessment report. Its prominent inclusion points not only to the willingness of NCA5 authors to promote alarm, but also the ineffectiveness of the review process in eliminating such statements.

3 OVERCONFIDENCE WHEN STATING CONCLUSIONS

3.1 Inflated certainty levels

Overconfidence in stating conclusions has been an endemic problem in NCA Reports. The authors frequently make sweeping generalizations without appropriate caveats or acknowledging uncertainties. The overconfidence begins with inflated language to describe certainty levels.

NCA5 uses specific terms to convey information about scientific confidence associated with important finds, observations and projections. The Front Matter of NCA5 states that

confidence in a finding is based on the type, amount, quality, strength, and consistency of evidence; the skill, range, and consistency of methods to detect, evaluate, attribute, and interpret climate trends; and the degree of agreement across scientific information sources.

and Table 1 from the Front Matter provides calibrated language for confidence assessment:

Very high:	Strong evidence (established theory, multiple sources, well documented and accepted methods, etc.). High consensus
High:	Moderate evidence (several sources, some consistency, methods vary and/or documentation limited, etc.). Medium consensus
Medium:	Suggestive evidence (a few sources, limited consistency, methods emerging, etc.) Competing schools of thought.
Low:	Inconclusive evidence (limited sources, extrapolations, inconsistent findings, poor documentation and/or methods not tested, etc.). Disagreement or lack of opinions among experts.

This terminology is obviously inflated: *high confidence* with just moderate evidence and medium consensus; *medium confidence* with only suggestive evidence and competing schools of thought. This inflation in terminology leads to overconfidence in conclusions.

Even with this inflated terminology, NCA5 further inflates the confidence level of many of its conclusions to well beyond what the IPCC has concluded with more thorough justification, and by incorrectly applying its own standards. An example is the conclusion about anthropogenic influence on climate change, arguably one of the most important scientific statements of the report.

Key Message 3.1 Traceable Account states:

There is *very high confidence* that emissions of GHGs from human activities, fossil fuel use in particular, have unequivocally caused *all* global warming observed over the industrial era.

while the corresponding statements from IPCC AR5 and AR6 are

AR5: It is *extremely likely* that *more than half* of the observed increase in global average surface temperature from 1951 to 2010 was caused by the anthropogenic increase in greenhouse gas concentrations and other anthropogenic forcings together. (IPCC, 2013)

AR6: The *likely* range of total human-caused global surface temperature increase from 1850–1900 to 2010–2019 is 0.8°C to 1.3°C, with a best estimate of 1.07°C. It is *likely* that well-mixed GHGs contributed a warming of 1.0°C to 2.0°C, other human drivers (principally aerosols) contributed a cooling of 0.0°C to 0.8°C, natural drivers changed global surface temperature by –0.1°C to +0.1°C, and internal variability changed it by –0.2°C to +0.2°C. It is *very likely* that well-mixed GHGs were the main driver of tropospheric warming since 1979.

The IPCC AR5 and AR6 statements provide likelihood assessments but not a confidence level. While NCA5 claims with *very high confidence* that fossil fuels have unequivocally caused *all* global warming, AR5 refers to “*more than half* of the observed warming” and AR6 provides a more nuanced statement that relates to a *likely* range of warming contributed to by human drivers.

3.2 Overstating findings in underlying sources

Although we did not do a detailed audit of NCA5’s use of source material, spot checks suggest a pattern of overstatement.

Chapter 2 page 11 reports on an increase in precipitation in some areas and attributes it to “climate change”:

Many eastern regions of the country are getting wetter (Figure 2.4). Average annual precipitation from 2002-2021 was 5%-15% higher relative to the 1901-1960 average in the central and eastern US, a trend attributable to climate change.

The claims were attributed to Knutson and Zeng (2018). But Knutson and Zeng did not say these things nor analyze this particular period. They said (emphasis added):

Observed annual precipitation trends over the recent 30-yr period 1981–2010 (Fig. 5) have generally stronger magnitudes (in mm yr⁻¹ decade⁻¹) than observed trends for 1901–2010 or 1951–2010. This is as expected since trends over a shorter period have stronger potential for occurrence of pronounced temporary trends due to internal variability alone. **Only 9% of the analyzed area has detectable and attributable anthropogenic influence (categories -3, -2, +2, +3), which represents less than one-third of the fractional areal coverage of such trends for 1901–2010 (29%).** ... Eight percent of the analyzed area has attributable anthropogenic *increases* (categories +2, +3), including parts of eastern Europe, northern Asia, northern Australia/Maritime Continent, and tropical Africa.

Thus for 1981–2010 they cautioned that observed changes are consistent with natural variability, and they did not claim attributable trends were observed over the United States.

As another example of overstating findings, NCA5's Chapter 2 page 18 states

There is robust evidence that human-caused warming has contributed to increases in the frequency and severity of the heaviest precipitation events across nearly 70% of the US.

This is apparently attributed to two papers Diffenbaugh *et al.* (2018) and Kirchmeier-Young and Zhang (2020). Diffenbaugh *et al.* presented ratios of observed occurrences of extreme events to simulated occurrence rates in CMIP6 models run with only natural forcings. They report that for 79% of the U.S., extreme rainfall events are currently more frequent than in simulations, and the ratio of occurrences will rise under future warming simulations. This is not “robust” evidence of anthropogenic influence since it requires readers to assume that climate models accurately simulate regional precipitation patterns, something that even IPCC AR6 admits is uncertain (AR6 WGI SPM Box SPM 1.2, SPM Section C.1.3).

Kirchmeier-Young and Zhang looked at extreme 1- and 5-day events in North America and, using three global climate models, found

A signal of combined anthropogenic and natural external forcing can be detected in the intensification of extreme precipitation at the continental scale over North America. In some cases, this intensification can be attributed to human influence specifically.

However, they noted large differences among the three models in their simulated precipitation changes, and two of the three models failed to yield attribution for 5-day events for the North America mean. In other words, the models disagreed on whether the observed changes were anthropogenic. Both of the cited references utilized the implausible RCP8.5 scenario and began their “historical” observational records only in 1961. Here again this is not “robust” evidence, it is merely suggestive, particularly since other studies have found natural variability predominant.

3.3 Technical example: Emergent constraints

Some of the overconfidence expressed by NCA5 arises by omitting discussion of counterevidence in the peer reviewed literature. An example is the topic of “emergent constraints”. Chapter 3 page 17 invokes this concept to support its claim that key uncertainties in climate model projections have been reduced:

An approach to reduce uncertainty in climate change projections has matured over the past decade. It is known as "emergent constraints." The term refers to strong statistical relationships between highly uncertain future climate parameters and observable trends or variations in the current climate, along with a physical explanation of this relationship.

Emergent constraints appeared in the climate modeling literature as an attempt to tie climate model behavior to climate observations in hopes of generating insight into real-world equilibrium climate sensitivity to a doubling of carbon dioxide (ECS). Climate models exhibit a wide range of ECS values. The idea of emergent constraints is to examine whether the models with shared sensitivity values also share a common representation of some climate process (usually related to clouds) that might influence ECS. Suppose, for instance, that in low-ECS models the total cloud fraction in the tropics is strongly correlated with that in the southern midlatitudes, but in high-ECS models the correlation is weak. Then researchers could examine whether the observed correlation is strong or weak and draw a conclusion about which type of model, and therefore which ECS value, is more likely.

A problem in the emergent constraint literature is that there has been a proliferation of proposed correlations with little physical motivation. NCA5 alludes to this in the Traceable Accounts section (p. 3-40) but not in the main Chapter text. As shown by Schlund *et al.* (2020) many emergent constraints patterned on CMIP5 models (Coupled Model Intercomparison Project No. 5, used for the IPCC 5th Assessment Report) failed to work on CMIP6 models (used for the IPCC 6th Assessment Report). In other words, the observed correlations in CMIP5 models between ECS and cloud processes failed to show up in CMIP6 models, indicating that they were likely spurious rather than clues to physical processes governing the climate system. As a result, emergent constraints estimated on CMIP5 models *increased* the uncertainty of ECS, rather than decreasing it as intended and as claimed by NCA5.

Schlund *et al.* summarized the problem as follows [**boldface** added for emphasis]

In this study, we systematically analyze 11 published emergent constraints on ECS that have mostly been derived from models participating in the Coupled Model Intercomparison Project Phase 5 (CMIP5) project... The focus of the study is on testing if these emergent constraints hold for [Earth System Models] participating in the new Phase 6 (CMIP6). Since none of the emergent constraints considered here have been derived using the CMIP6 ensemble, CMIP6 can be used for cross-checking of the emergent constraints on a new model ensemble. **The application of the emergent constraints to CMIP6 data shows a decrease in skill and statistical significance of the emergent relationship for nearly all constraints, with this decrease being large in many cases.** Consequently, the size of the constrained ECS ranges (66% confidence intervals) widens by 51% on average in CMIP6 compared to CMIP5.

Yet NCA5 described the use of emergent constraint analysis as an advance that reduced uncertainty of ECS and climate model projections (p. 3-38):

Emergent constraints have been widely applied to reduce the simulated spread in the climate feedbacks that shape climate sensitivity.

4 ARBITRARY AND AMBIGUOUS TREND ANALYSIS

In CWG25 Chapter 6 we discussed the need for care in the choice of time scale when making claims about trends in extreme weather. We provided examples from analysis of river flooding, heatwaves, and extreme precipitation to show how assertions about apparent trends based on short data sets fall apart when looking at longer versions of the same data sets. NCA5 uses varying time scales for trend discussions, especially in Chapter 2 (which is focused on trends), without explaining why shorter or longer intervals were chosen. For example, page 2-4 defines temperature trends with respect to the interval since the late 1800s whereas page 3-29 presents temperature trends over 1972-2021. Page 2-11 compares rainfall over the 2002-21 versus 1901-1960 intervals while page 2-18 uses the interval since the late 1950s to compute trends.

A further problem is that NCA5 uses term “trend” repeatedly without indicating if it is statistically significant (*i.e.* distinguishable from random variability); the ambiguity creates the impression all the observed changes were significant, but readers are left to speculate. All climate variables will, in principle, drift up or down over any particular time interval. Identifying the change as a trend requires showing that it is statistically significant using a variance estimator suitable for the type of data being analysed. Of particular importance is choosing a method robust to serial correlation which is frequently present in climate data. Throughout Chapters 1 to 3 references are made to “trends” without explaining how they were computed. The exception was on page 1-19 which refers to urban temperature trends being calculated using the Theil-Sen estimator, although that method does not seem to have been used elsewhere. The one case where reference is made to statistical significance was on Page 2-9 referring to a trend in Atmospheric Optical Depth but no explanation is given of how significance was determined. Page 2-19 refers to regional trends in droughts being “robust” but again fails to explain how the term was defined.

In this chapter when we refer to calculation of trends in precipitation, we use ordinary least squares to compute the linear trend over the entire sample length and we use the Vogelsang and Franses (2007) variance estimator which is robust to general forms of serial correlation.

4.1 Average and Extreme Precipitation

McKittrick and Christy (2019) showed that trends in U.S. average and extreme precipitation that appeared to be positive and significant over the NCA4 interval (1901 – 2016) lost significance and/or changed sign on longer samples (extending back as far as the 1840s, depending on location) or on a shorter interval coinciding with the final 40 years of the data then available (1978 – 2017). In CWG25 Section 6.4 we presented analyses of updated data for the same locations (and many others), in all cases finding both considerable regional heterogeneity and that trends detectable on one time scale are often not detected on longer or shorter time scales. These findings argue against simple attribution of changes to anthropogenic influences.

NCA5 did not continue the NCA4 precedent of using the post-1901 interval as the time scale for its precipitation discussion. Instead it refers to changes “since the 1950s”; NCA5 Figure 2.8 suggests 1958 might be the intended start date. Under Key Message 2.2 “Extreme Events Are Becoming More Frequent and Severe” (p. 2-16) NCA5 states (p. 2-18):

Rainfall is becoming more extreme

Since the 1950s, there has been an upward trend in heavy precipitation across the contiguous US. This increase is driven largely by more frequent precipitation extremes, with relatively smaller changes in their intensity.

In addition to our concerns that inferences about trends are contingent on choice of time scale, if the intended meaning of this statement is that positive and significant trends are found in heavy precipitation events across CONUS we cannot reproduce it. We supplemented the data used in CWG25 (McKittrick and Christy 2025a) with 61 stations across the Midwest (Christy 2025). Many post-1950 trends are positive but statistically insignificant depending on the measure of extreme precipitation. It is possible to find locations that positive and significant trends, such as the Southeast post-1950, but those results are not robust to selecting longer or shorter time scales of analysis nor do they generalize across all regions (see details in McKittrick and Christy 2025b).

The following is a more accurate summary of what U.S. precipitation data show.

There are regionally heterogeneous changes in average and extreme precipitation. Positive trends in annual average rainfall and extreme events (defined either as the annual maximum or the number of exceedances of the 99th percentile value) are generally weak or non-existent when looking at samples exceeding 130 years in length. Significant trends can be found when beginning the sample in the 1950s although in most cases the trends weaken and become insignificant when looking at the post-1980 interval. This argues against attribution of changes in precipitation to anthropogenic warming.

4.2 Heatwaves

In contrast to NCA5, NCA4 examined heat extremes across the entire 20th century, pointing out that peak heat wave activity was reached in the 1930s and has decreased since then, and that recent increases are confined to areas west of the Rockies. NCA4 stated:

[The] warmest daily temperature of the year increased in some parts of the West over the past century, but there were decreases in almost all locations east of the Rocky Mountains. In fact, all eastern regions experienced a net decrease, most notably the Midwest (about 2.2°F [1.2°C]) and the Southeast (roughly 1.5°F [0.8°C]).

Since the mid-1960s, there has been only a very slight increase in the warmest daily temperature of the year (amidst large interannual variability). Heat waves (6-day periods with a maximum temperature above the 90th percentile for 1961–1990) increased in frequency until the mid-1930s, became considerably less common through the mid-1960s, and increased in frequency again thereafter. As with warm daily temperatures, heat wave magnitude reached a maximum in the 1930s.
(NCA4 pp. 190-191)




FIGURE 3. U.S. Heat waves since 1900. Reproduction of NCA4 Figure 6.4

Contrast this careful and factual language from NCA4 with NCA5 Key Message 2.2: “Extreme Events Are Becoming More Frequent and Severe.” This blanket statement is followed by NCA5 Figure 2.7 (reproduced as Figure 4 below) which shows the number of hot days, cold days and warm nights (respectively in panels a—c) over the period 2002-2021 compared to 1901-1960. [Presumably these are annual counts, although not stated in the caption.]




FIGURE 4: Reproduction of NCA5 Figure 2.7.

Close inspection shows that this figure actually contradicts the NCA5 Key Message with respect to heatwaves and instead reinforces the position of NCA4. Notwithstanding Key Message 2.2, the leftmost panel in NCA5 Figure 2.7 above shows that over most of CONUS the number of extreme heat days has *decreased* in recent decades compared to the prior century.

The accompanying discussion in NCA5 becomes convoluted as it appears to be aimed at supporting the Key Message despite the contrary evidence. NCA5 Chapter 2 elaborates on Key Message 2.2 by stating (page 16):

Observations show an increase in the severity, extent, and/or frequency of multiple types of extreme events. Heatwaves have become more common and severe in the West since the 1980s (*high confidence*).

But while it mentions the increase in hot days in the West, it fails to mention the decrease everywhere east of the Rockies. On the next page (page 17) it raises the comparison to the 1930s and then switches to discussing global counts:

By some measures, the most extreme heatwaves on record in the United States occurred during the Dust Bowl era of the 1930s. ... Globally such heatwaves are becoming more frequent, and in recent decades the western United States has been following those trends.

The reference to a global trend seems to be an attempt to tie heatwaves to anthropogenic climate change. But the obvious question, unaddressed by NCA5, is why one region follows the global pattern of increasing heatwaves while most of the country is trending in the opposite direction. These issues are addressed in Chapter 6 of CWG25, demonstrating that the range and magnitude of extreme temperatures (10s of degrees) are driven more by natural variability than by the small anthropogenic forcing. The analysis in CWG25 shows that for both (a) hot days ($\geq 95^{\circ}\text{F}$) and (b) number of days in heatwaves, the current decade is no hotter than a century ago in the U.S. Thus, the NCA5 claim that

“multiday heatwaves have become hotter, more frequent, larger, and longer lasting in recent decades,” is simply false when expanding the temporal and spatial domains to encompass all available data (Chapter 6 and Figure 6.3.6 of CWG25).

NCA5 then pointed to the dramatic Pacific Northwest heatwave of 2021. CWG2025 Section 8.6.1 discussed this event in detail, reviewing the multiple peer reviewed studies that explained the unusual meteorological context was unrelated to human-caused climate change, a point omitted from the NCA5 discussion.

NCA5 finally addresses the declining heatwave trend in CONUS east of the Rockies but attempts to reframe it by focusing on urban warm spells.

[The] number of very hot days has actually decreased across the central and eastern regions due to summer cooling trends in the region (Figure 2.7; Ch. 22). This does not, however, mean the central and eastern US are not affected by heat. The impacts of extreme high temperatures are more severe if such conditions persist for several days, and overall, multiday heatwaves have become hotter, more frequent, larger, and longer lasting in recent decades. Across 50 large US cities, the US Global Change Research Program heatwave indicator (<https://www.globalchange.gov/indicators/heat-waves>) shows that the average number of heatwaves has doubled since the 1980s, and the length of the heatwave season has increased from about 40 days to about 70 days.

We discussed this part of NCA5 in CWG25 Section 6.3. The figure referenced by NCA5 uses a short time period which begins in the 1960s, the coldest decade in the last 100 years, thus suppressing any comparison to the 1930s. Further, nighttime temperatures in urban areas have increased faster than daytime temperatures (which NCA5 acknowledges), but this is a signature of urban heat islands and not a good metric of climate change. CWG25 found that the occurrence of hot days ($\geq 95^{\circ}$ F) has recently increased in the West, but declined in the remainder of the country, and in the nationwide average, since 1899.

5 INADEQUATE TRACEABLE ACCOUNTS

NCA3 began the practice of including a section on Traceable Accounts at the end of each chapter. This is the description from NCA5:

Each chapter concludes with a section entitled Traceable Accounts, which provides information on the overall process used to develop the chapter as well as a separate Traceable Account section for each Key Message. These Traceable Accounts describe the supporting evidence behind each Key Message, the process and rationale authors used in reaching their conclusions, and the author team’s expert assessment of the confidence in and, where applicable, likelihood of these conclusions. As such, Traceable Accounts provide information about the state of the science, document sources of uncertainty, identify research gaps, and allow traceability to data and resources.

The Traceable Accounts for each Key Message includes the following sections:

- Description of Evidence Base

- Major Uncertainties and Research Gaps
- Description of Confidence and Likelihood

The Traceable Accounts could have addressed the statutory requirement that the NCA discuss scientific uncertainties. They could also help satisfy the Executive Order for Restoring Gold Standard Science by addressing reproducibility, transparency, communication of error and uncertainty, and skepticism of findings and assumptions. However, the Traceable Accounts in NCA5's Chapters 2 and 3 are uneven, often superficial and uninformative.

An example of a deficient Traceable Account is NCA5's Key Message 2.2: "Extreme Events are Becoming More Frequent and Severe". The "Description of the Evidence Base" offers two paragraphs, the first summarizing the findings and the second generically describing the types of data that can be used for storms. "Major Uncertainties and Research Gaps" focuses on the uneven distribution across U.S. populations due to differences in exposure and vulnerability. The only limitations in data referred to are the lack of a long-term record of lightning and the lack of homogenized daily and hourly temperature data apart from reanalysis products.

But the most astonishing deficiency is in the "Description of Confidence and Likelihood." One of the statements in Key Message 2.2 is:

Hurricanes have been intensifying more rapidly since the 1980s (*high confidence*) and causing heavier rainfall and higher storm surges (*high confidence*).

The Description of Confidence and Likelihood includes the following statement:

Basic physical understanding and climate models both provide robust explanations for the links between climate change and observed changes in these extremes: this is why the authors also have high confidence that storms are delivering more rainfall and high confidence that storm surges are becoming higher.

No reference to journal publications or a specific data set is made anywhere in the chapter regarding hurricane-induced precipitation or storm surge to support this *high confidence* conclusion.

The Traceable Accounts for the various Key Findings of Chapter 3 have some substantial inconsistencies. Consider

Key Message 3-1: Human activities have caused global warming.
Confidence and Likelihood: "There is *very high confidence* that emissions of GHGs from human activities, fossil fuel use in particular, have *unequivocally* caused all global warming observed over the industrial era. There is also *very high confidence* that changes in natural climate drivers have had globally small and regionally variable long-term effects over this period."

The Uncertainties section addresses only uncertainties in radiative forcing by greenhouse gases, ignoring the great uncertainty in model response (CWG25 Chapters 4&5) In contrast, the Traceability Analysis of Key Message 3-4 ("Humans Are Changing Earth System Processes") raises the problem that relatively little is known about the role of natural variability at the regional scale. From its Description of Evidence Base:

In the last decade, there has been increasing recognition of the role of regional-scale natural variability in the past and future evolution of climate, as well as the fact that regional natural variability signals can be competitive with forced anthropogenic signals.

And from its Major Uncertainties and Research Gaps:

For example, the true natural variability of the climate system on decadal to centennial timescales could be larger than what we estimate from GCMs, which would further increase the irreducible uncertainty stemming from natural variability.

These statements are among the most scientifically credible in the entire NCA5 and they directly undercut the confidence assigned to Key Message 3-1.

There is also a generic problem in tracing NCA5's Traceable Accounts. The Front Matter of NCA5 states:

Each figure and some tables are accompanied by a metadata survey, which can be accessed in the online version of the report by clicking on the eyeball icon above the figure or table (see the table below for explanations of additional icons used throughout the report). The metadata survey describes data sources, figure or table development methods, copyright information, and other important documentation.

Clicking on the 'eyeball' icon does not reveal the metadata information; instead a "Contact Us" email appears, addressed to the USGCRP. This is an inadequate way to archive the metadata information.

In summary, the Traceable Analyses examined here, with few exceptions, fail to meet the stated objectives of the NCA.

6 OVEREMPHASIS ON EXTREME SCENARIOS

To respond to the GCRA mandate to "project major trends for the subsequent 25 to 100 years," NCA5 relies on projections of future climate change based on estimates of future emissions (the so-called SSP scenarios) and/or specified levels of radiative forcing (the RCP scenarios). For the most part, NCA5 uses scenarios developed for the IPCC Assessment Reports.

In CWG Section 3.2.1 we presented a critical assessment of past and current IPCC scenario envelopes, showing that they have tended to overshoot observed emissions. IPCC AR6 published in 2021 stated that "The likelihood of high-emission scenarios like RCP8.5 and SSP5-8.5 is considered low in light of recent energy sector developments." and that same year UN Climate Negotiators (COP) stopped using scenarios above 4.5.

NCA5 makes the following observations about the scenarios they use:

NCA5 authors were advised to assess the full range of scenarios available. (Front Matter)

Scenarios of future emissions and land-use change are developed as plausible alternatives, but no relative likelihood is attached to them. Some recent studies, however, have argued that

the highest scenario, SSP5-8.5, is no longer plausible without a reversal of current trends in the adoption of renewables and energy efficiency. (Key Message 3.3 Traceable Analysis)

Despite acknowledging the issue, NCA5 otherwise ignored the problem. A keyword search of NCA5 of mentions of RCP/SSP scenarios finds over 50% referred to the extreme emissions scenarios RCP8.5/SSP5-8.5. Further, nine NCA5 figures (5.2, 6.9, 7.4, 11.3, 14.6, 22.13, 22.19, 27.9, 29.13) included *only* the 8.5 scenarios.

The problem of relying on extreme scenarios is particularly acute for NCA5's sea level rise projections. The sea level rise scenarios used in NCA5 do not relate directly to the SSP emissions scenarios, but rather to specified amounts of sea level rise by 2100. Table 4 of NCA5's "Guide to Using this Report" reproduced below shows that Low relates to 0.3 m (0.98 feet), Intermediate-Low relates to 0.5 m (1.64 feet), Intermediate relates to 1.0 m (3.28 feet), Intermediate-High relates to 1.5 m (4.92 feet), and High relates to 2.0 m (6.56 feet).

Table 4. Descriptive Terms for Common Sea Level Rise Scenarios Used in NCA5						
Future global mean sea level rise and sea level rise along United States coastline are shown for five scenarios in feet (and meters), relative to a 2000 baseline. The US values shown in the right half of the table are averaged across the US coastal regions, including the contiguous US, Alaska, Hawaii, and the US-Affiliated Pacific Islands, and the US Caribbean. The national values shown in the table differ substantially from regional values. For example, sea level rise is higher in the Gulf Coast and lower or even negative in some parts of Alaska. In the next 30 years (2020–2050), sea level along the contiguous US coasts is expected to rise 0.92 feet (0.28 m), the same amount of sea level rise observed over the last 100 years (1920–2020). See Chapter 9 for regional sea level information. Adapted from Sweet et al. 2022.”						
Sea Level Rise Scenario Descriptor	Global Mean Sea Level			United States		
Year	2050	2100	2150	2050	2100	2150
Low	0.49 (0.15)	0.98 (0.3)	1.31 (0.4)	0.59 (0.18)	0.98 (0.3)	1.64 (0.5)
Intermediate-Low	0.66 (0.20)	1.64 (0.5)	2.62 (0.8)	0.75 (0.23)	1.64 (0.5)	2.95 (0.9)
Intermediate	0.92 (0.28)	3.28 (1.0)	6.23 (1.9)	0.89 (0.27)	3.28 (1.0)	6.89 (2.1)
Intermediate-High	1.21 (0.37)	4.92 (1.5)	8.86 (2.7)	1.12 (0.34)	4.92 (1.5)	8.86 (2.7)
High	1.41 (0.43)	6.56 (2.0)	12.14 (3.7)	1.38 (0.42)	6.56 (2.0)	12.46 (3.8)

TABLE 1: Reproduction of Table 4 from the Front Matter of NCA5

Considering only projections representing processes quantified with at least medium confidence, IPCC AR6 assessed global mean sea level projections for all emissions scenarios (including implausible extreme emissions scenarios) for 2100 to fall between 0.2 and 1.6 m (5th—95th percentile range). Nevertheless, the range of sea level rise scenarios considered by NCA5 extends to 2.0 m. Further, NCA5 fails to include plausible lower scenarios, such as 0.2 m cited by the IPCC, which also corresponds to the average rate of sea level rise over the past century.

Chapter 9 of NCA5 offers the following predictions of sea level rise:

“Under a range of potential global warming levels, average sea level along US coastlines is *likely* to be between 12 and 20 inches above 2000 sea levels in 2050 (Figure 9.2).”

The projected sea level rise values span from the Intermediate scenario to values slightly greater than the High scenario. They can be put into context by considering the observed global sea level rise from 2000-2024, which is only approximately 3.4 inches (0.28 feet). **NCA5 is thus projecting a sea level rise during 2025 and 2050 that is 3.5 to 5.8 times the rate observed during 2000-2024, an exceedingly implausible outcome.** Chapter 9 further states:

“Looking toward the future, an 11 inch (28 cm) average rise along the contiguous US coastline is expected by 2050 (relative to 2020), with a *likely* range of 9-13 inches.”

...

“There is *high confidence* and it is *likely* that sea levels will rise about 11 inches (likely range of about 9–13 inches) between 2020 and 2050”

In summary, NCA5 presents future sea level rise scenarios that are not directly relatable to either SSP/RCP emissions scenarios or to plausible global warming levels. The NCA5 scenarios include two that are higher than the 95th percentile for the extreme emissions scenario RCP8.5 and include hypothesized ice sheet processes in which there is low confidence. Projections to 2050 require rates of sea level rise from 2025 to 2050 that are more than triple the rate observed during 2000-2024. Further, NCA5 ascribes *high confidence* to these sea level rise projections. CWG25 Section 7.3 provides further context for USGCRP projections of extraordinary sea level rise in the next 25 years.

More generally, NCA5’s inclusion of, and even focus on, implausible extreme scenarios supports projections of alarming impacts that are very unlikely to be driven by anthropogenic warming.

7 CONFLATION OF NATURAL AND ANTHROPOGENIC CLIMATE CHANGES

NCA5’s Glossary defines “climate change” as a change in climate for any reason. This is technically correct, but at variance with the (now) popular perception of all climate changes being anthropogenic. The distinction is not academic- it is central both to attribution questions and to projections, since anthropogenic changes would respond to mitigation, while natural changes are unavoidable.

The discussion on NCA5 page 2-4 begins “Human activities are changing the climate”, then goes on to list evidence for anthropogenic GHG emissions and climate warming. The next paragraph begins “Climate change is happening now in the United States. Including Alaska, the continental US has been warming about 60% faster than the planet as a whole since 1970.” In context the implication is that the warming of CONUS faster than the planetary average is an effect of GHG emissions. But it is an artifact of the Earth’s surface being 70 percent ocean. Since the ocean warms more slowly than land, comparison of any land surface outside the tropics with the global average would show the land surface to be warming faster than the global average, regardless of the cause of the warming.

The same page goes on to say “Climate change is already affecting people in the United States. Extreme heat was estimated to be responsible for more than 700 deaths per year between 2004 and 2018 although some estimates put heat-related mortality closer to 1,300 deaths annually.” But extreme heat, whether natural or not, has always been a risk to human health. As to trends, the data show that adaptation has caused the mortality burden of extreme heat to drop substantially in recent decades (CWG25 Section 10.3.1).

Consider also the NCA5 statement on precipitation that we cited in Section 3.2:

Many eastern regions of the country are getting wetter (Figure 2.4). Average annual precipitation from 2002-2021 was 5%-15% higher relative to the 1901-1960 average in the central and eastern US, a trend attributable to climate change.

The attribution is superfluous because the statement is a tautology by NCA5's definition of climate change. But its gratuitous inclusion effectively attributes the observed trend to greenhouse gas emissions.

Exploitation of the ambiguity in "climate change" is endemic to NCA5. The NASEM review (pp. 23-24) commented on this problem as follows:

The draft NCA5 report frequently describes "change" but does not consistently distinguish between natural variability and climate change. This lack of clarity can be misleading for the reader by creating the misconception that some impacts that are the result of the combination of natural variability and climate change are instead solely due to climate change... it is important for NCA5 authors to accurately identify the multiple drivers of change (including global change), in addition to climate change, and accurately attribute impacts. Clearly describing impacts due to natural variability, climate change, or a combination of the two is important not only for informing policy decisions, but also for building public understand and trust in the report's messages. Climate change and natural variability should be clearly defined in the glossary and natural variability could be introduced in chapter introductions when the concept is important for understanding key messages.

This could have been easily addressed by reserving the term "climate change" for anthropogenic changes and using "climate variability" for natural changes. But the NCA5 authors seemingly ignored their recommendation.

A separate but related issue is use of the term "climate change" to refer to increases in socioeconomic vulnerabilities that have anthropogenic but non-GHG-related causes. NCA5 page 1-19 states "Climate change exacerbates inequalities." But the issue discussed is the negative correlation between neighborhood-level income and land surface temperatures (LST) in urban areas. To the extent the correlation is real the most likely explanation is that low-income households concentrate where housing is less expensive, which may be in neighborhoods with less nearby parkland and tree cover, which has nothing to do with CO₂ emissions. Similarly the NCA5 makes repeated reference to the Billion Dollar Disaster dataset which is dominated by trends in population and economic growth, as we discussed in CWG25 Section 10.2.

8 SUMMARY EVALUATION OF NCA5 AND RECOMMENDATIONS FOR NCA6

This chapter summarizes our evaluation of NCA5 relative to OSTP's charge: whether NCA5 (1) complied with the USGCRP's statutory obligation; (2) adheres to the core tenets of Gold Standard Science; and (3) can suitably inform federal climate and energy policy. The points made in this summary analysis refer to specific sections of this report as well as to the CWG25 report.

8.1 Compliance with USGCRP's statutory obligation

As cited in Section 1.2 of this report, the Global Change Research Act requires an assessment that:

- (1) integrates, evaluates, and interprets the findings of the [Global Change Research] Program and discusses the scientific uncertainties associated with such findings;
- (2) analyzes the effects of global change on the natural environment, agriculture, energy production and use, land and water resources, transportation, human health and welfare, human social systems, and biological diversity; and
- (3) analyzes current trends in global change, both human-induced and natural, and projects major trends for the subsequent 25 to 100 years.

We deem NCA5 to have fallen short when evaluated against these criteria because of the following major shortcomings (each is detailed in the chapters cited):

- Inadequate assessment of the relevant topics in climate science and the hazard components of climate impacts (such as hurricanes and floods). NCA5 instead focuses on evaluation of societal vulnerabilities (such as flood infrastructure, access to air conditioning, etc.), “Climate Action,” and policy recommendations with a social justice emphasis (Section 1.2).
- Overconfidence in conclusions and inadequate assessment of uncertainties (Section 3.1.5)
- Inadequate discrimination between natural and human-induced causes of trends and impacts (Chapter 7)
- Inadequate analysis of climate trends by relying on short time periods (Chapter 6)
- Uncritical evaluation of emission scenarios and climate simulations in making projections for the next 25 to 100 years (Chapter 6; Section 3.2.1 and Chapter 5 of CWG25).

8.2 Adherence to the tenets of Gold Standard Science

With respect to the tenets of Gold Standard science listed in Section 1.2 of this report, we have identified the following major shortcomings in NCA5:

- **Lack of reproducibility.** In the few examples that we examined closely, we identified an instance where we could not reproduce the findings of the NCA5, a task that should require only rudimentary data analysis (Section 4.1)
- **Lack of transparency.** In many instances, the Traceable Account fails to justify the Key Finding. Metadata was not readily accessible. (Chapter 5)
- **Inadequate communication of error and uncertainty.** The Traceable Account for each Key Message provides a section on Major Uncertainties and Research Gaps; most of the accounts that we examined were superficial and uninformative. (Chapter 5)
- **Not skeptical of findings and assumptions.** The main text of the chapters is, with few exceptions, not skeptical of its findings and assumptions. (Chapters 3, 5)
- **Some hypotheses and Key Messages are not falsifiable.** Key Message 13.4 is but one of many examples: “Equitable Distribution of Transportation Trade-offs and Benefits Requires Community Involvement.”
- **Failure to acknowledge benefits** of warming and additional atmospheric CO₂ (CWG25 Section 2.1, Chapter 9)
- **Inadequacy of the peer review process.** The review of the NCA5 Third Order Draft by National Academies of Science, Engineering and Medicine (NASEM) identified some of the deficiencies we have noted, but the NCA5 authors were not obligated to adopt the NASEM recommendations. More importantly, we’ve identified many other deficiencies in NCA5; the NASEM review did not point them out and thus lacked the rigor expected of NASEM reports. (Section 1.4, Chapter 7)

8.3 Suitability for informing Federal climate and energy policy

We assess that the selection and presentation of information in NCA5 makes it an unreliable basis for U.S. climate and energy policymaking. For each of the elements of knowledge characterization suitable for risk management listed in Section 1.3, here are specific examples where NCA5 falls short:

- The degree to which the assumptions made are reasonable/realistic based on background knowledge
 - NCA5 focused on implausibly extreme scenarios for emissions and sea level rise scenarios and made policy recommendations without adequate consideration of the costs
- The degree to which data/information exists and are reliable and relevant
 - NCA5's assumptions about model reliability, and its use of short time periods for analysis, does not support its high-confidence conclusions
- The degree to which there is disagreement among experts, including those from different environments or academic fields
 - While NCA5 was an interdisciplinary effort, the concerns we have raised all point to a failure to canvass or communicate the full range of evidence including data and research findings that speak against Key Messages.
- The degree to which the phenomena involved are understood and accurate models exist
 - NCA5 is not appropriately skeptical of the ability of climate models to attribute the causes of warming and extreme events or to project future climates
- The degree to which the knowledge has been thoroughly examined with respect to unknown knowns
 - A major NCA5 shortcoming is its dismissal of natural climate variability in many contexts where it should have been a major consideration

NCA5 is also problematic for informing policy because its hyperbolic language promotes alarm and advocates for specific policies without proper analysis of marginal costs and benefits. In recent decades, climate science has engendered passion and emotion. Indeed, some younger climate scientists have gone into the field to "save the planet." The NASEM Review of the NCA5 underscores the endemic nature of this problem - their review appears to be written by authors who share many of the biases of the NCA5 team, with an inordinate focus on social and policy issues. It is then not surprising that objectivity can suffer when climate scientists communicate their understanding to non-experts. Further, the field of climate science has been politicized by NetZero mandates. This lack of objectivity usurps the public's right to make informed choices and undermines their confidence in the scientific enterprise.

Philip Handler, President of the National Academy of Sciences, identified the problem forty-five years ago in an [editorial](#) that still resonates:

Difficulty arises in the scientific community from confusion of the role of scientist *qua* scientist with that of scientist as citizen, confusion of the ethical code of the scientist with the obligation of the citizen, blurring the distinction between intrinsically scientific and intrinsically political questions. When scientists fail to recognize these boundaries, their own ideological beliefs, usually unspoken, easily becloud seemingly scientific debate.

8.4 Recommendations for NCA6

While we have assessed NCA5 to be scientifically inadequate and unsuitable as a basis for decision-making, one can imagine a different National Climate Assessment that meets statutory and policymaker requirements. The goal should be not to persuade, but to impart understanding by being objective, transparent and complete. A suitable NCA would be an important complement to IPCC assessments to the extent that its analysis and conclusions differed from the oracular “consensus” of the IPCC reports and additional topics are addressed. Such a report would promote a more open scientific discussion and a more informed policy debate.

We recommend that the NCA6 have a narrower scope than NCA5, refocusing on climate science, including natural climate variability, with topics to include:

- How the climate has changed in the past. The U.S. has the best and longest meteorological records and they should be examined and displayed in their entirety. Trend analyses should include longest periods available and employ suitable statistical methods. Local paleoclimate data should be considered where available. Analysis should demonstrate not only how the climate has varied and changed, but also regional heterogeneity. Standard, easily understandable metrics should be employed.
- The range of climatic and weather conditions, including extremes, that have occurred in the past and can reasonably be expected to occur in U.S. regions over the near future.
- A critical survey of current global and regional modeling capabilities and their suitability for regional impact assessments and projections. This would include an assessment of their ability to reproduce observed changes. It should also discuss alternative methods for generating projections and forecasts on regional and decadal scales, along the lines of Chapter 10 of the IPCC AR6 WGI Report.
- Assessment of the impacts of recent historical climate variability and change on ecosystems and society. Where long-term data indicate a detectable regional change, regional projections can be made for the period 2030-2050. Emissions scenarios exceeding RCP4.5/SSP2-4.5 and ECS values exceeding 4°C should not be used. Regional projections to 2100 have no credibility at this time.

It is important that climate projections be assessed in context of current science capabilities. The pressures toward spatially or temporally precise projections should be resisted as the current models are nowhere up to the task (an example is discussed in CWG25 Section 5.8). In that light, the many pages of NCA5 devoted to projected regional and sectoral impacts have little value.

Given the substantial deficiencies in NCA5 that we have identified, along with the endemic bias of the climate community towards alarmism and policy advocacy, producing an objective NCA6 report with the current NCA organizational construct would be a challenge. A NASEM review could help, but only if the selected participants reflect full viewpoint diversity and do not simply mirror the assumptions and biases of past NCA teams. We regard the CWG25 report to be a good start on NCA6 in terms of scope and objectivity, including adherence to the tenets of Gold Standard Science.

More broadly, we recommend a rewrite of the Global Change Research Act. The current mandate is vague and off point, allowing successive NCA teams to interpret the mandate as they see fit. Moreover, it is becoming increasingly clear that current and foreseeable models are not fit-for-purpose to describe regional climates, let alone usefully project them for 25 and 100 years. Further, the use of the term “global change” is ambiguous and outdated.

9 REFERENCES

- Aven, Terje (2017) "Improving Risk Characterisations in Practical Situations by Highlighting Knowledge Aspects, with Applications to Risk Matrices," *Reliability Engineering & System Safety* 167 (November 2017): 42–48
<https://www.sciencedirect.com/science/article/pii/S0951832016306950>
- Christy, John (2025) Precipitation Data Archive for Midwest Stations.
https://www.nsstc.uah.edu/data/ushcn_jrc/US_MidW_61stn_dly_ppt_1893_2024_250704.xlsx
- Climate Working Group (2025) Impacts of Carbon Dioxide Emissions on the U.S. Climate. Washington DC: Department of Energy, May 27, 2025.
- Dellawalla, Colette, Victor Ambrose, Carl Bergstrom *et al.* (2025) "Trump's new 'gold standard' rule will destroy American science as we know it." *The Guardian* May 29, 2025
<https://www.theguardian.com/commentisfree/2025/may/29/trump-american-science>
- Diffenbaugh, N.S., D. Singh, and J.S. Mankin, 2018: Unprecedented climate events: Historical changes, aspirational targets, and national commitments. *Science Advances*, 4 (2), 3354.
<https://doi.org/10.1126/sciadv.aao3354>
- Handler, Philip (1980) Public Doubts About Science. *Science* 208(4448) p. 1093
<https://www.science.org/doi/10.1126/science.208.4448.1093>
- Kirchmeier-Young, M.C. and X. Zhang, 2020: Human influence has intensified extreme precipitation in North America. *Proceedings of the National Academy of Sciences of the United States of America*, 117 (24), 13308–13313. <https://doi.org/10.1073/pnas.1921628117>
- Knutson, Thomas and Fanrong Zeng (2018) "Model Assessment of Observed Precipitation Trends over Land Regions: Detectable Human Influences and Possible Low Bias in Model Trends" *Journal of Climate* 31(12) <https://journals.ametsoc.org/view/journals/clim/31/12/jcli-d-17-0672.1.xml>
- Koonin, Steven (2024) *Unsettled: What Climate Science Tells Us, What It Doesn't, and Why It Matters*. New York: Basic Books.
- McKittrick, R., and J.R. Christy (2019). Assessing changes in US regional precipitation on multiple time scales. *Journal of Hydrology*, 578, 124074. <https://doi.org/10.1016/j.jhydrol.2019.124074>
- McKittrick, R., and J.R. Christy (2025a) "Data and Code for DoE Report 2025", Mendeley Data, V1, doi: 10.17632/8n9ks93vn7.1
- McKittrick, R., and J.R. Christy (2025b) "Supplement to NCA5 Critique: Examining the reproducibility of NCA5 claim of extreme precipitation trends." Available online at [\[insert URL\]](#)
- National Academies of Sciences, Engineering, and Medicine. 2023. *Review of the Draft Fifth National Climate Assessment*. Washington, DC: The National Academies Press.
<https://doi.org/10.17226/26757>.
- NCA4. (2017). *Climate science special report: Fourth National Climate Assessment, Volume I* (D. J. Wuebbles, D. W. Fahey, and K. A. Hibbard, Eds.). U.S. Global Change Research Program.
<https://doi.org/10.7930/J0J964J6>
- NCA5. (2023). *Fifth National Climate Assessment* (A. R. Crimmins *et al.*, Eds.). U.S. Global Change Research Program. <https://doi.org/10.7930/NCA5.2023.CH1>
- Pielke Jr., R. (2024). Scientific integrity and U.S. 'billion dollar disasters'. *npj Natural Hazards*, 1, Article 12. <https://www.nature.com/articles/s44304-024-00011-0>
- Pielke Jr., R. (2025, May 8). RIP NOAA's billion dollar disasters. The Honest Broker [Substack].
<https://rogerpielkejr.substack.com/p/rip-noaas-billion-dollar-disasters>

Schlund, Manuel, Axel Lauer, Pierre Gentine, Steven C. Sherwood, and Veronika Eyring (2020)
Emergent constraints on equilibrium climate sensitivity in CMIP5: do they hold for CMIP6?
Earth System Dynamics 11 1233-1258 <https://doi.org/10.5194/esd-11-1233-2020>

APPENDICES

APPENDIX 1: Charge to the review

ABOUT THE AUTHORS

[Insert updated bios]

DOCUMENT WITHHELD FOR PRIVILEGE

DOCUMENT WITHHELD FOR PRIVILEGE


From: Judith Curry [curry.judith@gmail.com]
Sent: 5/23/2025 6:15:55 PM
To: Travis Fisher [travis.scott.fisher@gmail.com]; Ross McKittrick [ross.mckittrick@gmail.com]; John Christy [climatemanager60@gmail.com]; Steven Koonin [steven.koonin@gmail.com]; Roy Spencer [roywspencer@hotmail.com]
Subject: problem with chapter 4
Attachments: 4 ClimateSensitivityMay21 Curry.dotx

See the attached version of chapter 4 that we discussed on Wed. The changes in red did not make it into the final version, we agreed on these changes in the phone call and they should be added to the text,

There are also some changes to ch 9 on attribution that were discussed on wed, that didn't make it into the final version (i'm not too concerned about these)

My bigger concern is that if the two chapters that i focused on were not the latest versions, are there others?

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

4 Climate Sensitivity to CO₂ Forcing

Chapter Summary

There is growing recognition that climate models are not fit for the purpose of determining the Equilibrium Climate Sensitivity (ECS) of the climate to increasing CO₂. The IPCC has turned to data-driven approaches using historical data and paleoclimate reconstructions, but their reliability is diminished by data inadequacies.

Data-driven ECS estimates tend to be lower than climate model-generated values. The IPCC AR6 upper bound for the likely range of ECS is 4.0°C, lower than the AR5 value of 4.5°C. This lowering of the upper bound seems well justified by paleoclimatic data. The IPCC AR6 lower bound for the likely range of ECS is 2.5°C, substantially higher than the AR5 value of 1.5°C. This raising of the lower bound is more weakly justified. Evidence since the AR6 finds the lower bound of the likely range to be around 1.8°C.

4.1 Introduction

The climate's response to increasing concentrations of CO₂ is central to the scientific debate on anthropogenic climate change, and so also to the public debate on "climate action." The simplest measure of that response is the rise in the global average surface temperature, quantified by the Equilibrium Climate Sensitivity (ECS). ECS is defined as the amount of warming expected in response to a doubling of CO₂ from its pre-industrial concentration of 280 ppm, after all climate components have had time to adjust. Some components, like temperatures in the lower atmosphere (troposphere), adjust rapidly, while others such as the deep ocean and cryosphere might take as long as centuries. A related measure, the Transient Climate Response (TCR), better describes the shorter time scales; it is defined as the amount of warming when the CO₂ concentration is doubled by rising one percent annually for 70 years.

The 1979 Charney Report for the US National Academy of Sciences (Charney *et al.* 1979) proposed that the most likely ECS was $3.0 \pm 1.5^\circ\text{C}$. The IPCC repeatedly reaffirmed that range with only minor variations until its most recent AR6. AR5 termed 1.5 -- 4.5°C as the *likely* range (66 percent probability) and stated that ECS is *extremely unlikely* (95 percent probability) to be below 1.0°C and very unlikely (90 percent probability) to exceed 6.5°C.

The uncertainty in ECS has remained stubbornly wide, despite many individual studies that claimed to narrow it. Most recently, AR6 narrowed the *likely* range to 2.5–4.0°C and deemed the *very likely* range to be 2.0 - 5.0°C. This narrowing on the low end is disputed, as will be discussed below.

Uncertainties in ECS are highly consequential for policy making. As will be discussed in Chapter 11, economic models use ECS values to project the costs of CO₂ emissions. The traditional value (3.0 °C) has typically yielded modest global social costs of CO₂ emissions, sufficient to justify some policy actions, but mostly deferred to later in this century. If ECS is very high (above 4.5°C) immediate aggressive emission controls become more imperative, whereas no CO₂ emission controls are economically justifiable for ECS below 2.0°C (Dayaratna et al. 2016, 2020). Obtaining a precise estimate is impossible, so policy making needs to account for the uncertainty.

Climate scientists use multiple lines of evidence to determine the Equilibrium Climate Sensitivity:

- Climate model simulations
- Historical observations

- Paleoclimatic observations
- Process understanding of feedbacks

By itself, the equilibrium warming effect of a doubling of atmospheric CO₂ is slightly more than 1°C.³³ The large values of ECS arise from positive feedbacks that amplify the warming effect of CO₂. Water vapor feedback is positive: a warmer atmosphere may have more water vapor, which itself is a powerful greenhouse gas. Water vapor feedback is partially canceled by the fact that atmospheric temperature decreases more slowly with height above the Earth's surface when atmospheric water vapor increases. Warmer temperatures also result in less snow and sea ice cover, allowing the Earth to absorb more of the sun's radiation. Some simple estimates of these three feedbacks increase the ECS to around 2°C (Sherwood et al., 2020).

Higher values of ECS arise primarily from positive cloud feedbacks. The magnitude, and even the sign, of cloud feedbacks is the greatest source of uncertainty. Elements of cloud feedback include changes in the latitudinal distribution of clouds, changes in the distribution of cloud height (changes in low versus high clouds), changes to the phase of clouds (ice versus liquid), changes in cloud particle size (associated with changes in concentration and/or composition of aerosol particles), changes in the precipitation efficiency of clouds, and even changes in how clouds are distributed over the daily solar cycle (Curry and Webster, 1999). It is difficult for climate models to simulate any of these cloud processes correctly owing to their small scale, let alone predict how they will change in the future. Further, cloud processes modulate the magnitudes of the water vapor, lapse rate, and the surface albedo feedbacks.

4.2 Model-based estimates of climate sensitivity

The ECS ranges given in IPCC AR4 and AR5 were obtained primarily by examining the behaviour of large-scale climate models, also called General Circulation Models (GCMs). However, the IPCC changed course in its AR6 when it turned to a more direct data-driven methodology. Here we discuss some of the pitfalls of using GCMs to try to determine the Earth's climate sensitivity.

ECS is an emergent property of climate models—that is, it is not directly parameterized or tuned. Historically, collections of parameterizations have been chosen to produce an expected value of climate sensitivity, particularly with regards to sub-grid-scale cloud processes, particularly in AR4 and to a lesser extent in AR5. Otherwise plausible GCMs and parameter selections have been discarded because of perceived conflict with this warming rate, or aversion to a model's climate sensitivity being outside an accepted range. For example, even in a CMIP6 model, Mauritsen and Roeckner (2020) state the following regarding their Max Planck Institute climate model (emphasis added):

“We have documented how we tuned the MPI-ESM1.2 global climate model to match the instrumental record of warming; an endeavor which has clearly been successful. Due to the historical order of events, the choice was to do this practically by *targeting an ECS of about 3 K using cloud feedbacks*, as opposed to tuning the aerosol forcing.”

In other words, the modelers chose an ECS value of 3°C and then tuned the cloud parameterizations to match their intended result.

Direct warming from CO₂ doubling is only about 1°C (Hall and Manabe 1997); further warming arises from climate feedbacks that are not explicitly resolved by the GCM but rely on parameterizations of physical processes. Higher values of ECS arise primarily from assumed positive cloud feedbacks,

whereas the magnitude and even the sign of the feedbacks are very uncertain. Elements of cloud feedback include changes in the latitudinal distribution of clouds, changes in the distribution of cloud height (changes in low versus high clouds), changes to the phase of clouds (ice versus liquid), changes in cloud particle size (associated with changes in concentration and/or composition of aerosol particles), changes in the precipitation efficiency of clouds, and even changes in how clouds are distributed over the daily solar cycle (Curry and Webster, 1999). It is difficult for GCMs to simulate any of these processes correctly owing to their small scale, let alone predict how they will change in the future. Further, cloud processes modulate the magnitudes of the water vapor, lapse rate, and the surface albedo feedbacks.

ECS can be determined from climate model simulations by doubling the concentration of CO₂ and allowing several centuries for the warming to equilibrate. To avoid the necessity of long simulations, “effective climate sensitivity” is commonly evaluated from a 150-year simulation in response to a sudden quadrupling of CO₂.

The spread of ECS values from the CMIP5 ensemble of climate models used in the AR5 was 2.0–4.7°C; that range increased for the CMIP6 models used in the AR6 to between 1.8 and 5.7°C (Chen et al, 2021, Scaffeta 2021, see Figure 5.1). Far from settling the model-based sensitivity of climate system uncertainties appear to be growing. The main cause of the overall upward shift in ECS in CMIP6 relative to CMIP5 is a larger positive cloud feedback, driven by changes to the cloud parameterizations in many CMIP6 models (Zelinka et al, 2020)




Figure 4.1 Equilibrium Climate Sensitivities in °C of 37 climate models from the CMIP6 ensemble. Identifiers for the various models appear along the horizontal axis. From (Scafetta, 2021) <https://www.mdpi.com/2225-1154/9/11/161>

Because of concerns about model tuning and the high sensitivity to cloud parameterizations, the IPCC AR6 (2021) did not rely on climate model simulations in their assessment of climate sensitivity, relying instead on other methods, chiefly a recent empirical estimate.

4.3 Data-driven estimates of climate sensitivity

Climate sensitivity can also be estimated from instrumental records of surface temperatures and ocean heat content, combined with estimates of how climate forcings (e.g., greenhouse gases, solar, volcanoes, aerosols) have changed in the past (Otto et al., 2013). Using this information, a simple empirical model called an Energy Balance Model can be employed. It requires estimating a feedback parameter whose uncertainties are highly amplified in the resulting ECS (Roe and Baker, 2007).

The accuracy of the data-driven methods depends on the quality of the input data. Assumptions are needed about ocean heat storage, and the ocean data is inadequate. The greatest source of uncertainty is the amount and composition of aerosol particles and their interactions with cloud radiative properties (the so-called aerosol indirect effect). Climate models exhibit warming in response to GHGs but cooling in response to aerosols (Schwartz et al. 2007). Observed 20th century warming can be shown to be consistent either with low ECS and low aerosol cooling, or high ECS and high aerosol cooling. Since fossil fuel use adds both GHGs and aerosols to the atmosphere, both effects need to be estimated to isolate the warming effect of CO₂. Over time, the IPCC's estimate of the cooling effect of aerosols has declined (Lewis 2022).

Paleoclimate proxies are also used to evaluate the sensitivity of past climates, by comparing paleoclimate changes in the Earth's temperatures to estimates of changes in forcings. The two most informative periods are the last glacial maximum (around 20,000 years ago), which was about 3–7°C colder than today, and a mid-Pliocene period (roughly three million years ago), which was 1–3°C warmer than today. The limits on cooling during the last glacial maximum give the best single evidence that high values of climate sensitivity are unlikely. However, paleoclimate estimates are associated with very large uncertainties in the estimated temperatures and forcings. Further, estimates of climate sensitivity based on past climate states may not be representative of the current state of the climate system.

A recurring theme in the climate literature is that ECS estimates based on historical data are smaller than ECS estimates inferred from climate models (Sherwood and Forest 2024). About 15 estimates based on historical data appeared in the peer-reviewed literature between 2012 and 2024 yielding ECS best estimates between 1.0°C and 2.5°C, although critics have questioned some of the methods and the data quality. For the AR6, the IPCC placed primary weight on the results of Sherwood et al. (2020) that combined historical and paleoclimate proxies with the process-based approach and yielded a best estimate of 3.1°C, with a *likely* range of 2.6–3.9°C. Lewis (2022) raised a number of concerns about this result, including methodological errors, outdated input values, and use of subjective versus objective Bayesian priors in the analysis. Lewis' analysis found that climate sensitivity is estimated to be much lower and better constrained than in the Sherwood et al. analysis – median 2.2°C (1.8–2.7°C in the 17–83 percent *likely* range, and 1.6–3.2°C in the 5–95 percent *very likely* range). The IPCC AR6 estimated only a 5 percent probability that ECS was below 2.3°C, whereas Lewis estimated it to be over 50 percent. The most recent publications on the debate between Sherwood et al. and Lewis further defend their respective positions: Sherwood and Foster (2024) and Lewis (2025).

An argument emphasized in AR6 is that data-driven ECS estimates might understate the future warming response to GHGs because of a so-called “pattern effect” (Forster et al 2021). The tropical Pacific is believed to strongly influence the overall efficiency with which the Earth radiates heat to space, but some regions remove heat more efficiently than others. If the west-to-east temperature gradient in the tropical Pacific is weakened in a warming climate, warming would concentrate where heat is less efficiently removed, raising ECS.

Most climate models simulate that rising GHGs will weaken the east-west temperature gradient, which led the IPCC in the AR6 to conclude that data-driven ECS estimates understated the *future* ECS value. However, Seager et al. (2019) pointed out that, contrary to models, the east-west temperature gradient has been strengthening over time. They further argued that the mechanism predicting otherwise in climate models was based on a faulty characterization of oceanic dynamics and there is no reason to expect the gradient to weaken. A similar argument was recently made by Lee et al. (2024), who concluded that “the trajectory of the observed trend reflects the response to increasing GHG loading in the atmosphere”; in other words, GHG warming should lead to a future strengthening rather than a weakening of the temperature gradient. Increased efficiency of atmospheric cooling implies, if anything, that the future ECS in a warming climate may be *lower* than current estimates.

5.5 Transient Climate Response

The Transient Climate Response (TCR) provides a more useful observational constraint on climate sensitivity. TCR is the amount of warming that results when CO₂ is increased at an annual rate of 1 percent over a period of 70 years (i.e., doubled gradually). Relative to the ECS, observationally determined values of TCR avoid the problems of uncertainties in ocean heat uptake and the fuzzy boundary in defining equilibrium arising from a range of timescales for the longer-term feedback processes (e.g., ice sheets). TCR is more generally related to peak warming and better constrained by historical warming, than ECS. IPCC AR6 judged the *very likely* range of TCR to be 1.2–2.4°C. In contrast to ECS, the upper bound of TCR is more tightly constrained. For comparison, the TCR values determined by Lewis (2023) are 1.25 to 2.0°C, showing much better agreement with the AR6 values than was seen in a comparison of the ECS values.

As will be discussed in Chapter 6, a real-world ECS smaller than the model average is supported by studies that show climate models on average overestimate the observed surface and troposphere warming over the past 50 years. Lower sensitivities also align with evidence from consistent fingerprinting (Section 8), where model responses need to be downscaled to align with observed trends.

Mauritsen, T., & Roeckner, E. (2020). Tuning the MPI-ESM1.2 global climate model to improve the match with instrumental record warming by lowering its climate sensitivity. *Journal of Advances in Modeling Earth Systems* 12, e2019MS002037; <https://agupubs.onlinelibrary.wiley.com/doi/epdf/10.1029/2019MS002037>

Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., et al. (2020). Causes of higher climate sensitivity in CMIP6 models. *Geophysical Research Letters*, 47, e2019GL085782. <https://doi.org/10.1029/2019GL085782>

1. Lan, X., Tans, P. and K.W. Thoning: Trends in globally-averaged CO₂ determined from NOAA

Global Monitoring Laboratory measurements. Version Monday, 14-Apr-2025 09:08:57 MDT
<https://doi.org/10.15138/9N0H-ZH07>.

2. Zeke Hausfather, "Explainer: How Scientists Estimate Climate Sensitivity," *Carbon Brief*, July 22, 2020, <https://www.carbonbrief.org/explainer-how-scientists-estimate-climate-sensitivity>.
3. Jule Charney et al., "Carbon Dioxide and Climate: A Scientific Assessment," 1979, <https://doi.org/10.17226/12181>.
4. Rajendra K. Pachauri and Leo Meyer, *Climate Change 2014: Synthesis Report* (Geneva, CH: Intergovernmental panel on climate change, 2015).
5. "In-Depth Q&A: The IPCC's Sixth Assessment Report on Climate Science," Carbon Brief, August 9, 2021, <https://www.carbonbrief.org/in-depth-qa-the-ipccs-sixth-assessment-report-on-climate-science>.
6. Nicola Scafetta, "Testing the CMIP6 GCM Simulations versus Surface Temperature Records from 1980–1990 to 2011–2021: High ECS Is Not Supported," *Climate* 9, no. 11 (2021): 161, <https://doi.org/10.3390/cli9110161>.
7. Brian J. Soden and Isaac M. Held, "An Assessment of Climate Feedbacks in Coupled Ocean–Atmosphere Models," *Journal of Climate* 19, no. 14 (July 15, 2006): 3354–3360, <https://doi.org/10.1175/jcli3799.1>.
8. S. C. Sherwood et al., "An Assessment of Earth's Climate Sensitivity Using Multiple Lines of Evidence," *Reviews of Geophysics* 58, no. 4 (July 22, 2020), <https://doi.org/10.1029/2019rg000678>.
9. Judith A. Curry and Peter J. Webster, "Thermodynamic Feedbacks in the Climate System," in *Thermodynamics of Atmospheres and Oceans* (New York, NY: Academic Press, 1999), 351–385.
10. Rajendra K. Pachauri and Leo Meyer, *Climate Change*.
11. D. Chen et al., "Framing, Context, and Methods."
12. Mark D. Zelinka et al., "Causes of Higher Climate Sensitivity in CMIP6 Models," *Geophysical Research Letters* 47, no. 1 (2020): e2019GL085782, <https://doi.org/10.1029/2019GL085782>.
13. Thorsten Mauritsen and Johann Jungclaus, "Tuning the MPI-ESM1.2 Global Climate Model to Improve the Match With Instrumental Record Warming by Lowering Its Climate Sensitivity," *Journal of Advances in Modeling Earth Systems* 12, no. 7 (2020): e2019MS002037, <https://doi.org/10.1029/2019MS002037>.
14. Alexander Otto et al., "Energy Budget Constraints on Climate Response," *Nature Geoscience* 6, no. 6 (May 19, 2013): 415–416, <https://doi.org/10.1038/ngeo1836>.
15. Steven C. Sherwood and Chris E. Forest, "Opinion: Can Uncertainty in Climate Sensitivity Be Narrowed Further?" *Atmospheric Chemistry and Physics* 24, no. 5 (2024): 2679–2686, <https://doi.org/10.5194/acp-24-2679-2024>.
16. "Why Low-End 'Climate Sensitivity' Can Now Be Ruled Out," Carbon Brief, July 22, 2020, <https://www.carbonbrief.org/guest-post-why-low-end-climate-sensitivity-can-now-be-ruled-out>.
17. S. C. Sherwood et al., "An Assessment of Earth's Climate Sensitivity."
18. D. Chen et al., "Framing, Context, and Methods."
19. Nicholas Lewis, "Objectively Combining Climate Sensitivity Evidence," *Climate Dynamics* 61, no. 9–10 (2023): 3155–3163, <https://doi.org/10.1007/s00382-022-06398-8>.
20. Ibid.

21. Steven C. Sherwood and Chris E. Forest, "Opinion: Can Uncertainty in Climate Sensitivity Be Narrowed Further?"
 22. Nicholas Lewis, "Comment on 'Can Uncertainty in Climate Sensitivity Be Narrowed Further?' by Sherwood and Forest (2024)," *EGUsphere* [preprint], 2025, <https://doi.org/10.5194/egusphere-2025-1179>.
 23. P. Forster et al., (2021): The Earth's energy budget, climate feedbacks, and climate sensitivity. In: *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V. et al. (eds.)], Cambridge University Press, 2021.
 24. Richard Seager et al., "Strengthening Tropical Pacific Zonal Sea Surface Temperature Gradient Consistent with Rising Greenhouse Gases," *Nature Climate Change* 9, no. 7 (2019): 517–522, <https://doi.org/10.1038/s41558-019-0505-x>.
 25. Sangyeop Lee et al., "Zonal Contrasts of the Tropical Pacific Climate Predicted by a Global Constraint," *Climate Dynamics* 62, no. 1–2 (2024): 229–246, <https://doi.org/10.1007/s00382-023-06741-7>.
 26. D. Chen et al., "Framing, Context, and Methods."
 27. Nicholas Lewis, "Objectively Combining Climate Sensitivity Evidence."
 28. Rajendra K. Pachauri and Leo Meyer, *Climate Change*.
 29. Nicholas Lewis, "Objectively Combining Climate Sensitivity Evidence."
 30. Nicholas Lewis, "Comment on 'Can Uncertainty in Climate Sensitivity Be Narrowed Further?' by Sherwood and Forest (2024)."
- Nierenberg, William et al. (1983) *Changing Climate: Report of the Carbon Dioxide Assessment Committee*. Washington, National Academies Press 1983.

From: Travis Fisher <travis.scott.fisher@gmail.com>
Sent: Monday, May 12, 2025 10:42 PM
To: Ross McKittrick
Cc: Steven Koonin; Judith Curry; John Christy; Roy W. Spencer
Subject: Process for edits & comments
Attachments: 0.Intro.May1 [SEK] TF comment.docx

Hi all,

I know some of us will spend the day tomorrow at the Heritage event or streaming it: <https://www.heritage.org/climate/event/the-sky-falling-reconsidering-the-endangerment-finding>

Very soon (probably later this week), I'll go through each chapter and offer comments regarding readability for non-experts and highlighting anything that seems out of place. What is the best way to do that? I don't seem to have edit/comment rights in Dropbox, which I think is the right approach because I'm sure I would only ruin your work.

Is it ok with you all if I send an email with an attachment once I've completed reading a chapter? My guess is I'll only have minor comments in bubbles to provide killer insights like "please spell out," "this could use more explanation," or "I don't understand this at all." Maybe I'll get bold and ask, "How does this square with [insert inferior analysis]?"

I promised you all that I won't try to change the substance at all. If that's how any of my comments come across, then I've clearly made a mistake, and you should feel free to ignore me or call me out. I intend to keep my promise and make sure you all are 100% comfortable with the report. I'm confident we'll get there, especially since we're keeping the whole effort under Secretary Wright. And the substance is truly impressive, by the way.

Attached is my read of the intro. In case you're reading on a phone or just don't want to open the file, my only comment was on the paragraph about this team not being wrapped up in the EPA process. I wrote:

You could say it's public knowledge that the EPA announced in March 2025 that it was "kicking off a formal reconsideration of the 2009 Endangerment Finding in collaboration with the Office of Management and Budget (OMB) and other relevant agencies. EPA also intends to reconsider all of its prior regulations and actions that rely on the Endangerment Finding."
<https://www.epa.gov/newsreleases/trump-epa-kicks-formal-reconsideration-endangerment-finding-agency-partners>

I think you can acknowledge that this effort is designed to inform the EPA's policymaking process.

If you want to reiterate your independence from EPA, especially EPA's political team, I think that's completely fair. They've been pretty hard for me to get a hold of, so even I have a level of independence that has basically happened by accident!

Best,
Travis

Introduction

This document originated in late March 2025 when Secretary Wright assembled an independent team to write a report on issues in climate science relevant for energy policymaking, with particular focus on the question of whether carbon dioxide emissions endanger the US public. The team agreed to undertake the work on the condition that there would be no editorial oversight by the Secretary, the Department of Energy, or any other government personnel. This condition has been honored throughout the process and the writing team has worked with full independence.

The team began working in early April with a deadline of May 28. The short timeline and the technical nature of the material meant that we could not comprehensively review all topics. While the report is intended to be accessible for non-experts, we have omitted some introductory or explanatory material that can easily be accessed elsewhere. Nor have we attempted to survey entire literatures related to the topics covered. The scientific and technical material herein only samples vast literature on climate change but is, we believe, more than sufficient to establish our conclusions.

The team writing this report was not involved in, nor briefed on, any Administration plans or processes related to a potential challenge to the 2009 EPA CO₂ Endangerment Finding. We were asked, however, to include some information related to the role of US motor vehicles in global climate change, since that is an issue of relevance to the Department of Energy.

The writing team is grateful to Secretary Wright for the opportunity to prepare this report and for his support for independent scientific inquiry.

From: Judith Curry <curry.judith@cfanclimate.com>
Sent: Saturday, May 17, 2025 8:51 AM
To: Ross McKittrick; John Christy; Roy W. Spencer; Steven Koonin; Travis Fisher
Subject: proposed chapter merger

I propose merging the risk management chapter with heat wave mortality chapter. Both chapters are a bit skimpy, and i think they would fit well together

From: Woods, Andrea <andrea.woods@hq.doe.gov>
Sent: Monday, August 4, 2025 10:24 AM
To: Steven Koonin; climateman60@gmail.com; roy.spencer@nsstc.uah.edu; curry.judith@gmail.com; ross.mckitrick@gmail.com
Cc: Barrios, Audrey
Subject: Question from New Yorker - AI use in DOE report

Hi All,

We received an inquiry from the New Yorker over the weekend with claims from unnamed scientists whose work was cited in the DOE report arguing that their work was not cited correctly, and this could indicate that AI was used to compile the report. The claims seem very flimsy and we'll still provide a statement urging these unnamed scientists to submit their comments to the docket, but I wanted to check in with the group. Were any AI tools used?

I don't see any major issues with us utilizing AI tools to complete citations, but if AI was not used at all, we'll note this for the reporter in our response.

Thanks!
Andrea

From: John Christy [climateman60@gmail.com]
Sent: 4/14/2025 10:20:31 PM
To: Roy Spencer [roywspencer@hotmail.com]
CC: Ross McKittrick [ross.mckittrick@gmail.com]; Steve Koonin [steven.koonin@gmail.com]
Subject: Re: [2.2.2 and 2.2.3]Apr14[JC] Figs. Heat/Cold waves
Attachments: CONUS_HtWvs.png; CONUS_CldWvs.png

Roy

See attached.

John C.

On Mon, Apr 14, 2025 at 10:54 AM Roy Spencer <roywspencer@hotmail.com> wrote:

John:

My preference would be to leave out the regional results in the 1st and 3rd plots... graphs are too busy. Just keep the USA48 results.

-Roy

From: John Christy <climateman60@gmail.com>
Sent: Monday, April 14, 2025 10:45 AM
To: Ross McKittrick <ross.mckittrick@gmail.com>; Steve Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>
Subject: [2.2.2 and 2.2.3]Apr14[JC] Figs. Heat/Cold waves

Roy

I'm sending these figures that now use 1211 USHCN stations. I'm not inserting these into the document as they are for consideration. The data can be displayed however we would like. The 20-year running averages give a better climate picture I think. I believe one of the complaints of NCA5 was that the NCA5 group did no research on their own, whereas this would show we had the expertise to do the actual investigation.


In some way the point should be made that the temperature extremes are 1+ orders of magnitude greater than the small rise due to GHGs, so they will still occur. The US has not seen the weather patterns that created the 1899 Arctic outbreak nor the heatwaves of the 1930s, but the climate system is entirely capable of throwing one of these at us at any time regardless of GHGs. Regionally, summer patterns favored heat in the West in the 21st century, but heat in the central/east in first-half of 20th century.

Ross.

I assume you will hold the "Master" that combines all sections once each author is comfortable with letting text/figures be included?


John C.

CONUS Average of 1211 USHCN Stations Heatwave Days Per Year
Value = number of days Daily Max > 90th percentile in 7 or more consecutive days



CONUS Average of 1211 USHCN Stations Coldwave Days Per Year

Value = number of days Daily Min < 10th percentile in 7 or more consecutive days



From: Roy Spencer <roywspencer@hotmail.com>
Sent: Tuesday, April 15, 2025 4:56 AM
To: John Christy
Cc: Ross McKittrick; Steve Koonin
Subject: Re: [2.2.2 and 2.2.3]Apr14[JC] Figs. Heat/Cold waves

Those look good to me.
-Roy

From: John Christy <climateman60@gmail.com>
Sent: Monday, April 14, 2025 5:20 PM
To: Roy Spencer <roywspencer@hotmail.com>
Cc: Ross McKittrick <ross.mckittrick@gmail.com>; Steve Koonin <steven.koonin@gmail.com>
Subject: Re: [2.2.2 and 2.2.3]Apr14[JC] Figs. Heat/Cold waves

Roy

See attached.

John C.

On Mon, Apr 14, 2025 at 10:54 AM Roy Spencer <roywspencer@hotmail.com> wrote:
John:

My preference would be to leave out the regional results in the 1st and 3rd plots... graphs are too busy. Just keep the USA48 results.

-Roy

From: John Christy <climateman60@gmail.com>
Sent: Monday, April 14, 2025 10:45 AM
To: Ross McKittrick <ross.mckittrick@gmail.com>; Steve Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>
Subject: [2.2.2 and 2.2.3]Apr14[JC] Figs. Heat/Cold waves

Roy

I'm sending these figures that now use 1211 USHCN stations. I'm not inserting these into the document as they are for consideration. The data can be displayed however we would like. The 20-year running averages give a better climate picture I think. I believe one of the complaints of NCA5 was that the NCA5 group did no research on their own, whereas this would show we had the expertise to do the actual investigation.

In some way the point should be made that the temperature extremes are 1+ orders of magnitude greater than the small rise due to GHGs, so they will still occur. The US has not seen the weather patterns that created the

1899 Arctic outbreak nor the heatwaves of the 1930s, but the climate system is entirely capable of throwing one of these at us at any time regardless of GHGs. Regionally, summer patterns favored heat in the West in the 21st century, but heat in the central/east in first-half of 20th century.

Ross.

I assume you will hold the "Master" that combines all sections once each author is comfortable with letting text/figures be included?

John C.

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Monday, April 14, 2025 11:54 AM
To: John Christy; Ross McKittrick; Steve Koonin
Subject: Re: [2.2.2 and 2.2.3]Apr14[JC] Figs. Heat/Cold waves

John:

My preference would be to leave out the regional results in the 1st and 3rd plots... graphs are too busy. Just keep the USA48 results.

-Roy

From: John Christy <climateman60@gmail.com>
Sent: Monday, April 14, 2025 10:45 AM
To: Ross McKittrick <ross.mckittrick@gmail.com>; Steve Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>
Subject: [2.2.2 and 2.2.3]Apr14[JC] Figs. Heat/Cold waves

Roy

I'm sending these figures that now use 1211 USHCN stations. I'm not inserting these into the document as they are for consideration. The data can be displayed however we would like. The 20-year running averages give a better climate picture I think. I believe one of the complaints of NCA5 was that the NCA5 group did no research on their own, whereas this would show we had the expertise to do the actual investigation.

In some way the point should be made that the temperature extremes are 1+ orders of magnitude greater than the small rise due to GHGs, so they will still occur. The US has not seen the weather patterns that created the 1899 Arctic outbreak nor the heatwaves of the 1930s, but the climate system is entirely capable of throwing one of these at us at any time regardless of GHGs. Regionally, summer patterns favored heat in the West in the 21st century, but heat in the central/east in first-half of 20th century.

Ross.

I assume you will hold the "Master" that combines all sections once each author is comfortable with letting text/figures be included?

John C.

From: John Christy [climateman60@gmail.com]
Sent: 4/15/2025 9:19:42 PM
To: Ross McKittrick [ross.mckittrick@gmail.com]
CC: Steve Koonin [steven.koonin@gmail.com]; Roy Spencer [roywspencer@hotmail.com]
Subject: Re: [2.2.4]Apr15[JC] Ext Precip
Attachments: PacCoast_Prcp_20241231.xlsx

Ross

Attached is file for the Pacific Coastal stations (29) San Francisco began in 1849, but this list starts in 1877 with 3 stations reporting back then. All stations are complete from 1893-2024.


It's common practice to use the Water Year calendar in the West. Hydrologists use Oct-Sep for water management tasks, I use Aug-Jul because 1 Aug is the climatological driest day on the West Coast.

Attached is an example answering the question, "When did the 13 heaviest events occur along the Pacific Coast? (i.e., 1-in-10yr occurrence in 130 years). This result indicates Atmospheric Rivers are variable, but not trendy. The big El Nino of 1997-98 stands out. 29 stations times 13 events divided by 26 5-yr intervals gives an expected value of 14.5 events per interval. Note, lowest value in 2015-2019.


Same plot for the 24 humid SE stations. You can see the rise in the last few intervals due to the 4 eastern-most stations (3rd plot). I've double checked the observations with other stations and this result is real ... and it goes up to Maine (as I mentioned due to a series of tropical systems that hit the area between 1995 and 2019).

Let me know if anyone writing about this stuff is interested in any associated plots.


John C.



Distribution of 15 Heaviest Events per 5-year period
Each station contributes 15 events (i.e., nominal 1 in 10-year occurrence)
24 Stations in Humid Southeast, 1874-2023 Austin TX to Wash DC



Distribution of 15 Heaviest Events per 5-year period
Each station contributes 15 events (i.e., nominal 1 in 10-year occurrence)
4 Stations in Humid Southeast, 1874-2023 eNC, eVA, DC



On Tue, Apr 15, 2025 at 3:18 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:
Sure I can run that. Last year you send me long daily SE Humid zone records and I wrote the code to analyse those but never got around to writing up the results.

On Tue, Apr 15, 2025 at 11:49 AM John Christy <climateman60@gmail.com> wrote:

Ross

Attached is the updated daily precip (no missings) of 24 stations in the humid SE which, if your code is still handy, could be used as an update to our 2019 paper. Data are Oct 1872 to Dec 2024, so you could use either Oct - Sep Water year or calendar year (152 total years). Station list is in second tab.

Note: The four eastern-most stations account for the increase in heavy (5-day) events in last 25 years. This is seen in observations from eastern North Carolina up to Maine due to several tropical systems that drifted that way in the past 25 years and evidently accounts for the upward trend in heavy events in that part of the US.

I will send an update of the 29 Pacific Coastal stations which show no change in say 5-day extremes suggesting Atmospheric Rivers show no intensification.

John C.

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Wednesday, April 16, 2025 3:30 PM
To: John Christy
Cc: Steve Koonin; Roy Spencer
Subject: Re: [2.2.4]Apr15[JC] Ext Precip
Attachments: SE_Humid_Apr.16.2025.zip

John, here's a folder of outputs from the SE data. The charts show the trends by location, open circles mean insignificant. In each of the tables I also show the trends in averaged locations (coastal, inland and all). Probably the trends in annual max rainfall are most useful for the present purposes but everything is there. I'll work on running the Pacific stations.

On Tue, Apr 15, 2025 at 5:20 PM John Christy <climateman60@gmail.com> wrote:
Ross

Attached is file for the Pacific Coastal stations (29) San Francisco began in 1849, but this list starts in 1877 with 3 stations reporting back then. All stations are complete from 1893-2024.

It's common practice to use the Water Year calendar in the West. Hydrologists use Oct-Sep for water management tasks, I use Aug-Jul because 1 Aug is the climatological driest day on the West Coast.


Attached is an example answering the question, "When did the 13 heaviest events occur along the Pacific Coast? (i.e., 1-in-10yr occurrence in 130 years). This result indicates Atmospheric Rivers are variable, but not trendy. The big El Nino of 1997-98 stands out. 29 stations times 13 events divided by 26 5-yr intervals gives an expected value of 14.5 events per interval. Note, lowest value in 2015-2019.

Same plot for the 24 humid SE stations. You can see the rise in the last few intervals due to the 4 eastern-most stations (3rd plot). I've double checked the observations with other stations and this result is real ... and it goes up to Maine (as I mentioned due to a series of tropical systems that hit the area between 1995 and 2019.


Let me know if anyone writing about this stuff is interested in any associated plots.

John C.


Distribution of Heaviest 5-day Events by 5-yr periods
29 Pacific Coast Stations from San Diego CA to Blaine WA
Each station contributes 13 events (i.e., nominal 1 in 10-year occurrence)
Water Year Aug-July



Distribution of 15 Heaviest Events per 5-year period
Each station contributes 15 events (i.e., nominal 1 in 10-year occurrence)
24 Stations in Humid Southeast, 1874-2023 Austin TX to Wash DC



Distribution of 15 Heaviest Events per 5-year period
Each station contributes 15 events (i.e., nominal 1 in 10-year occurrence)
4 Stations in Humid Southeast, 1874-2023 eNC, eVA, DC



On Tue, Apr 15, 2025 at 3:18 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

Sure I can run that. Last year you send me long daily SE Humid zone records and I wrote the code to analyse those but never got around to writing up the results.

On Tue, Apr 15, 2025 at 11:49 AM John Christy <climateman60@gmail.com> wrote:

Ross

Attached is the updated daily precip (no missings) of 24 stations in the humid SE which, if your code is still handy, could be used as an update to our 2019 paper. Data are Oct 1872 to Dec 2024, so you could use either Oct - Sep Water year or calendar year (152 total years). Station list is in second tab.

Note: The four eastern-most stations account for the increase in heavy (5-day) events in last 25 years. This is seen in observations from eastern North Carolina up to Maine due to several tropical systems that drifted that way in the past 25 years and evidently accounts for the upward trend in heavy events in that part of the US.


I will send an update of the 29 Pacific Coastal stations which show no change in say 5-day extremes suggesting Atmospheric Rivers show no intensification.

John C.


Trends in Annual Rainfall Totals (inches)



Trends in Annual Average 2-day Rainfall Totals (inches)



Trends in Annual Maximum Rainfall since 1873



	"Trend"	"CI-lower"	"CI-upper"	"Sig"
"Austin"	-0.0056	-0.0511	0.0398	0
"Galveston"	-0.0169	-0.097	0.0632	0
"Shreveport"	0.049	-0.0447	0.1427	0
"Vicksburg"	0.0208	-0.0682	0.1097	0
"New Orleans"	0.0428	-0.0148	0.1004	0
"Memphis"	0.0506	-0.0491	0.1502	0
"Mobile"	0.0421	0.004	0.0803	1
"Nashville"	0.0151	-0.0609	0.0911	0
"Montgomery"	0.0043	-0.038	0.0466	0
"Atlanta"	0.0188	-0.0252	0.0627	0
"Knoxville"	0.0112	-0.0693	0.0916	0
"Macon"	-0.0171	-0.0778	0.0436	0
"Quitman"	-0.0364	-0.0621	-0.0107	1
"Augusta"	-0.0107	-0.0719	0.0504	0
"Key West"	0.016	-0.0057	0.0377	0
"Jacksonville"	0.0139	-0.0456	0.0733	0
"Lenoir"	-0.0023	-0.0455	0.041	0
"Savannah"	-0.0014	-0.0536	0.0508	0
"Charleston"	0.0327	-0.0929	0.1583	0
"Lynchburg"	0.0081	-0.0578	0.074	0
"Wilmington"	0.0828	-0.0251	0.1908	0
"Weldon"	0.0403	-0.0376	0.1181	0
"Washington"	-0.018	-0.0721	0.0361	0
"Norfolk"	0.0084	-0.0947	0.1114	0
"Coastal"	0.2204	-0.299	0.7399	0
"Inland"	0.1279	-0.6884	0.9443	0
"All"	0.3484	-0.9415	1.6382	0

	"Trend"	"CI-lower"	"CI-upper"	"Sig"
"Austin"	-0.0031	-0.028	0.0218	0
"Galveston"	-0.0092	-0.0531	0.0347	0
"Shreveport"	0.0269	-0.0245	0.0782	0
"Vicksburg"	0.0114	-0.0373	0.0601	0
"New Orleans"	0.0235	-0.0081	0.055	0
"Memphis"	0.0277	-0.0269	0.0823	0
"Mobile"	0.0231	0.0022	0.044	1
"Nashville"	0.0083	-0.0333	0.0499	0
"Montgomery"	0.0024	-0.0209	0.0256	0
"Atlanta"	0.0103	-0.0138	0.0344	0
"Knoxville"	0.0061	-0.0379	0.0502	0
"Macon"	-0.0094	-0.0426	0.0239	0
"Quitman"	-0.0199	-0.034	-0.0058	1
"Augusta"	-0.0059	-0.0394	0.0276	0
"Key West"	0.0088	-0.0031	0.0207	0
"Jacksonville"	0.0076	-0.025	0.0402	0
"Lenoir"	-0.0012	-0.0249	0.0225	0
"Savannah"	-0.0007	-0.0293	0.0278	0
"Charleston"	0.0179	-0.0509	0.0868	0
"Lynchburg"	0.0044	-0.0316	0.0405	0
"Wilmington"	0.0454	-0.0138	0.1046	0
"Weldon"	0.0221	-0.0206	0.0647	0
"Washington"	-0.0098	-0.0395	0.0198	0
"Norfolk"	0.0046	-0.0518	0.0611	0
"Coastal"	0.121	-0.1636	0.4056	0
"Inland"	0.0703	-0.377	0.5176	0
"All"	0.1913	-0.5154	0.898	0

	"Trend"	"CI-lower"	"CI-upper"	"Sig"
"Austin"	-0.1446	-0.3277	0.0385	0
"Galveston"	0.0146	-0.0215	0.0506	0
"Shreveport"	0.0612	0.0033	0.119	1
"Vicksburg"	0.0544	-0.0265	0.1353	0
"New Orleans"	0.0638	0.0178	0.1099	1
"Memphis"	0.0557	0.0204	0.091	1
"Mobile"	0.0456	0.008	0.0833	1
"Nashville"	0.0285	0.0086	0.0483	1
"Montgomery"	0.0334	-0.0053	0.0721	0
"Atlanta"	0.022	-0.036	0.0799	0
"Knoxville"	0.0271	-0.0086	0.0628	0
"Macon"	-0.1608	-0.4328	0.1113	0
"Quitman"	-0.2412	-0.3368	-0.1457	1
"Augusta"	0.0113	-0.0166	0.0391	0
"Key West"	0.0405	0.0263	0.0547	1
"Jacksonville"	0.0481	-0.0005	0.0968	0
"Lenoir"	-0.2276	-0.4022	-0.053	1
"Savannah"	0.0313	-0.0028	0.0653	0
"Charleston"	0.0422	-0.0467	0.131	0
"Lynchburg"	0.0207	-0.0243	0.0657	0
"Wilmington"	0.0825	0.0244	0.1407	1
"Weldon"	-0.0112	-0.1767	0.1542	0
"Washington"	0.015	-0.0023	0.0323	0
"Norfolk"	0.0402	-0.0085	0.089	0
"Coastal"	0.1162	-0.0143	0.2467	0
"Inland"	-0.0261	-0.3261	0.2738	0
"All"	0.0968	-0.2752	0.4688	0

	"Trend"	"CI-lower"	"CI-upper"	"Sig"
"Austin"	-0.0193	-0.0647	0.0262	0
"Galveston"	-0.0168	-0.0645	0.0308	0
"Shreveport"	0.0313	-0.0282	0.0907	0
"Vicksburg"	0.0251	-0.0191	0.0694	0
"New Orleans"	0.0442	-0.0047	0.0931	0
"Memphis"	0.0218	-0.0155	0.059	0
"Mobile"	0.0444	0.0108	0.0779	1
"Nashville"	0.0156	-0.0142	0.0454	0
"Montgomery"	0.0065	-0.0155	0.0286	0
"Atlanta"	0.0155	-0.0154	0.0465	0
"Knoxville"	0.0084	-0.0258	0.0427	0
"Macon"	-0.0298	-0.1134	0.0539	0
"Quitman"	-0.0438	-0.0972	0.0096	0
"Augusta"	0.0095	-0.0178	0.0368	0
"Key West"	0.0397	-0.0109	0.0904	0
"Jacksonville"	0.0219	-0.0134	0.0571	0
"Lenoir"	-0.0397	-0.1119	0.0324	0
"Savannah"	0.0155	-0.0098	0.0407	0
"Charleston"	0.018	-0.0637	0.0997	0
"Lynchburg"	0.0078	-0.017	0.0327	0
"Wilmington"	0.0914	-0.006	0.1888	0
"Weldon"	0.0156	-0.0653	0.0966	0
"Washington"	-0.0037	-0.0236	0.0163	0
"Norfolk"	0.0397	-0.0153	0.0948	0
"Coastal"	0.5466	-0.0855	1.1788	0
"Inland"	-0.1725	-2.0315	1.6865	0
"All"	0.9239	-2.6116	4.4595	0

	"Trend"	"CI-lower"	"CI-upper"	"Sig"
"Austin"	-0.4126	-1.3394	0.5142	0
"Galveston"	-0.3131	-1.1154	0.4893	0
"Shreveport"	0.1345	-0.7354	1.0043	0
"Vicksburg"	0.4425	0.0254	0.8595	1
"New Orleans"	0.2529	-0.5796	1.0855	0
"Memphis"	-0.0984	-0.6483	0.4515	0
"Mobile"	0.5101	-0.0286	1.0487	0
"Nashville"	0.153	-0.3503	0.6563	0
"Montgomery"	-0.0362	-0.7458	0.6734	0
"Atlanta"	0.2255	-0.2161	0.667	0
"Knoxville"	0.1281	-0.1064	0.3627	0
"Macon"	-0.5225	-1.6686	0.6237	0
"Quitman"	-0.5879	-1.4677	0.292	0
"Augusta"	0.3468	-0.1117	0.8052	0
"Key West"	0.6592	-0.2485	1.5668	0
"Jacksonville"	0.4021	-0.3265	1.1307	0
"Lenoir"	-0.5949	-1.6687	0.4788	0
"Savannah"	0.192	-0.344	0.7281	0
"Charleston"	0.3682	-0.2388	0.9751	0
"Lynchburg"	0.2251	-0.1788	0.6289	0
"Wilmington"	1.1429	-0.2246	2.5105	0
"Weldon"	0.3309	-1.1542	1.8159	0
"Washington"	0.0016	-0.6074	0.6106	0
"Norfolk"	1.0065	0.0957	1.9173	1
"Coastal"	1.39	0.3461	2.4339	1
"Inland"	-0.522	-2.6805	1.6364	0
"All"	0.8218	-1.1368	2.7804	0

	"Trend"	"CI-lower"	"CI-upper"	"Sig"
"Austin"	-0.0006	-0.0028	0.0017	0
"Galveston"	0.0003	-0.0012	0.0019	0
"Shreveport"	0.0022	-0.001	0.0054	0
"Vicksburg"	0.0002	-0.0016	0.002	0
"New Orleans"	0.0009	-0.0017	0.0034	0
"Memphis"	-0.0001	-0.0035	0.0033	0
"Mobile"	0.001	-0.0006	0.0026	0
"Nashville"	-0.0007	-0.0027	0.0014	0
"Montgomery"	-0.0007	-0.0023	0.0009	0
"Atlanta"	0.0016	0.0001	0.003	1
"Knoxville"	0.0007	-0.0009	0.0022	0
"Macon"	-0.0011	-0.0039	0.0017	0
"Quitman"	-0.0017	-0.0041	0.0007	0
"Augusta"	0.0026	0.0008	0.0044	1
"Key West"	0.0011	-0.0005	0.0027	0
"Jacksonville"	0.0004	-0.0031	0.0039	0
"Lenoir"	-0.0019	-0.0056	0.0019	0
"Savannah"	-0.0009	-0.0028	0.001	0
"Charleston"	0	-0.0026	0.0025	0
"Lynchburg"	0.0004	-0.0019	0.0027	0
"Wilmington"	0.0021	-0.001	0.0053	0
"Weldon"	-0.0001	-0.0051	0.0049	0
"Washington"	-0.0004	-0.0026	0.0018	0
"Norfolk"	0.0034	0.0003	0.0065	1
"Coastal"	0.0139	0.0035	0.0243	1
"Inland"	-0.0052	-0.0268	0.0164	0
"All"	0.0082	-0.0114	0.0278	0

	"Trend"	"CI-lower"	"CI-upper"	"Sig"
"Austin"	-0.0001	-0.0017	0.0015	0
"Galveston"	-0.0007	-0.0013	-0.0001	1
"Shreveport"	0	-0.0005	0.0006	0
"Vicksburg"	0.0002	-0.0003	0.0007	0
"New Orleans"	0.0002	-0.0001	0.0005	0
"Memphis"	0.0001	-0.0006	0.0008	0
"Mobile"	0.0002	-0.0001	0.0004	0
"Nashville"	0.0001	-0.0002	0.0004	0
"Montgomery"	0.0002	-0.0002	0.0005	0
"Atlanta"	0	0	0	0
"Knoxville"	0	0	0	0
"Macon"	0	-0.0004	0.0004	0
"Quitman"	0.0002	-0.0004	0.0007	0
"Augusta"	0	0	0	0
"Key West"	0.0002	-0.0001	0.0005	0
"Jacksonville"	0.0001	-0.0003	0.0004	0
"Lenoir"	0	0	0	0
"Savannah"	0	0	0	0
"Charleston"	0.0001	-0.0003	0.0004	0
"Lynchburg"	0	0	0	0
"Wilmington"	0.0003	-0.0003	0.001	0
"Weldon"	0.0001	-0.0002	0.0004	0
"Washington"	0	0	0	0
"Norfolk"	0.0002	-0.0004	0.0008	0
"Coastal"	0.0139	0.0035	0.0243	1
"Inland"	-0.0052	-0.0268	0.0164	0
"All"	0.0082	-0.0114	0.0278	0

From: John Christy [climateman60@gmail.com]
Sent: 4/16/2025 8:47:30 PM
To: Ross McKittrick [ross.mckittrick@gmail.com]
CC: Steve Koonin [steven.koonin@gmail.com]; Roy Spencer [roywspencer@hotmail.com]
Subject: Re: [2.2.4]Apr15[JC] Ext Precip

Got it. Roy - this is mostly for you since you have the extreme sections. Looks like nothing really remarkable.

John C.

On Wed, Apr 16, 2025 at 2:30 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

John, here's a folder of outputs from the SE data. The charts show the trends by location, open circles mean insignificant. In each of the tables I also show the trends in averaged locations (coastal, inland and all). Probably the trends in annual max rainfall are most useful for the present purposes but everything is there. I'll work on running the Pacific stations.

On Tue, Apr 15, 2025 at 5:20 PM John Christy <climateman60@gmail.com> wrote:

Ross

Attached is file for the Pacific Coastal stations (29) San Francisco began in 1849, but this list starts in 1877 with 3 stations reporting back then. All stations are complete from 1893-2024.

It's common practice to use the Water Year calendar in the West. Hydrologists use Oct-Sep for water management tasks, I use Aug-Jul because 1 Aug is the climatological driest day on the West Coast.


Attached is an example answering the question, "When did the 13 heaviest events occur along the Pacific Coast? (i.e., 1-in-10yr occurrence in 130 years). This result indicates Atmospheric Rivers are variable, but not trendy. The big El Nino of 1997-98 stands out. 29 stations times 13 events divided by 26 5-yr intervals gives an expected value of 14.5 events per interval. Note, lowest value in 2015-2019.

Same plot for the 24 humid SE stations. You can see the rise in the last few intervals due to the 4 eastern-most stations (3rd plot). I've double checked the observations with other stations and this result is real ... and it goes up to Maine (as I mentioned due to a series of tropical systems that hit the area between 1995 and 2019.


Let me know if anyone writing about this stuff is interested in any associated plots.

John C.


Distribution of Heaviest 5-day Events by 5-yr periods
29 Pacific Coast Stations from San Diego CA to Blaine WA
 Each station contributes 13 events (i.e., nominal 1 in 10-year occurrence)
 Water Year Aug-July



Distribution of 15 Heaviest Events per 5-year period
Each station contributes 15 events (i.e., nominal 1 in 10-year occurrence)
24 Stations in Humid Southeast, 1874-2023 Austin TX to Wash DC



Distribution of 15 Heaviest Events per 5-year period
Each station contributes 15 events (i.e., nominal 1 in 10-year occurrence)
4 Stations in Humid Southeast, 1874-2023 eNC, eVA, DC



On Tue, Apr 15, 2025 at 3:18 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:
Sure I can run that. Last year you send me long daily SE Humid zone records and I wrote the code to analyse those but never got around to writing up the results.

On Tue, Apr 15, 2025 at 11:49 AM John Christy <climateman60@gmail.com> wrote:
Ross

Attached is the updated daily precip (no missings) of 24 stations in the humid SE which, if your code is still handy, could be used as an update to our 2019 paper. Data are Oct 1872 to Dec 2024, so you could use either Oct - Sep Water year or calendar year (152 total years). Station list is in second tab.

Note: The four eastern-most stations account for the increase in heavy (5-day) events in last 25 years. This is seen in observations from eastern North Carolina up to Maine due to several tropical systems that drifted that way in the past 25 years and evidently accounts for the upward trend in heavy events in that part of the US.

I will send an update of the 29 Pacific Coastal stations which show no change in say 5-day extremes suggesting Atmospheric Rivers show no intensification.

John C.

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Tuesday, April 15, 2025 4:19 PM
To: John Christy
Cc: Steve Koonin; Roy Spencer
Subject: Re: [2.2.4]Apr15[JC] Ext Precip

Sure I can run that. Last year you send me long daily SE Humid zone records and I wrote the code to analyse those but never got around to writing up the results.

On Tue, Apr 15, 2025 at 11:49 AM John Christy <climateman60@gmail.com> wrote:
Ross

Attached is the updated daily precip (no missings) of 24 stations in the humid SE which, if your code is still handy, could be used as an update to our 2019 paper. Data are Oct 1872 to Dec 2024, so you could use either Oct - Sep Water year or calendar year (152 total years). Station list is in second tab.

Note: The four eastern-most stations account for the increase in heavy (5-day) events in last 25 years. This is seen in observations from eastern North Carolina up to Maine due to several tropical systems that drifted that way in the past 25 years and evidently accounts for the upward trend in heavy events in that part of the US.

I will send an update of the 29 Pacific Coastal stations which show no change in say 5-day extremes suggesting Atmospheric Rivers show no intensification.

John C.

From: Fisher, Travis [travis.fisher@hq.doe.gov]
Sent: 7/31/2025 9:17:59 PM
To: Judith Curry [curry.judith@gmail.com]; Roy Spencer [roywspencer@hotmail.com]; Park, Charles [charles.park@hq.doe.gov]
CC: Ross McKittrick [ross.mckittrick@gmail.com]; John Christy [climatemanager60@gmail.com]; Steven Koonin [steven.koonin@googlemail.com]; Loucks, Joshua [Joshua.Loucks@hq.doe.gov]; Cohen, Seth [seth.cohen@hq.doe.gov]
Subject: RE: [EXTERNAL] DOE reviews
Attachments: DOE Report Review Master Ordered.xlsx

I'm attaching the version of the internal review comments that was compiled by the Office of Science. Note that I haven't checked to see if they numbered the reviewers the same way we did.

Even though we haven't had our introductory call yet, I'm also including Charles Park. Let me know if you're available tomorrow afternoon for a call to introduce Charles.

And I just saw Steve's email—yes, we should compile the anonymous reviews and your responses, and we should strongly consider releasing that document (and I believe we would have to if anyone requests it).

From: Judith Curry <curry.judith@gmail.com>
Sent: Thursday, July 31, 2025 5:06 PM
To: Roy Spencer <roywspencer@hotmail.com>
Cc: Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climatemanager60@gmail.com>; Steven Koonin <steven.koonin@googlemail.com>; Josh Loucks <loucksj14@gmail.com>; Cohen, Seth <seth.cohen@hq.doe.gov>; Fisher, Travis <travis.fisher@hq.doe.gov>
Subject: [EXTERNAL] DOE reviews

Ross, if you can send me the latest version of the 8 DOE review responses, i can clean those up, verify changes were made, and send around for approval. Would be good to get those out of the way before the public comments

On Thu, Jul 31, 2025 at 12:27 PM Roy Spencer <roywspencer@hotmail.com> wrote:

Ironic that no one cared when MY work was mischaracterized by the 97-Percenters' as being part of the 3%, even though by their criteria I was part of the 97% consensus.

I complained, but all they said was, "well, we put you in the 3 percent category".

The bias is rampant.

-Roy

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 31, 2025 11:27 AM
To: Roy W. Spencer <roywspencer@hotmail.com>; Judith Curry <curry.judith@gmail.com>; John Christy <climatemanager60@gmail.com>; Steven Koonin <steven.koonin@googlemail.com>; Josh Loucks <loucksj14@gmail.com>; Cohen, Seth <seth.cohen@hq.doe.gov>; Fisher, Travis <travis.fisher@hq.doe.gov>
Subject: Fwd: Fw: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

John and I got this from Tim Vogelsang at MSU. Carbon Brief is contacting everyone whose work was cited trying to gather up claims of misrepresentation. As Tim notes, there is no option on the form for saying your work was correctly characterized.

From: Vogelsang, Timothy <tjv@msu.edu>
Sent: Thursday, July 31, 2025 12:19 PM
To: Ross McKittrick <rmckitri@uoguelph.ca>; John Christy <climateman60@gmail.com>
Subject: RE: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

CAUTION: This email originated from outside of the University of Guelph. Do not click links or open attachments unless you recognize the sender and know the content is safe. If in doubt, forward suspicious emails to IThelp@uoguelph.ca.

If you look at the pdf I was sent with the instructions, there is no option to report that someone's research has been correctly characterized. That's not a very scientific factchecking exercise, is it!

Tim

From: Ross McKittrick <rmckitri@uoguelph.ca>
Sent: Thursday, July 31, 2025 12:15 PM
To: Vogelsang, Timothy <tjv@msu.edu>; John Christy <climateman60@gmail.com>
Subject: Re: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

I've heard of Carbon Brief but only read snippets of their articles cited elsewhere. They seem to be heavily into promoting renewables.

It's interesting to watch the machine swing into gear. There is a public comment docket opening up and I expect there will be quite the avalanche of material submitted, which we will need to prepare responses to, all of which will form part of the public record. If the comments are substantive we'll readily make revisions. We responded to a lot of comments from the internal DoE staff during the pre-publication review stage. But some of the most hostile comments didn't provide any data or citations in support so we couldn't do anything with them.

Ross

Dr. Ross McKittrick
Professor of Economics
Department of Economics and Finance
rossmckittrick.com



University of Guelph | Guelph ON Canada N1G 3P9

From: Vogelsang, Timothy <tjv@msu.edu>
Sent: Thursday, July 31, 2025 12:02 PM

To: Ross McKittrick <rmckitri@uoguelph.ca>; John Christy <climateman60@gmail.com>
Subject: FW: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

CAUTION: This email originated from outside of the University of Guelph. Do not click links or open attachments unless you recognize the sender and know the content is safe. If in doubt, forward suspicious emails to ITheelp@uoguelph.ca.

Ross and John,

I just got this email. Maybe you know about this group. I'm sure this is just the tip of the iceberg of pushback coming.

Tim

From: Leo Hickman <leo.hickman@carbonbrief.org>
Sent: Thursday, July 31, 2025 11:49 AM
To: Leo Hickman <leo.hickman@carbonbrief.org>
Subject: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

Hello,

I'm contacting you because we have noticed that at least one of your studies has been cited in a new report published this week by the US Dept of Energy titled, "A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate."

A number of scientists cited in the report have already publicly highlighted that their findings have been falsely or misleadingly characterised. It seems that the report is intended to be used as scientific justification by the Trump administration in its efforts to revoke the US Environmental Protection Agency's "endangerment finding" on greenhouse gases.

Carbon Brief is now seeking your help to assess whether your own study, or indeed your area(s) of expertise more widely, have been mischaracterised in the report. This is part of a wider 'crowdsourced' fact-checking effort we are now convening to draw on inputs from experts in their respective fields.

I have attached a PDF with some instructions on how you can participate with your own insights and reflections, should you wish. (The instructions include ways to do this anonymously and securely, if that is your preference.)

Additionally, if you know of a colleague or co-author who might wish to participate, please feel free to forward this email to them.

Our deadline for submissions is Monday 11 August 2025 - 0900 GMT.

Many thanks in advance.

Best wishes, Leo

--

Leo Hickman

Editor, Carbon Brief

Phone: + 44 (0) 7731 403979

Social: Twitter; BlueSky; LinkedIn

Web: www.carbonbrief.org

Need a digest of the latest climate change-related headlines sent to your inbox? You can sign up to Carbon Brief's free daily, weekly, Cropped and China Briefing newsletters here.

--



Judith Curry, President
CFAN - Climate Forecast Applications
Network
Reno, NV USA
curry.judith@cfanclimate.com |
+1.404.803.2012
<http://www.cfanclimate.net>

This message does not originate from a known Department of Energy email system.
Use caution if this message contains attachments, links or requests for information.

Reviewer Number	Chapter Number	Page Number	Approximate Paragraph Number
--------------------	-------------------	-------------	------------------------------------

1	General comment	Overarching	Overarching
---	--------------------	-------------	-------------

2	General comment		
---	--------------------	--	--

3	General comment		
---	--------------------	--	--

6	Title Page	i	
6	Title Page	iii	1
8	Foreword	vii	4
8	Foreword	vii	8
6	Forward	vii	Forward
8	Executive Summary	viii	2
8	Executive Summary	viii	2

8	Executive Summary	viii	3
---	----------------------	------	---

8	Executive Summary	viii	4
---	----------------------	------	---

8	Executive Summary	viii	5
---	----------------------	------	---

8	Executive Summary	viii	6
---	----------------------	------	---

8	Executive Summary	viii	6
---	----------------------	------	---

8	Executive Summary	viii	9
---	----------------------	------	---

1	Executive Summary	viii	1
---	----------------------	------	---

3	Executive Summary	viii	3
---	----------------------	------	---

3	Executive Summary	viii	5
---	----------------------	------	---

3	Executive Summary	viii	6
---	----------------------	------	---

7	Executive Summary	viii	2
---	----------------------	------	---

6	Executive Summary	viii	2
---	----------------------	------	---

6	Executive Summary	viii	3
---	----------------------	------	---

6	Executive Summary	viii	4
---	----------------------	------	---

6	Executive Summary	viii	6
6	Executive Summary	viii	8
6	Executive Summary	viii	9
7	Executive Summary	viii	3
7	Executive Summary	viii	4
7	Executive Summary	viii	5
7	Executive Summary	viii	5,6,7

7	Executive Summary	viii	9
1	Preface	ix	3
1	Preface	ix	3
1	Preface	ix	3
8	Preface	ix	1
1	1	2	1

1	1	2	1
---	---	---	---

1	1	2	4
---	---	---	---

4	1	2	full text
6	1	2	

7	1	2	3
---	---	---	---

1	2	3	1
---	---	---	---

4	2	3	3
---	---	---	---

4	2	3	3
---	---	---	---

4	2	3	3
---	---	---	---

4	2	3	4
---	---	---	---

4	2	3	4
---	---	---	---

4	2	3	4
---	---	---	---

4	2	3	3 and 4
---	---	---	---------

4	2	3	3 and 4
---	---	---	---------

4	2	3	5
---	---	---	---

4	2	3	5
5	2	3	2.1.1, para. 2
	2	3	2.1.1, para. 3
5			
	2	3	2.1.1, para.1
5			
	2	3	2.1.1, para.2
5			
	2	3	2.1.1, para.2
5			
	2	3	2.1.1, para.3
5			
8	2	3	1
8	2	3	2
7	2	3	1/Chapter summary

7	2	3	3
---	---	---	---

7	2	3	5
---	---	---	---

4	2	4	1
---	---	---	---

4	2	4	2
---	---	---	---

	2	4	2.1.2
--	---	---	-------

5

	2	4	2.1.1, last para.
--	---	---	-------------------

5

6	2	4	1
---	---	---	---

7	2	4	1
---	---	---	---

2	2	5	Fig 2.2
---	---	---	---------

6	2	5	Figure 2.2
---	---	---	------------

8	2	5	1
---	---	---	---

7	2	5	Fig 2.2
---	---	---	---------

7	2	6	1
---	---	---	---

4	2	7	4
5	2	7	2.1.3, para. 2

6	2	7	3
---	---	---	---

4	2	8	2
8	2	8	1
8	2	8	1
7	2	8	1
6	2	9	3
8	2	9	4
5	2	6-7	2.1.3

3	2.1	3-6
---	-----	-----

3	2.2	7
---	-----	---

7	3	12	Summary
---	---	----	---------

8	3	13	2
---	---	----	---

4	3	14	1
---	---	----	---

8	3	14	4
---	---	----	---

8	3	14	5
---	---	----	---

1	3	15	3
---	---	----	---

8	3	15	1
---	---	----	---

1	3	16	4
7	3	16	2,3,4,5
1	3	17	2
4	3	18	1
	3	19	3.2.2, last para.
5			
6	3	19	last
7	3	19	5
8	3	20	2

1	3	22	6
7	3	22	3
2	3	11,12	Sec 3.1.1
2	3	21,22	Sec 3.3
3	3.1	14	4
3	3.2	17	figure
3	3.2	19	
3	3.2	19	5
3	3.2	20	
3	3.3		
1	4	26	1
1	4	26	2

1	4	26	3
---	---	----	---

1	4	26	6
---	---	----	---

1	4	26	6
---	---	----	---

1	4	26	6
---	---	----	---

8	4	26	6
---	---	----	---

1	4	28	2
---	---	----	---

7	4	28	2
---	---	----	---

1	4	29	3
---	---	----	---

8	4	29	1
---	---	----	---

8	4	29	2
---	---	----	---

7	4	29	2
---	---	----	---

2	4	30	Sec 4.4
---	---	----	---------

8	4	30	1
---	---	----	---

3	4.2	27	2
---	-----	----	---

3	4.2	27	3
---	-----	----	---

3	4.3		
---	-----	--	--

1	5	32	1
---	---	----	---

8	5	32	1
---	---	----	---

7	5	32	3
---	---	----	---

1	5	33	Figure
---	---	----	--------

1	5	34	1
---	---	----	---

1	5	34	4
---	---	----	---

8	5	34	1
---	---	----	---

8	5	34	2
---	---	----	---

7	5	34	1
---	---	----	---

8	5	35	1
---	---	----	---

8	5	36	1
---	---	----	---

7	5	36	1
---	---	----	---

1	5	37	3
---	---	----	---

8	5	37	1
---	---	----	---

7	5	38	3
---	---	----	---

1	5	39	Figure 5.7
---	---	----	------------

1	5	39	3
---	---	----	---

8	5	39	1
---	---	----	---

8	5	39	1
---	---	----	---

1	5	40	1
---	---	----	---

1	5	40	5
---	---	----	---

7	5	40	4
---	---	----	---

8	5	41	1
---	---	----	---

8	5	42	4
---	---	----	---

7	5	42	2
---	---	----	---

6	5	44	1
---	---	----	---

6	5	44	1
---	---	----	---

7	5	44	3
2	5	37, 38	Sec 5.4
3	5.1	33	
3	5.2	34	figure
3	5.3	36	figure 5.4
3	5.4	37	4
3	5.4	38	
3	5.7	42	
1	6	47	1
1	6	47	1
1	6	47	3
6	6	47	chapter summary

6	6	47	chapter summary
8	6	47	5
7	6	47	2
7	6	47	3
7	6	47	5
1	6	50	Figure
1	6	52	4
1	6	53	Table
1	6	53	5
8	6	53	4
1	6	54	1

8	6	56	1
2	6	57	para 1
6	6	57	6.3.2
7	6	57	2
1	6	58	Figure
1	6	58	4
1	6	58	4
8	6	58	3
7	6	58	3
1	6	59	3
7	6	59	4

7	6	60	1,2
1	6	61	3 to 6

8	6	61	3
---	---	----	---

8	6	61	7
---	---	----	---

7	6	61	4
---	---	----	---

8	6	62	1
7	6	62	2

7	6	63	box
---	---	----	-----

7	6	64	1
1	6	67	2

1	6	68	7
---	---	----	---

8	6	68	3
1	6	69	1

1	6	69	3
---	---	----	---

1	6	69	4
---	---	----	---

7	6	69	2
---	---	----	---

7	6	70	2
---	---	----	---

7	6	71	3
---	---	----	---

3	6		6.4, 6.6, 6.7
---	---	--	---------------

3	6.2		
---	-----	--	--

3	6.3	60	
3	6.3	56-57	
6	7	74	2
1	7	75	1
7	7	80	1
6	7	all	
3	7		
1	8	83	3
1	8	84	6
1	8	84	6
2	8	85	para 5
6	8	86	all
7	8	88	3

1	8	90	3
1	8	92	2
7	8	92	4
1	8	93	4
7	8	93	3
3	8	94	table 8.1
2	8	85,86,87	"Natural variability of large-scale ocean circulations"
2	8	88,89	Optimal fingerprinting discussion
2	8	90,91,92	Section 8.4
3	8.3	84	3
3	8.3	87-88	

3	8.4	91	2 and 3
3	8.6	95	2
3	8.6		
1	9	104	1
	9	104	all
5			
6	9	104	chapter summary
	9	105	9.2, para. 2
5			
	9	105	9.2, para. 3, 4, 5
5			
	9	105	9.2, para. 3
5			

7	9	105	2,3,4,5
---	---	-----	---------

	9	106	9.2, para. 2
--	---	-----	--------------

5

7	9	106	1
---	---	-----	---

	9	107	1st para.
--	---	-----	-----------

5

	9	108	1st
--	---	-----	-----

5

	9	108	9.4, para. 1
--	---	-----	--------------

5

	9	109	
--	---	-----	--

5

3	9.1	104	
---	-----	-----	--

1	10	110	4
---	----	-----	---

1	10	111	1
---	----	-----	---

1	10	111	2
---	----	-----	---

7	10	111	1,2
---	----	-----	-----

1	10	112	1
---	----	-----	---

1	10	113	7
---	----	-----	---

3	10.2	111	
---	------	-----	--

3	10.3	114	
---	------	-----	--

3	10.3		
---	------	--	--

1	11	116	1
---	----	-----	---

1	11	117	5
7	11	117	3

7	11	118	3
---	----	-----	---

1	11	119	3
---	----	-----	---

1	11	119	6
---	----	-----	---

7	11	120	1
---	----	-----	---

1	11	121
---	----	-----

1	11	121	3
---	----	-----	---

1	11	121	4
---	----	-----	---

1	11	122	6
---	----	-----	---

1	11	122	7
---	----	-----	---

1	11	123	2
---	----	-----	---

1	11	125	5
---	----	-----	---

1	11	116-120	Section 11.1
---	----	---------	--------------

1	11	123-125	Section 11.1
---	----	---------	--------------

1	11	Chapter	Chapter
---	----	---------	---------

3	11.1		
---	------	--	--

3	11.2		
---	------	--	--

1	12	129	1
---	----	-----	---

1	12	129	2
---	----	-----	---

1	12	129	3
---	----	-----	---

1	12	129	4
---	----	-----	---

1	12	129	5
---	----	-----	---

1

12

129

Chapter

1

12

129

Chapter

1

12

129

Chapter

1	12	129	Chapter
---	----	-----	---------

4	12	129	3
---	----	-----	---

4	12	129	5
---	----	-----	---

	12	129	
--	----	-----	--

5			
---	--	--	--

7	12	130	4
---	----	-----	---

3	12		
---	----	--	--

Comment or Suggested Edit

- I would like to thank the authors for the opportunity to provide comment on this report. Providing decision-makers with information about the impacts of carbon dioxide emissions on the U.S. climate is critical as they face consequential decisions about the future of energy, agricultural, and environmental policy. Assessments of the science such as IPCC reports and the NCA benefit from undergoing robust review including expert review, public review, and government review with review comments and responses made publicly available. Similarly, important government reports and regulations undergo inter-agency review and are published in the Federal Register for a full notice and comment process and publish a public response to comment. While these processes are far from perfect, they offer the opportunity for improvement and help lend authority to the final products, and I highly encourage you to undertake a similar robust and open peer review process for this report to enhance the report's credibility in the eyes of decision-makers and the public.

In the limited time allowed, this reviewer concentrated on sections related to the physical climate and climate change. Frankly, the document was discouraging. Secretary Wright wants a " ... a more thoughtful and science-based conversation about climate change and energy." I find that this report distorts the science in much the same way as the media coverage about which he complains. Most notable are the omissions or under-reported significant climate changes, such as sea-ice decline, changes in the global water cycle and increases in heat-stress due to increases in absolute humidity (from water-vapor feedback). Further, the examples used by the report authors are narrow and appear to be selected to emphasize uncertainty, while the more rigorously evaluated science that contradicts their limited discussion is ignored, perhaps because it does not fit their narrative. This is unfortunate, because I believe it detracts from the much needed discussion about the future, including energy policies, technologies and trade-offs. I encourage DOE to start the discussion by accepting the well-reviewed and broadly accepted scientific conclusions of the IPCC Working Group 1, with respect to the observations and analyses of climate change in the 20th and 21st centuries. Skepticism regarding predictions/projections of future climate based on posited scenarios is justified. As a member of the physical climate research community, I believe there are better uses for the models (and the computer resources they consume) than scenario projection simulations, to inform the discussion that is urgently needed about the future.

The report indicates numerous examples of uncertainties in climate science. It would be extremely helpful if these uncertainties were made more explicit, listed in a concluding section, and prioritized in some way - what are the top 10 priority research areas? If the opening remarks by Secretary Wright and the title pointed to uncertainties and the need for more research, I believe this report will be better received. As I read the report and left review comments below, **I bolded items where I gather the authors imply that research is needed.** Consider including these kinds of statements throughout and adding a summary.

In general, criticism and re-analysis of past research is always welcome. This report summarizes some substantial concerns from a small group of researchers. The report should be reframed to reflect this and I provide some suggestions in a later comment. I suspect that simply publishing this report will not offset the hundreds of reports and publications on CO2 and Climate that have come before it. But, new analyses and communication within and outside the research community could make a positive difference, should the findings in this report be substantiated. The DOE Office of Science BER could be provided an opportunity to address the research uncertainties in this document in a thorough and robust manner.

As someone who used to read the DOE reports on CO2 as a graduate student in the 1990s and then conducted numerous analyses of carbon on many topic areas which were later confirmed by numerous other analyses, it seems somewhat cavalier for DOE to issue one brief conclusive report on the "impacts of carbon dioxide emissions". I suggest the following option: Change the title to "Topics of Uncertainty on the Impacts of Carbon Dioxide Emissions on the U.S. Climate" or "**An Assessment of CO2 Impacts on Climate to Guide Future Research**", and then use this report to drive research/analyses for each report chapter which would conclude in separate topical reports that could be finished within a year or two. In doing so, the DOE would positively contribute to addressing some of these standing issues and hopefully enable the nation to move forward in a more productive manner.

The stated charge to the team was to summarize the current state of climate science, but in the Preface the team states that it chose to focus on topics that are downplayed in or absent from current assessment reports. To accurately describe the contents of the report, the Foreword should also state that the report is a summary of the things that the author team thinks have been downplayed in major climate assessment reports.

The statement about climate change not being the greatest threat facing humanity is a value judgement. Since the Secretary criticizes others for blurring the line between scientific conclusions and policy recommendations in the second, fourth, and sixth paragraphs of this Foreword, it is hypocritical to offer a value judgement without clearly labeling it as such and noting that such conclusions must necessarily involve one's beliefs in addition to scientific information.

I appreciate the perspective. It could be useful to add or edit the Secretary's Forward to indicate that "**the report has been purposefully commissioned to researchers with alternative perspectives in an effort to address these perspectives and provide some closure for the American public, the US Economy, and US energy policy.**" I suggest this, because it will be very obvious that a handful of skeptics were chosen out of literally hundreds of extremely active and well-respected scientist. Addressing that upfront will immediately reduce criticism, and hopefully generate a positive environment to address these issues. Everyone would like to see convergence of findings on these issues.

It is misleading (cherry picking) to only mention "global greening" (i.e., the influence of elevated CO2 on plant growth) here without also noting the impacts of elevated temperatures and changing precipitation patterns on ecosystem health and agricultural productivity. (This comment is also true of the underlying content on pages 3-7)

It is misleading to state that ocean pH levels are within the range of natural variability. This statement is not true on the timescales governing the impacts of ocean acidification on ocean ecosystems and fisheries. (This comment is also true of the underlying content on pages 8-9)

It is inaccurate to say that "scenarios of CO2 emissions have tended to overestimate observed levels". As the authors themselves note on page 15, it is impossible to reliably predict societal behavior regarding energy production and other sources of emissions, so the climate science community has never provided or suggested that there is a single predictive scenario against which future emissions could be judged. Rather, there has always been a range of projected emissions scenarios, and emissions have always tracked within the envelope of those scenarios. The authors make a reasonable point in the full text about the large number of climate studies that have used RCP 8.5, whereas actual emissions have recently been closer to RCP 4.5. This would be a more appropriate finding to include here in the executive summary.

Stating that climate models "offer little guidance on how the climate responds to elevated CO2" is at best an unjustified (and at worst a biased) judgement given that the authors chose to focus only on the shortcoming and deficiencies of these models, and do not account for the progress made in improving the ability of these models to reproduce many key aspects of observed climate change. The same is true for all of Chapter 5.

Stating that climate models "generally run hot" is an inaccurate and misleading summary statement. A large number of papers and scientific assessments have validated the ability of climate models to accurately simulate many major aspects of global climate. Like all models, climate models are imperfect, and some models have recently shown a tendency to produce higher climate sensitivities--the IPCC AR6 acknowledged this "hot model problem" and there has been considerable discussion in the scientific literature about it and its implications for impacts studies. But this is a relatively recent phenomenon, and there could be other reasons for the recent mismatch (i.e., under-estimates of the reduction in aerosol cooling due to air pollution controls). A balanced scientific assessment would also note, for example, that climate model projections from the 1970s and 80s have accurately predicting the observed trend in global temperature over the past few decades.

It is misleading and inaccurate to say that most extreme weather events do not show long terms trends. It is true that some kinds of extreme events, like tornadoes, have not shown appreciable or detectable changes, and that changes in wildfire must account for changes in land management. But a "summary of the current state of the science" should acknowledge that there is clear evidence that heat waves are becoming more frequent and more extreme, extreme precipitation and nuisance flooding are becoming more common, etc. (This comment is also true of the underlying content in Part II)

It is misleading to state that the aggregate sea-level rise trend in U.S. tide gauges is not accelerating, because as noted in the same sentence, there are significant regional differences in SLR (primarily due to differences in land subsidence), so only looking at domestic gauge data is insufficient to determine if sea-level rise--an inherently global process--is accelerating, with regional factors superimposed on that global trend. There is clear evidence that global sea level rise is accelerating. Moreover, while land subsidence has offset sea level rise in some U.S. regions, the opposite is true in others, and in those regions the combined effects of sea level rise and subsidence are posing considerable risks to people and infrastructure in those regions...but this important context is omitted. (The same is true of the sea level rise analysis on pages 75-78.)

Most of the sentences in this paragraph are not factual in nature. For example, the call for uninterrupted observations of the global climate system is a research recommendation that is outside the remit of this report and is not supported by text within the document. The same sentence also implies that current emissions scenarios are not "realistic", which is not a defensible statement given that historical emissions have tracked within the envelope of previous emission projections (see comment above and Figure 3.2.2), and argues that climate models should address biases and uncertainties, which is something that every climate modeling team already spends an enormous amount of time doing. If retained, the non-scientific portions of this paragraph should be moved to the Preface, and the misleading statements should be corrected.

- There is a disconnect between the stated intent of the report by the authors in the preface, "whether carbon dioxide emissions endanger the U.S. public," and this executive summary, as well as the organization of the report. Would suggest aligning by either reframing the focus in the preface to better match the organization of the report, or better aligning the report organization with the stated focus.

This would be more correct: "Some scenarios of future CO2 emissions overestimate observed levels"

This would be more correct: Some detailed climate models run "hot" in their simulation of the climate of the past few decades, while many have underestimated the amount of surface warming. **More research is needed to select the most skillful and reliable models.**

Need reference for: "Claims of increased frequency or intensity of hurricanes, tornadoes, floods, and droughts are not supported by historical data." **More research is needed to identify how frequency and intensity of weather events are affected by environmental conditions.**

The greening effect can't be due to increase in CO2 alone. The fertilization effect needs Nitrogen and phosphorous. There is observational[1] evidence that nitrogen available to plants is decreasing with increasing CO2 and thus acting to limit fertilization effect. Secondly, the greening of some portions of the planet are likely due to changes in precipitation patterns associated with changing climatic conditions and not entirely attributable to fertilization effect. We will also need to consider that the temperature effect on photosynthesis will also be a limiting mechanism to any continued greening due to fertilization in the future [2]. I recommend acknowledging that the greening is not attributable to CO2 alone and continued greening in the future with increasing CO2 is uncertain.

Any increase in greening from additional CO2 will likely be more than offset by longer growing seasons with higher temperatures (i.e., more days over 100). This causes serious risk to the US midwestern "bread basket", which is why this has been considered a national security concern. The data supporting this trend has been published a few times now in Science by a fellow out of Stanford, among others. Sources: Lobell and Field. 2007, doi: 10.1088/1748-9326/2/1/014002; also Sloat et al. 2020, doi:10.1038/s41467-020-15076-4 has a thorough introduction with many citations.

My understanding is that the upper bound of emissions (i.e., very high) is business as usual, with no mitigation, and a high-end of growth. It's a worst, yet plausible, case scenario. Similarly, the lowest emissions (i.e., very low) scenario includes technologies that don't yet exist including significant CO2 removal. Both are there to understand plausible extremes. The more likely scenarios are the low to high scenarios. I'm sure researchers are open to revisiting how things are bounded and presented if there are reasonable objections.

This may be true. But data driven analyses may not capture anticipated feedbacks.

I am a big fan of data and meta-analysis. However, I'll note that I do not think forest management practices are overlooked. In fact, it is one of the main driving story lines (i.e., too much fuelwood exists, so we need to harvest more forest). There remains strong debate on the efficacy of fuelwood removal over large areas as an effective means of wildfire reduction. The key to this analysis is to look at whether the extent (not the numbers) of fires has grown and whether the burned areas are returning to forests or returning to brush or grassland. If it's the latter, then that is more likely due to regional changes in climate (temp and precipitation).

I think we can point to sea-level rise in south florida, crop damage, and disease vectors fairly easily. I'm not aware of what costs are associated with these 3 items, but I suspect it's quite large.

Yes, this is an important paragraph. Most scientists are not activists. Just because there is a science finding, scientists don't expect that to turn into policy. The public and the democratic system can determine the policy using an evidence-based risk/benefit analysis, just as noted here. Opening up this conversation (which should have been done decades ago) would be welcome and would better engage the public on this issue. DOE could in fact take a leading role here and help address a major national issue.

I am glad we acknowledge that CO2 causes warming, though I am not sure why we are not considering other energy sector related emissions that also cause warming (CFCs/HFCs, methane, aerosol effects). It is not clear what is meant by "CO2 emissions have tended to overestimate observed trends". This is not correct, the cumulative emissions track the RCP8.5 scenario the closest [3,4]. While there was a decrease in emissions during the Covid-19 period the emissions are now back on track following RCP8.5. This statement should clarify what is meant by the statement in the text.

This paragraph seems to be talking about equilibrium climate sensitivity. The most recent model inter-comparison gives a wider range of climate sensitivities in the range of 1.8 to 5.7 and is wider than estimates of climate sensitivities that have been estimated from paleoclimatic and more recent historical datasets, in the range could be in the range of 2.3 to 4.5 K[5]. The higher sensitivity in the most recent model assessments are an area of active investigation by modelers, the feasibility of a higher range needs to be confirmed with additional observational and process scale modeling [6]. The range for climate sensitivities, 1.8-2.7, should be corrected to the most recent findings, 2.3-4.5. We should also recognize that the (a)earth is a complex system of systems (b)we can't do an experiment to find out what is correct or reduced the uncertainty. As we get better at extracting temperature records and CO2 concentrations from paleo data, there will be better reconstructions and possibly narrower range along with improvements in modeling that resolve many physical, biological and chemical processes as we get to higher and cloud resolving spatial resolutions.

This is again not true for the majority of the models in the AR6. Figure 7.19 in the IPCC AR6 report shows that majority of the models exhibit a range within the observational range. The models running hot in the most recent decades are the ones with climate sensitivities greater than 5K, which are 3 out of a total of 40 and it should be noted that about the same number of models also are running cold' compared to observed[7]

There is very little dispute that these are challenging issues for climate science. Many of the issues identified here related to compound effects (subsidence and sea-level rise; wildfire and forest management) that requires further scientific investigation. Identifying further research avenues/focus areas will be helpful to firm up some of the challenges faced in using the climate projections as discussed in the paragraph 9.

The focus on uncertainty is relevant and should be emphasized in future research. However, as a reasonable policy pathways, making decisions in the face of these uncertainties is critical. Most of the climate uncertainties that we have confidence at this time can be considered Level 2 and Level 3 uncertainties, approaches such as scenario analysis (and others) are suggested and followed for making hazard and risk estimates for future conditions. For deep uncertainty hazards (for example tornadoes etc.) that we don't have enough data or capability to model, developing methods for assessing the uncertainty and risk should be a focus of future research.

- “While the report is intended to be accessible to non-experts, we have omitted some introductory or explanatory material that can easily be accessed elsewhere. Nor have we attempted to survey entire literatures related to the topics covered.”
 - While this approach is understandable given the time constraints this report was produced under, given the level of visibility this report will have, it potentially opens up the report to criticism of being misleading by omission – focusing on areas where the authors disagree with the broader scientific literature and omitting areas large areas that are known about climate where there is less disagreement.
-
- “We have focused as much as possible on literature published since 2020 and referenced previous IPCC and NCA assessment reports. We have also used data through 2024 where possible.”
 - This is a good approach for this report; however, much of the literature cited is from pre-2020, and many of the references to IPCC and NCA reports are to older reports (e.g. NCA 4 and IPCC AR 4&5) instead of NCA 5 and IPCC reports from the 6th assessment cycle. In these sections that heavily cite this older literature, it is not clear if the newer literature does not address these points or if the issues raised in the older literature are resolved or significantly changed in the newer literature.
-
- The IPCC’s mandate is to assess the scientific literature comprehensively, including a range of views, and report disagreements. This report has a tendency to selectively quote the IPCC focusing on disagreements and excluding areas where scientists agree. This approach risks giving a misleading picture of the state of the science through omission. A possible way to address this is to have a section summarizing areas of agreement by pointing to all statements of fact and high confidence statements from IPCC AR6 before digging into the more controversial areas with disagreement. This could be a short discussion about areas of agreement up front before explaining the focus of this report on these areas of disagreement, with the IPCC statements of fact and high confidence statements included in an appendix.
 - The report also includes only a very small subset of the literature since 2020. The authors should describe the approach to how they selected the literature included, and if possible, point to any systematic ways excluded literature could lead to different conclusions.
 - It is also notable that there a fair amount of new analysis in this report done by the authors that has not yet been peer reviewed. It would be good to flag these spots with a note about why these are needed while other related peer reviewed articles are excluded.
- The focus on "whether carbon dioxide emissions endanger the U.S. public" is slightly different from the stated focus in the Foreword on "the Nation's climate". This distinction is important because climate change and its impacts outside of U.S. borders can endanger the U.S. public both directly (e.g., global processes affecting sea level rise in the United States, or the impacts of climate change on overseas military installations) and indirectly (i.e., impacts of climate change on international agricultural production and trade dynamics).
- If the goal of this report is to discuss links between CO2 and climate, this is an odd way to begin the report with a focus on differences between CO2 and criteria air pollutants.

- “It is an issue of concern because of its potential effects on the global climate, a complex issue that occupies much of this report.” Suggest deleting the word potential as the existence of the effect is not disputed.
- “But the primary concern about CO2 is its role as a greenhouse gas (GHG) that alters the earth’s energy balance.” Would add “warming the planet” to the end of this sentence as the qualitative effect is not debated and the concern is not about the energy balance itself but about the warming effects of the imbalance.

This chapter points out that the primary concern around human emissions of CO2 is that it changes the radiative balance of the atmosphere. The chapter also makes the points that CO2 has no toxicological effects on humans at current atmospheric levels, and that CO2 is necessary for photosynthesis, with the potential to stimulate plant growth at higher concentrations. These points are well-established scientific facts.

This is all generally correct. I would like to think this is common knowledge.

The reason CO2 was identified as a pollutant is the health consequences of increased temperature, flooding and other climate impacts and not because of any direct affect on health due to exposure to atmospheric CO2[8].

- “The recent decline in pH is within the range of natural variability on millennial time scales. Most ocean life evolved when the oceans were mildly acidic. Decreasing pH might adversely affect corals, although the Australian Great Barrier Reef has shown considerable growth in recent years.”
- Note that the IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (SROCC) finds that pH decline has very likely emerged from natural variability. SROCC SPM A.2.5, “Open ocean surface pH has declined by a very likely range of 0.017–0.027 pH units per decade since the late 1980s, with the decline in surface ocean pH very likely to have already emerged from background natural variability for more than 95% of the ocean surface area.”

It would be more reflective of the literature to say here that “Many studies over the past decade have confirmed a global greening pattern (increase in LAI) attributable to rising CO2 levels, changes in climate, changes in nitrogen deposition, land cover change, and other factors.”

While the Zhu et al. 2016 study did find that 70.1% (+/- 29.4%) of the integrated global change in LAI was attributable to changes in CO2, most of the observed change in LAI is concentrated in the tropical forest regions of South America, Africa, and Indonesia, as shown in this report's Figure 2.1, reproduced from Zhu et al. In terms of the fraction of global land area influenced most by the factors considered in Zhu et al. 2016, effects on LAI from changing climate is the dominant factor (positive effects for 28.4% of land area plus negative effects on 4.7%, for a total area with dominant climate influence of 33.1%), followed by other factors not represented explicitly in the study or in the models (positive effects on LAI from other factors over 25.0% of land area plus negative effects over 5.3%, for a total area with other factors as the dominant effect of 30.3%), and then CO2 effects (23.2% of land area dominated by positive effects and 0.1% with negative effects, for a total of 23.4%). Since the report is focused on the U.S. it would be helpful to indicate the global context for the numbers reported.

The dominant factors related to changes in LAI over the U.S. (including Alaska) are land cover change (+LCC), climate effects (+CLI), other factors (+OF and -OF), and then CO₂ (+CO₂). There are more extensive regions where +CO₂ is the dominant factor for the boreal forest zone in Canada. Given that, and given the explicit focus of the report on the U.S., it would be useful to call out in this section that the global greening pattern has distinct geographic variation in driving factors, and that over the target geography of the U.S. it seems to be dominated by factors other than increasing CO₂. See my powerpoint slide 1 for a zoom on the area of interest from the Zhu et al. 2016 paper.

"Zheng et al. (2017) confirmed the pattern..." should be modified to say "Zhen et al. (2017) confirmed the pattern of observed greening..." As written, and in the context of the previous paragraph, the current text suggests that Zheng et al. 2017 confirmed the pattern of greening being related mainly to increased CO₂, a factor which was not examined in their study.

Without the addition of the appropriate context in the preceeding paragraph (my notes 3, 4, and 5) the remainder of this paragraph could be misinterpreted to mean that the impacts of greening on air temperature are attributable back to a positive effect of CO₂ on greening.

The sentence "Greening has been observed globally but Chen et al. (2019) noted that China and India were exploiting it through land management changes" is misleading. Chen et al. (2019) shows that China and India are *causing* the observed greening in large areas by intentionally increasing the land area occupied by forests (China) and increases in harvested area in croplands, mainly due to multi-cropping, heavy fertilizer use, and expanded irrigation (China and India). It would be more accurate to say "Greening has been observed globally, but Chen et al. (2019) represent a disproportionate fraction of the total greening trend, accomplished through intentional increases in forest area (China), and expansion of multi-cropping, fertilizer use, and irrigation in croplands (China and India)."

Given the changes suggested in these two paragraphs, it raises the question of whether the section title (CO₂ and global greening) is itself misleading. That title wrongly suggests that all the content to follow is reflective of a direct influence of CO₂ on the observed global greening trends, when the literature presents a much more *nuanced* interpretation.

It would be worth noting in this section that the multiple remote sensing-based datasets for LAI changes have remarkable differences, especially when comparing the figure 2.1 reproduced from Zhu et al. 2016 with the MODIS-based data record from Chen et al. 2019 (their figure 1). That means it is difficult to place a lot of confidence in the attribution of changes in different regions to different factors. See my powerpoint slide 2 for a comparison of the four datasets used in Zhu et al. 2016 and Chen et al. 2019.

The authorship for the Haverd et al. 2020 citation is not correct in the References section for this chapter. The correct citation is: Haverd, V., B. Smith, J. G. Canadell, M. Cuntz, S. Mikaloff-Fletcher, G. Farquhar, W. Woodgate, P. R. Briggs and C. M. Trudinger 2020. Higher than expected CO₂ fertilization inferred from leaf to global observations. *Global Change Biology* 26(4): 2390-2402 DOI: <https://doi.org/10.1111/gcb.14950>.

A more recent and more comprehensive study (Keenan et al. 2023) reexamined multiple data constraints and concluded that the model used in Haverd et al. 2020 (CABLE) is outside the likely range of CO₂ fertilization effects, with higher CO₂ fertilization than indicated by multiple independent constraints. At the very least this more recent paper should be cited in the report, with text indicating that the central estimate from the TRENDY models is more likely to be in the range of observational constraints than is the CABLE result. As it stands the report suggests that the latest science indicates that most models have underestimated the fertilization effect, and the literature doesn't currently support that conclusion. See: Keenan, T. F., X. Luo, B. D. Stocker, M. G. De Kauwe, B. E. Medlyn, I. C. Prentice, N. G. Smith, C. Terrer, H. Wang, Y. Zhang and S. Zhou 2023. A constraint on historic growth in global photosynthesis due to rising CO₂. Nature Climate Change 13(12): 1376-1381 DOI: 10.1038/s41558-023-01867-2.

Zheng not in reference list. Do you mean Zeng?

"nearly double"? Why exaggerate? - it's a 76% increase. And they reported that future increase in GPP in extra-tropical northern hemisphere (e.g., U.S.) for [CO₂] from 300 to 600 ppm of 42%, or 32% more than previous predictions (not 76%).

Zhu also explained that "other factors (OF) not considered in models, such as forest management, grazing, changes in cultivation practices and varieties, irrigation and disturbances such as storms and insect attacks, can be a cause of mismatch between observed and simulated LAI trends.... OF contributes the most to the observed LAI trend over 25.0% (increase) and 5.3% (decrease) of the vegetated area"

This report is meant to focus on the U.S., so it would be more appropriate to discuss greening trends in the U.S. Fig. 2.1 shows that the trend in LAI across much of the U.S., with the exception of the Southeast, is less than the global average. Negative OF effects are mainly found in northern high latitudes (e.g., Alaska). Growing season warmth is the dominant factor in cold regions (10.1038/s41558-018-0258-y). Significant browning also is occurring in parts of Alaska (10.1175/EI-D-20-0025.1)

Greening of the Arctic can have important adverse consequences. Direct observations support that warming and increased vegetation cover in the Boreal-Arctic region showed a "robust increasing trend of CH₄ emissions (+8.9%) with strong inter-annual variability. The majority of emission increases occurred in early summer (June and July) and were mainly driven by warming (52.3%) and ecosystem productivity (40.7%)." Yuan et al 2024 (10.1038/s41558-024-01933-3).

Wang et al 2020 (10.1126/science.abb7772) report a decline over recent decades in CO₂ effects on global photosynthesis, possibly due to increasing nutrient interactions or interactions with effects on LAI. The two summary statements provide an extremely narrow and misleading view of "direct impacts of CO₂ on the environment" (the chapter title). The most obvious direct impact of CO₂, in the context of climate change, is its role in global radiative forcing, which is covered in the next chapter but not mentioned here. The chapter should at a minimum be retitled, but better yet combined with the following chapter which discusses other aspects of increasing CO₂.

It is misleading to state that ocean pH levels are within the range of natural variability. This statement is not true on the timescales governing the impacts of ocean acidification on ocean ecosystems, fisheries, and other processes.

The summary should note that CO₂ fertilization effects are limited by the availability of other critical plant nutrients (Nitrogen, Phosphorous). The fertilization effect is also constrained by higher temperature affecting photosynthesis[9]

The attribution of greening to primarily CO₂ fertilization is questionable as per recent research. Specifically, in the Northern high latitudes (where the greening is most prominent), increases in temperatures due to climate change has been shown to provide the best explanation[10]. In the sub-Saharan region reforestation efforts and changing precipitation patterns lead to some of the observed greening (the great green wall) [11]. The attribution to CO₂ fertilization should be qualified by noting the other present and emerging constraints.

This summarization ignores the compound effects of temperature, moisture and CO₂ fertilization in assessing the impacts of CO₂ increase on vegetation and crop yields. Many studies have shown that [12]. Making note of these compensating factors is required for a balanced view on CO₂ fertilization effects on plants.

The attribution analysis in the Chen et al. 2024 paper is very sensitive to the choice of multiple regression vs. partial correlation (their figures 3 vs S7, respectively), and in both cases they seem to misattribute the significant increases in LAI in India and China to CO₂ rather than to changes in landcover and cropping practices. This kind of regression-based attribution is less informative than the model-based inference used for example by Zhu et al. 2016. The description of sensor calibration issues in this paper is also very cursory. It would probably be better to refer to the review paper of Piao et al. 2020 to support the claim of continued greening. Piao, S., X. Wang, T. Park, C. Chen, X. Lian, Y. He, J. W. Bjerke, A. Chen, P. Ciais, H. Tømmervik, R. R. Nemani and R. B. Myneni 2020. Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth & Environment* 1(1): 14-27 DOI: 10.1038/s43017-019-0001-x.

The section 2.1.2 (Photosynthesis and CO₂ levels) lacks coherence. It seems to be a collection of uncontroversial points about the ability of increased CO₂ concentration to stimulate plant growth and reduce water loss through stomates. Figure 2.2 seems intended to show that the current levels of CO₂ are not high compared to levels experienced in Earth's deep geologic past. A balanced treatment of the topic might call out the rate of change in CO₂ concentration from human-caused emissions, and how both that rate and the absolute levels diverge radically from rates of change and ranges of variability over the past hundreds, thousands, and hundreds of thousands of years. See my powerpoint slide 3 for an example set of figures that might be included.

Much of this information in this section is appropriately covered in chapter 9 on agriculture. Chapter 2 purports to have a wider focus (global and environment responses), so it should not dwell on responses of individual crop plants (crops account for only 15% of global GPP). There has been plenty written on ecosystem response to eCO₂, but evidence such as Fig. 2.3 is not relevant. All analyses of direct eCO₂ effects reveal an overriding interaction with other environmental factors (nitrogen and temperature), and global models show that eCO₂ only partially compensates for the negative effects of climate change on ecosystem carbon response.

Zhang, 2025 (10.1016/j.ecolind.2025.113762) reported that although vegetation cover was increasing, resilience against external disturbances (drought, heatwaves, wildfires) has simultaneously declined.

Why use studies based on satellite sensors? I'm quite sure there are studies (again, out of Stanford) that use historic yield data (county-scale USDA data and global FAO data) with associated temperature and precipitation data. Those analyses do not agree with your findings here, I think. But, we can certainly pose these questions to our scientists that study and develop land models and develop a report based on all available data. I'll also note that for sometime data synthesis wasn't considered fundamental science and was therefore largely ignored. Enabling the researchers to conduct robust data analyses on this and other topics would be largely welcomed.

Chen et al. (2024)[13] conclude that "In the attribution analysis of LAI trend and growth rate trend, there were some differences between multiple linear regression model and partial correlation analysis at high latitudes in the northern hemisphere. Partial correlation analysis showed that LAI trend and growth rate trend in this region were mainly affected by temperature and radiation, while multiple linear regression suggested that CO₂ was the dominant driver. The results of the partial correlation analysis seem more reasonable because some previous studies have shown that vegetation change in the high latitudes of the northern Hemisphere is mainly positively affected by climate warming. Multiple linear regression model performed relatively poorly in this region and might not fully identify the contribution of individual drivers. In addition, our model does not include the contribution of land management, which has been highlighted in some previous studies. Of course, the contribution of land management to LAI trend and growth rate trend may be implied in other drivers, such as the positive contribution of CO₂ to LAI trend in PP and PN areas".

The graph is not relevant to the discussion. It should only, at most, cover the time period since the last glacial maximum, which occurred ~20 kyr BCE, and common practice is to concentrate on the last 1000 years.

All data is interesting and useful for perspective and understanding. But isn't the main question whether we are driving up CO₂ and temp at a rapid pace in the post-industrial era? This graph does not help answer that question.

It is misleading to show (only) a graph of CO₂ levels over hundreds of millions of years. These timescales are generally not relevant to processes and issues surrounding anthropogenic CO₂ increases. For example the authors in multiple places make statements about CO₂ levels during the last glacial maximum, but those fluctuations are undetectable on a graph with such a long timescale.

A simple calculation will show us how misleading this figure is. It shows that over 600 Million years CO₂ changed by around 6000ppm. That is a rate of change of 0.00001/yr; CO₂ changed from 275 to 430 in the past 175 years, that is rate of change of 0.88ppm/yr. A change in the rate of over 10⁵ in magnitude - there lies the problem. We are seeing a rapid increase in CO₂ that has never been witnessed and the ecosystem and human society (which is at best 10K years old), may not be able to adapt to this rapid change.

This again suffers from not acknowledging the effects of compound and often compensating phenomena that is occurring with the on-going climate change. For example [14] note that much of the Midwest any benefit from CO₂ fertilization will be negated by temperature increases. [15] note that "Our results show that long-term physiological acclimation dampens the initial stimulation of plant net carbon assimilation to elevated [CO₂], and of plant water use to warming."

The report suggests that the topics of CO₂ fertilization and increased water use efficiency are not adequately treated in recent IPCC reports. In IPCC's AR6 WG I report there is an entire chapter devoted to an assessment of carbon cycle processes (Chapter 5, Global Carbon and Other Biogeochemical Cycles and Feedbacks). This is precisely where one would expect to find information about CO₂ fertilization and water use efficiency. In the Executive Summary for the chapter is the following bolded headline: **Increases in atmospheric CO₂ lead to increases in land carbon storage through CO₂ fertilization of photosynthesis and increased water use efficiency (high confidence)**. The associated Executive Summary text includes reference to two figures, three chapter sections, and a cross-chapter box all dealing directly with this topic. The Executive Summary text also references the important topic of nutrient dynamics and their interactions with fertilization and ecosystem-climate feedbacks, a critical nuance of the interaction of CO₂ and vegetation productivity that is ignored in the report reviewed here.

Cheng did NOT report "increased plant growth"; they reported increased GPP. The distinction is fundamentally important, has been recognized for decades, and is completely ignored in this chapter.

Stating that increased aridity in drylands will not lead to decreased productivity seems a bit questionable. We can point to precipitation gradients from high to low precipitation that supports forests to grasslands to shrublands, respectively. Less water in general will mean less productivity; this can be seen in different forest compositions as well. The authors cite Zhang et al. (2014), but Zhang's main point is that the direct CO₂ effect may be causing greening in drylands that offsets the decreased precipitation. This may be more specific to dryland vegetation, and Zhang et al. specifically state that this finding is relevant to north-east Brazil, Namibia, western Sahel, Horn of Africa, and central Asia.

Section 2.2.2 (Coral reef changes) has a strong focus on past examples of early investigations that are later updated. There is also a note about the difficulty of publishing negative results, with a quote that this pattern "is true across all of science". Given the objective of this report, I suggest a shift in tone away from complaints about the kinds of papers that do and don't receive media attention, and instead highlight the latest synthesis of results about the likely future of coral reefs. A relatively recent synthesis paper (Cornwall et al. 2021) in a well-respected journal (PNAS) based on collated data from 233 locations worldwide projects that 37% of reefs will stop accreting and thus fail to keep up with sea level rise by 2050 under the low emissions RCP2.6 scenario, and no reefs would be able to accrete at a rate to stay ahead of sea level rise by 2100 under either RCP4.5 or RCP8.5. This analysis finds that most of the loss of net accretion rate is due to bleaching events, and a minority of the impacts are due to "direct physiological impacts of ocean warming and acidification on calcification or bioerosion". C.E. Cornwall, S. Comeau, N.A. Kornder, C.T. Perry, R. van Hooidonk, T.M. DeCarlo, M.S. Pratchett, K.D. Anderson, N. Browne, R. Carpenter, G. Diaz-Pulido, J.P. D'Olivo, S.S. Doo, J. Figueiredo, S.A.V. Fortunato, E. Kennedy, C.A. Lantz, M.T. McCulloch, M. González-Rivero, [...] & R.J. Lowe, Global declines in coral reef calcium carbonate production under ocean acidification and warming, *Proc. Natl. Acad. Sci. U.S.A.* 118 (21) e2015265118, <https://doi.org/10.1073/pnas.2015265118> (2021). acidification is not a misnomer...the word literally means to make something more acidic, which is what is happening. The authors may choose to argue that the term is misleading since the oceans are slightly basic, but that is not the same thing as a misnomer.

It is misleading and largely irrelevant to discuss ocean pH levels tens of thousands or millions of years ago, because that timescale is not relevant to the impacts associated with anthropogenic ocean acidification on ocean life. It is well documented that recent changes in ocean pH are happening ~100 times faster than natural fluctuations prior to human CO₂ emissions, and marine species must adapt to changes in conditions on timescales of a few of their lifetimes, which is clearly not millions of years.

The fallacy of these arguments in terms of pH changes and evolution is that the changes described in this paragraph occurred over millennia and the evolutionary process had time to adapt; here with climate change we are talking about changes in our lifetime, the speed of change is the issue. This will lead to collapse of some ecosystem and by the time adaptation takes place it will be millions of years and in a different climate - not the one we have now and that enabled the rise of humans.

I do not disagree with this observation of a decline effect, and I have seen it in some of the early biomass energy analyses that appeared in prominent journals and were then corrected over a few years. While it is unfortunate that some try to grab headlines with unsupported assumptions, it does seem that other researchers tend to provide counter arguments and analyses fairly quickly, indicating that the system does in fact work fairly well.

Similar to previous comment--changes in marine life over hundreds of millions of year has little relevance to the survival of marine species on decadal timescales due to anthropogenic ocean acidification. Are the authors trying to suggest that U.S. society would be well served by going back to conditions during the Jurassic?

This material is repeated in chapter 9 where it belongs. Here, it would be more appropriate to focus on how eCO₂ interacts with drought on ecosystem response (e.g., 10.3390/atmos12020212)

It is misleading to separate the photo-chemical effects of CO₂ from the temperature effects. Extreme heat is hard on plants, causing them to lose water, close stomata, etc. It depends on the plant and the degree of heat. **More research is needed to understand how the temperature (mean and variability) affects plant productivity and resilience.**

Again it is misleading to separate the CO₂ and the temperature effects on the oceans. The warming surface waters is having significant impact on ocean fish and habitats due to the overall increase in global temperatures

The misuse of RCP8.5 as business as usual is acknowledged. However, this was never intended to be a no-policy or business as usual scenario but as a high-end baseline scenario. When dealing with interaction of physical systems with socio-economic projections (which tend to be highly uncertain and policy dependent), the development of this high baseline scenarios is intended to explore the uncertainty ranges that represent a high-end scenario[16,17]. As any engineer would acknowledge, this is a prudent approach when dealing with L3 and L4 type deep uncertainties and provide valid envelopes for assessing uncertainties and risks from high impact events/hazards. The rest of the statements here are uncertainties acknowledged by the climate scientists and included in developing projections and discussing the results[4]. Does the IPCC downplay the role of sun in climate change? The IPCC includes sun variability in all its assessments from the beginning and has been acknowledged as a potential but insignificant contributor.

The IPCC probably discounted the results reported in Connolly (2021) because that study included methodological errors that were documented in several subsequently publications, which the authors do not acknowledge. A serious assessment of TSI would not cherry-pick a single study, especially one that has been shown to include serious errors.

The study by Connolly et al. 2021 uses a linear regression approach to estimate the amount of variability in surface temperature trends that could be explained by total solar irradiance time series. Their approach is to first model the temperature trend with TSI as the dependent variable, and then compare the residuals from that fit to anthropogenic forcing factors. This is an obviously biased approach, and in fact a full paper was published the following year (in the same journal) laying out the erroneous methods and the faulty conclusions they produce (Richardson and Benestad 2022). Richardson, M. T. and R. E. Benestad 2022. Erroneous use of Statistics behind Claims of a Major Solar Role in Recent Warming. Research in Astronomy and Astrophysics 22(12): 125008 DOI: 10.1088/1674-4527/ac981c.

It is unclear what the authors mean by "other natural sources of energy imbalance". If the authors are aware of a plausible additional source of energy imbalance they should state what it is rather than making vague statements without providing any evidence that such sources might exist.

This is one of several places where the authors discuss the CO₂ levels at which C3 plants begin dying in the context of the last glacial maximum. This is misleading, because there is no evidence that CO₂ levels would have continued falling below the levels of the last glacial maximum, and irrelevant to the stated focus of this report, because they rebounded to 280 ppm many centuries before human activities began producing large amounts of CO₂.

- "Instead, the IPCC has created various sets of scenarios meant to span a plausible range of possibilities"
- Suggest changing "created" to "assessed" as the scenarios are created by the science community not the IPCC.

The oceans and land surface absorb more like 60% of CO₂ emissions. The 50% figure is repeated multiple times in the ensuing pages and should be corrected.

- “The IEA trends, which are based on plausible extrapolations of current economic and demographic trends, run below the entire envelope of RCP projections, implying not only that RCP8.5 is dubious, but that even lower emission scenarios are unlikely.”
- Burgess compares IEA to the two highest SSP-RCP scenarios, not to the full range. The last sentence of the paragraph is incorrect – the range in Burgess is only no climate policies which doesn’t include the lower RCPs.

Please see above. The business-as-usual may be a misnomer but this is a 95% case that is typically included in any type of prudent uncertainty analysis, in particular given the difficulty of predicting the evolution of socio-economic and political conditions over the next 100 years. It should be noted the difference between RCP8.5 and RCP4.5 for mid-century conditions that is of particular focus for adaptation studies is not significant and most of the divergence in these two scenarios occurs later in the century.

- “As of 2023, global CO2 emissions have been tracking well below SSP7.0 and are even below SSP2-4.5.”
- Note that near-term emissions are not necessarily a good predictor of long-term emissions, as the latter will depend on changes in population, demographics, technology, and policy that do not have the same influence on near-term emissions.

Section 3.2.2 (The carbon cycle relating emissions and concentrations). There is no new information or analysis presented in the chapter, and it relies heavily on Global Carbon Project information. Figure 3.2.3 is described as showing a seven-fold difference in the recent trend in land carbon sink estimates across land models, and this is given as a potential driver of uncertainty in future carbon cycle and climate change predictions. Variation among models is well known and is one of the reasons exercises like CMIP and GCP rely on multi-model ensembles when assessing the state of the science. The distribution of results looks fairly typical in this diagram, with a strong central value around 1.3 to 1.3.

Regarding the uncertainty in the land sink, a primary reason for this uncertainty is that the primary response of photosynthesis to CO2 does not predict C sequestration because of strong interactions between CO2 and climate, nutrients, stand dynamics, and other factors. That is why the almost complete focus in this report on crop plants and plant responses in greenhouses does not inform us about "CO2 uptake by land processes"

Yes, CO2 uptake by land processes have uncertainty associated with them. It's worth noting that DOE researchers (many of whom are retired now) were world-leading and making progress in this area, but nearly all work on CO2 uptake and emissions stopped 10-15 years ago due to political pressure and political concerns. The Carbon Dioxide Information and Analysis Center at ORNL, now cancelled, was a main contributor to the Global Carbon Project. Now the US is woefully behind and ceding expertise on this to the Europeans (as cited in this report). We are also ceding expertise on carbon monitoring via satellites to other countries. There's nothing necessarily wrong with this if this is where the US wants to go, but it's important to recognize that we have lost significant capacity due to policy changes over many years. That the dynamic vegetation models need more development is acknowledged and is a defining challenge for Earth System Modeling for the next decade.

The final sentence on this page is misleading and confusing as written. Most studies of the land carbon sink suggest that the rate at which it is increasing is declining, and several studies suggest that the ocean sink is declining. The authors should also explain the relevance of their statements here, because it isn't clear.

- “In summary, while there is clearly warming in the land record, there is also evidence that it is biased upward by patterns of urbanization and that these biases have not been completely removed by the data processing algorithms used to produce climate data sets.”
- Quantitative info would be helpful here. How large is the warming? How large might the bias be?

As noted, Parker[18,19,20] has investigated the Urban effects on temperature records. The finding establish that by comparing warm and wind conditions world wide that the impact is negligible. However, as noted by authors, this literature hasn't progressed much over the last decade and much more attention needs to be paid to this question by employing newer models, datasets and analytical techniques available to us at present.

There is no quantification of the effects of TSI and volcanic aerosols that would lead to modifications of the radiative forcing shown in the figures

Many of the arguments and references are dated. Further, the challenge posed is to prove a negative, i.e. that the record used by AR6 is not biased by urban heat island effects. On the contrary, while verbally stating uncertainty, the authors cannot make a claim that the AR6 statement is false. Even if one were to accept the Soon, et al value of 0.89 deg C/century, the focus on the land surface warming completely neglects the contribution of the oceans to global surface temperature change, which must be considered in any discussion of surface temperature change.

More accurate to say "because their effect remain largely unknown". Good examples include natural aerosols (sea salt, dust, biogenic aerosols). Are there others to mention? **More research is needed to understand radiative effects of both natural and anthropogenic sources of radiative variability throughout the atmospheric column.**

What scenarios are the models using in this figure?

Vegetation is only one way that CO2 is accommodated by land processes, the other is through soils and microbial processes Microbial dynamics is complex and challenging to understand but may also be a way to boost CO2 absorption by soils (which also enriches the soil nutrients!). **More research is needed to better understand how plants and soil systems remove CO2 from the atmosphere.**

More research is needed to improve the understanding and representation of land systems, carbon and nutrient fluxes in models.

Carbon uptake by oceans and ocean ecosystems is less well studied and modeled. The lack of spread of carbon in ocean models may result from insufficient information, ocean biogeochemistry, ecosystems and model development. **More research is needed to study and model carbon, nutrient and ecosystem processes and dynamics in ocean models.**

More research is needed to map warming trends in urban, non-urban and other environments and to ensure accurate representation of regional and global trends in temperature.

- The Chapter Summary doesn't really capture the full tax of the chapter. Would suggest: Defining ECS/TCS, reporting values for ECS and uncertainties/methods, then reporting values for TCS, preferably with a sentence on what those values imply
- “This raising of the lower bound is less justified; evidence since the AR6 finds the lower bound of the likely range to be around 1.8°C.”
- This isn't really discussed comprehensively in the chapter.

- “The climate’s response to increasing concentrations of CO2 is central to the scientific debate on anthropogenic climate change”
 - Suggest rewording this to, “The magnitude of the climate’s response...” As currently worded it could be a bit misleading since the qualitative effect (i.e. increasing CO2 concentrations lead to increases in global average temperature) is not disputed – the uncertainty is around the magnitude of the effect.
 - The text cites Dayaratna (2016), but this should be a reference to Dayaratna (2017) as listed in the reference section.
 - “If ECS is very high (above 4.5°C) immediate aggressive emission controls become more imperative, whereas no CO2 emission controls are economically justifiable for ECS below 2.0°C (Dayaratna et al. 2016, 2020).”
 - This could use additional clarification as there are low and now carbon technologies that can be economically justified based on cost of energy alone, depending on technology, location, etc.
-
- “Obtaining a precise estimate is impossible, so policy making needs to account for the uncertainty.”
 - The Social Cost of CO2 (SC-CO2) as used by the USG in regulatory impact analyses has always accounted for uncertainties in the ECS. EPA (2016) used a probability distribution for the ECS calibrated to IPCC AR4 findings (the most recent values available at the time) to harmonize the approach for the three models used (FUND, DICE, and PAGE). EPA (2023) updated the SC-CO2 in response to NAS recommendations (NAS 2017) to include a much fuller representation of the uncertainties in both ECS and TCR, and utilized a modular approach incorporating the Finite amplitude Impulse Response (FaIR) climate model, a reduced complexity climate model, with ECS and TCR distributions calibrated to IPCC AR6 to represent this uncertainty.
 - The Dayaratna (2017 & 2020) papers referenced here analyze sensitivity of a subset of the older models used in EPA (2016) to just ECS estimates. It would be helpful to address the advances in the broader literature that are discussed and represented in EPA (2023) as the focus of this report is on literature published since 2020.
 - EPA (2016) “Technical Support Document: - Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis - Under Executive Order 12866 - Interagency Working Group on Social Cost of Greenhouse Gases, United States Government.” https://www.epa.gov/sites/default/files/2016-12/documents/sc_co2_tsd_august_2016.pdf
 - EPA (2023) “Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, ‘Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review’ – EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances.” Docket ID No. EPA-HQ-OAR-2021-0317. https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf
 - NAS (2017) “Valuing Climate Damages. Updating Estimation of the Social Cost of Carbon Dioxide” <https://nap.nationalacademies.org/catalog/24651/valuing-climate-damages-updating-estimation-of-the-social-cost-of>
- The inclusion of a paragraph with high-level summary statements about climate policy here is confusing and inappropriate. This content belongs in chapter 11.
- “Because of concerns about model tuning and the high sensitivity to cloud parameterizations, the IPCC AR6 (2021) did not rely on climate model simulations in their assessment of climate sensitivity, relying instead on data-driven methods.”
 - AR6 WGI 7.5.5 states this a little differently. IPCC’s assessment wasn’t “directly informed” but “ESMs remain essential tools.”

We can all agree that there is a desperate need for developing higher-resolution climate models that can get to cloud resolving scale so that we can represent the cloud feedbacks correctly in the model and avoid uncertainties from parameterization and scale dependency of the parameters. This doesn't make these models useless. The models provide directions for further improvements and necessary changes to reduce the uncertainty bands over the ECS.

- "For AR6, the IPCC placed primary weight on the results of Sherwood et al. (2020)..."
- The IPCC assessed a large number of studies, not just Sherwood (2020).

The point about observed global temperature trends being consistent with both high-ECS high-aerosol and low-ECS low-aerosol worlds is an important one. It would be good to acknowledge that uncertainty in ECS is not just due to uncertainties in climate models, but also the lack of accurate measurements of pre-industrial aerosols and thus anthropogenic changes in aerosol forcing. And this uncertainty cuts both ways--if we live in a high-ECS world that has been "masked" by strong aerosol cooling, reducing air pollution could result in rapid warming (and this may be linked to the "hot model" problem as well).

The point about ECS values derived from past climate states (especially cooler ones) possibly not being totally representative of the current climate is another good point, one that would be good to acknowledge in their earlier and later discussions about uncertainty and the rationales for emissions reductions (e.g., on page 26 and in Chapter 11).

The databased approaches have mostly been robust with changes at the high end around 3-4C and at the low end around 1.8 to 2.5C as noted in the chapter. It will be challenging to make this range shorter given the data requirements and uncertainties in data over paleo/holocene periods. It seems reasonable to estimate that we may have ECS in the range of 3 to 4K (higher than 3.5K being less likely [21]) at the higher end and this should not prevent us from planning and thinking on ways to prevent this possible scenario as it could have major consequences for humans across the planet.

The statement "TCR is more generally related to peak warming and better constrained by historical warming, than ECS." requires further context. While true that the "peak warming" rate achieves a maximum around 70 years, temperature increases continue for several centuries and approaches the ECS value asymptotically. Further, the Lewis (2023) arguments were placed in context by Sherwood and Forest (2024), that identified multiple recent ECS studies that showed, collectively, that the PDF in their original Sherwood, et al paper could not be narrowed. Additionally, the reference to Sherwood et al (2020) is incorrect. It omits many authors and the authors that are listed are not in the correct order.

The relatively shallow treatment on TCR relative to ECS here is puzzling, since the authors acknowledge that it is better constrained, and it is also a more relevant value for projected warming over the next several decades. The first sentence here also describes it as an observational constraint, but it is not, it is a metric that can be produced from either data or models (or both), and climate models are routinely used to run TCR experiments.

"This practice" - need to clarify whether this refers to "emergent property" or discarding. I believe the authors mean the latter, but it should be made more clear

"assumed positive cloud feedback" - this is an emergent property too, based on simulation/observation. Should be rephrased to not use the word "assumed". **Further research is needed to better understand and model how clouds change with temperature and what the feedback effects are on the radiative balance. Further research is needed to narrow the uncertainty on climate sensitivity, both using models and observations.**

- It would be helpful for the reader for this chapter to start with a summary figure (e.g. like IPCC AR6 WG1 – SPM.1 or Fig. 3.4). Currently, the chapter jumps right into details that will be difficult for anyone who is not a climate scientist or already versed in these issues to understand or interpret.

A more accurate title of this chapter would be "deficiencies in climate models" because the authors focus almost exclusively on processes and features that models have struggled to reproduce, rather than a balanced assessment that also indicates what these models do well.

The models range over a factor of three in ECS comparing the most sensitive to the least sensitive models. However vast majority of the models are in 3 to 3.5K sensitivity range and it is probably not correct to assign all models the same amount of importance, given the fact that some of institutes participating in the exercise have different levels of capacity to develop and test models.

- "Figure 5.1 CMIP6 Average surface temperature range across 33 models and standard deviation using SSP5-85 scenario."
- SSP5-85 is an excessively high scenario that is not representative of current expectations of emissions projections. Suggest changing the figure for on with data from a less misleading and more representative scenario.
- There have been extensive published comments and responses on the approach used in Scaffeta (2023). Can these be reflected here?
- G.A. Schmidt, G.S. Jones, and J.J. Kennedy, "Comment on "Advanced Testing of Low, Medium, and High ECS CMIP6 GCM Simulations Versus ERA5-T2m" by N. Scafetta (2022)", *Geophysical Research Letters*, vol. 50, 2023. <http://dx.doi.org/10.1029/2022GL102530>
- N. Scafetta, "Reply to "Comment on 'Advanced Testing of Low, Medium, and High ECS CMIP6 GCM Simulations Versus ERA5-T2m' by N. Scafetta (2022)" by Schmidt et al. (2023)", *Geophysical Research Letters*, vol. 50, 2023. <http://dx.doi.org/10.1029/2023GL104960>
- Z. Hausfather, K. Marvel, G.A. Schmidt, J.W. Nielsen-Gammon, and M. Zelinka, "Climate simulations: recognize the 'hot model' problem", *Nature*, vol. 605, pp. 26-29, 2022. <http://dx.doi.org/10.1038/d41586-022-01192-2>
- S. Jain, A.A. Scaife, T.G. Shepherd, C. Deser, N. Dunstone, G.A. Schmidt, K.E. Trenberth, and T. Turkington, "Importance of internal variability for climate model assessment", *npj Climate and Atmospheric Science*, vol. 6, 2023. <http://dx.doi.org/10.1038/s41612-023-00389-0>

- "Spencer (2024) has also provided a useful summary of the Model-Observation mismatch by comparing trends in surface temperature data products with those in individual climate models, as summarized in Figure 5.3; most climate models show substantially more warming than the observations since 1979."

• It is worth noting that IPCC projected temperatures do not use all models and instead focus on those that best reproduce observations. So while some models may show warmer or cooler temperatures than observations, that does not imply that projections from large assessments reflect those biases. Scaffeta (2023) has been widely criticized in the scientific literature for its methods and conclusions. These results should not be presented without acknowledging these concerns, or better yet, the authors should attempt to survey the literature on this topic rather than presenting results from a single controversial paper.

Spencer (2024) is not a peer reviewed journal paper and is not really appropriate to include here when there are many papers on this topic that have been peer reviewed.

While this analysis for the CMIP6 model set is accurate and focuses on a dataset that has shown higher ECS than previous iterations. It should be noted the models have exhibited considerable skill in predicting the subsequent GMST warming[22]. The criticism of the CMIP6 based high ECS models may be premature and probably will have to wait additional time before passing judgement. As noted in [22] these models have shown considerable skill when evaluated on their projections over the past 30-40 years after the projections were published.

The statement that the signal for anthropogenic greenhouse warming emerges first and most strongly is not supported and is inconsistent with my understanding of climate dynamics. The following sentence is also dubious (and probably backwards) since models run with observed surface temperatures tend to produce much better mid-tropospheric temperature simulations.

In most places the authors correctly describe the discrepancy between models and observations as a mismatch, which is appropriate because observations of mid-tropospheric temperatures are also subject to uncertainty (as indicated in the error bars in Figure 5.4). This is one of the reasons for the IPCC confidence assessment noted in the following paragraphs. However, here and in a few other places in this section the writing implies that differences can be solely attributed to model deficiencies.

A recent study attributes the bias to inter-annual availability and residual bias in the satellite record. The pre- and post 2000 have temperature records from two separate satellites and the discrepancy in the trends needs to be resolved[23]

- “Hence, we assess with medium confidence that CMIP5 and CMIP6 models continue to overestimate observed warming in the upper tropical troposphere over the 1979–2014 period by at least 0.1°C per decade,”

“Notably, despite the accumulation of evidence of excess model warming the IPCC assigns only medium confidence to the existence of a warming bias.”

- This reflects a more general issue with IPCC confidence statements where the inclusion of a quantification reduces the confidence they include. Here the full statement with the quantification of 0.1C per decade is assigned medium confidence whereas a statement about the existence of the bias without a specific quantification would receive higher confidence.

This section should be combined with the previous one as it is really just elaborating on the same discrepancy between climate models and observational estimates of tropospheric temperatures.

This is challenge we need to face and work on improving the parametrizations. The new era of super high resolution models would resolve the complex interactions of turbulence, transport and clouds and will provide less uncertainty[24]. However, this can't be used to invalidate the entire set of models, the models are performing remarkably well given the complex system that is being modeled. It should be noted the models do cover the range of the observations and it is not outside the envelope of model predictions.

- Does this update with data through 2024 use the same models that Christy and McNider (2017) used, or does it use updated versions for the models as well? This would be helpful to clarify here and in other places where figures are produced with updated data.

- “The IPCC AR6 did not assess this issue.”

- Was this assessed in IPCC AR5? Many of the citations here could have been assessed then. Is there any more recent evidence on these issues that should be referenced here?

There has indeed been considerable controversy about this particular metric. It is curious that the authors choose to cite two older papers here (2008 and 2009) rather than the rich recent literature, especially since they went to the trouble of updating on of their graphics with 2024 data. Recommend including a more balanced assessment of this discrepancy, it's potential causes, and recent efforts to address it, as well as noting that climate models are able to accurately capture many other aspects of climate change.

This entire section is very misleading as it cherry-picks a specific result and takes it out of important context. The reason why stratospheric cooling has slowed (and, at least in one study, even warmed) in the past two decades is due to the recovery of the ozone layer. This is clearly stated in the Philipona et al article cited by the authors and in the IPCC report. Additionally, satellite-based estimates of stratospheric temperature (which is data the authors highlight in the previous section) do not show the same trends. This section should be removed or amended to note that stratospheric temperatures are influenced by multiple factors and that a comprehensive analysis of stratospheric temprature trends is fully consistent with those projected under climate warming scenarios.

- Are there more recent papers or data available? All the info here is AR6 or pre-AR6, but more recent years have shown more surface warming. It would be interesting to show stratospheric temperatures for these more recent years as well to see how this relationship holds.

- Trends in snow cover depend on both changes in precipitation and changes in temperature. It would be helpful if these were discussed separately rather than showing the net effect for one region/season. Part of the problem here may be which part of the stratosphere is sampled and the importance of interannual a variabilities in the lower part of the stratosphere. [25] have shown that including the stratosphere as defined from 25 to 50km there is high confidence that the cooling signal caused by CO2 increase is detectable and beyond the noise levels of the natural and inter annual variability in the lowest part of the stratosphere. We should put this debate to rest.

Many observational studies have shown that snow cover is decreasing, and have explained why it is more meaningful and useful to look at spring trends in snow cover, as well as why observations of snow cover can be uncertain.

The authors provide a relatively accurate survey of results regarding hurricane numbers and landfall frequency, which have indeed not shown any trends. But their analysis of hurricane intensity, which has been shown in many studies to increase, is lacking...the chart showing just estimated windspeeds of the top ten landfalling storms is really weak given the analytical talents of the author team. They also do not provide any discussion of the physical basis for why hurricane numbers may not be changing while the intensity of the strongest hurricanes is expected to increase, nor analyze any data in the context of those conclusions. Given the large uncertainties in obtaining data over much of the Northern Hemisphere, using a smaller regional analysis to discount the models may be inappropriate. We should that this remains a question we need to resolve developing better datasets with more seasonal and sub-seasonal coverage to test the model on a global scale

Some of the early work on this topic was done by researchers at Stanford. But there are plenty of other papers. Here's one that came up in a quick search: <https://www.pnas.org/doi/10.1073/pnas.0906865106> Contradicting a number of large meta-analyses on the topic of decreased yields with increased temperature is ok. That's what science is all about. But it's going to have to be done in a more robust way than a couple paragraphs here with one or two citations.

This statement from [26] refers to the fitness of global scale models to resolve regional scale features and argues for developing model at higher spatial resolutions. Using downscaling and other techniques additional insights can be gained at regional scales and provide information for a set of 'purposes'. However, we should work on getting to regional and local scales with GCM as quickly as possible. The analysis presented of the IPCC AR5 figure, Fig 5.6 in the current document, is outdated and should be removed. The issue IS addressed in the AR6, but not graphically, in the text reproduced by the authors. If the section is retained, it should be moved ahead of Section 5.3 to present the material in the correct order chronologically.

The text and figure refer to 33 models, which would include the best and worst. How much does the uncertainty/range change for a subset of best-performing models?

How many models are included in each case?

How are the lower and mid-troposphere defined?

"excess amplification" - amplification of what? And Figure 5.6 shows tropics, is this bias only tropical?

Further research is needed to improve how models represent clouds and dynamics; additional observations are also needed to improve these representations.

Further research is needed to improve how models represent planetary albedo to align with observations.

- This chapter relies solely on observations and does not include other lines of evidence, it would be helpful to discuss what these other lines of evidence indicate to avoid introducing potential biases (e.g. what do process based lines of evidence show?)
- The chapter summary does not fully capture the chapter or the science. Suggest talking about the extremes affected by climate, the extremes not affected by climate, and those where the data isn't sufficient to establish a trend.

- "Over geologic time the climate system has generated an (essentially) infinite variety of weather patterns and extremes..."

- Just because an event occurred before humans does not mean humans are not influencing the frequency or intensity of such events now. Additionally, if something has happened in the geologic record, but humans have never experienced it before, they may not be prepared for such an event.

This is very important. I have not dug into the data on extreme storm data (both numbers and intensity).

Getting researchers together that disagree on a topic, analyze the data, and publish findings together can be very productive. DOE could sponsor a workshop and bring together a broad range of experts and find answers to extreme weather trends.

I conducted some analyses on wildfire years ago. I saw the bias in the university textbooks and in the wildfire working groups, who essentially went by textbook teachings. It was rare that people went back to the original data. The last time I checked the actual data (a good 10 years ago), I found that...you're correct...the number of fires had not changed much. What does seem to be changing is that the extent of fires has grown, and this is mostly attributable to hotter and drier conditions. I do not believe there is more fuel load now than there was before the US was settled. Arguing that harvesting will decrease fire is nonsensical; you have no idea where the lightning is going to strike and subsidizing the thinning of all forest in the western fire zone would be cost prohibitive. I think we need to stop using wildfire management as a justification for harvesting. If we want to harvest more, than do it, and stop using fuel load arguments (which is more of a concept than proven reality) as a reason to do it. The biggest issue here is encroachment of housing communities into the Wild-Urban Interface (WUI) and how we want to handle that with policies. Fire is really only bad if it's burning someone's house down. Otherwise, it's a natural progression in forest dynamics. Sources: Campbell et al. 2012. doi:10.1890/110057.; Schoennagel and Nelson. 2011. doi:10.1890/090199.

Long term persistence is not a recognized property or phenomenon associated with rainfall data. The use of such a statistical technique, especially one that is not widely recognized or used in the hydroclimate community, demands additional justification.

Looking for trends or estimating trends in extreme events in current and future climate is not necessarily an attribution study. Estimating frequency, duration and intensity of events helps understand the changing range of the hazard (if there is one). An attribution would be to find to what extent a particular event is influenced or affected by increase in CO2. I would consider these two not to be the same.

Attributing any single event to changing climate is perilous, agreed.

As noted by [26] there is not yet any strong evidence that LPT is a fact

- The figure heading has “@RyanMaue” in the title. Looks odd to have a social media handle in the figure. Is the source peer reviewed?

- Insufficient data is different than no effects. It would be helpful to be clear in instances like this whether a conclusion is because the data supports it, or because there is insufficient data to say the opposite.
- Could you also show a similar table with hurricane precipitation?
- The last two paragraphs show findings from NCA4. With the interest in focusing on recent literature, what does NCA5 say about these issues? Does it not cover this issue, does it say the same thing, does it contradict this finding?

The authors have cherry picked results throughout this section. For example, they have selected only a few quotes from IPCC and NCA describing extreme heat. The absolute single daily maximum temperature for a given year, for example, is a poor statistic for assessing statistical changes in extreme heat (it would be like comparing the tallest person in each country to try to calculate change in average height). A complete and comprehensive review of the underlying assessments and the data it cites clearly shows that extreme heat has been increasing in North America.

- The heading of section 6.3.1 is a bit misleading as this is partly driven by warmer winters and fewer cold days.

The "climate less prone to extremes" is a misleading statement. If minimum temperatures are increasing faster than maximum temperatures (which is an expectation of climate change based on basic physical principals), then the difference between warm and cold extremes on many timescales is expected to decrease. There are reasonable points to be made regarding observational deficiencies, regional variations in extreme events, and the potential benefits of reductions in extreme cold, rather than coming up with misleading statistics.

The conclusion "The rise in Tmin has been strongly related to the growing presence of manufactured surfaces around the weather stations over the last 100+ years (the so-called urban heat island effect; Section 3.3 and Spencer et al. 2025)." is only found in an author's self-citation of Spencer, et al, (2025) that does not appear in the references at the end of Chapter 6. Further, the trend of greater increases in Tmin relative to Tmax is documented and explained in Easterling, et al (1997).

<https://doi.org/10.1126/science.277.5324.364>; Thorne, et al (2016) <https://doi.org/10.1002/2015JD024584>. The trend cannot be attributed to UHI effects.

The NCA5 is cited here and elsewhere. However, the USGCRP website that hosts the NCA reports has been removed. Hence, there is no way to read and review the referenced findings. This DOE report is therefore arguing against something that no longer exists, and that may appear questionable.

I am not sure how one can conclude that the temperature extremes have decreased based on figure 6.3.4. If I understand correctly the Tmax-Tmin for each year was tabulated and a mean of this difference averaged over the 100+ years was subtracted from these values to get some thing like a anomaly. This anomaly seems to have moved from +3F to -3F over the course of q00 years and now trending upwards again. This suggests to me that there has been significant changes in the extremes over the years and the most recent trend is towards warmer highs?

- Figure 6.3.5 shows the number of days greater than or equal to 95F in 6-year periods. Why show this in 6-year periods? That seems like an unusual choice. Would this look different with 5, 10, or 15 year periods?
- "We measure "Heatwave Days" here as the count of all days in May-Sep each year that exceed the 90th percentile for that day and that lie within a period of at least six consecutive days."
- Why six consecutive days? Would this look different for 5 days, a week, 10 days? Six days seems like an unusual period to pick.
- The dependence on and subsequent choice of reference periods is worrying. Can multiple reference periods be shown to show the dependence?

The NCA5 heading is not a controversial statement, it simply says that things are changing, which they are by the author's own admissions and graphs. It does not suggest pervasive positive trends, as the authors suggest it does.

The NCA5 is suggesting that the western US is experiencing increase in extreme heat (page 57, para 4); it seems like figure 6.3.5 establishes the fact and not contradict. There was no claim of pervasive (all CONUS) increase in the report

- "The NCA5 directs readers to the website <https://www.globalchange.gov/indicators/heat-waves> (USGCRP 2023)"
- Broken link.

The definition of heat wave as consecutive 6 days seems to be a special case. The standard definition is exceedance of 85 or 90 percentile for 2 or more days. I can't check with NCA5 as it was taken down from the web

Figure 6.3.6 establishes that heatwaves as defined here have increased in the west and have shown increasing trend all over CONUS in the past 50 years. [27] shows the results for the last 60 years and there is little doubt that there is an increase over most of the CONUS, again the NCA5 report is off the web; figure 6.3.7 shows the heat waves are monotonically increasing from the 1960. Including the longer data is only interesting in the dust bowl 1930 camouflage the signal somewhat for the Midwest and central plains. The problem of taking averages over large regions (for example entire CONUS) is that it kills the peak signals and smooths out the emergent climate change indicators. Doing continental averages to look for extremes is really not very useful. Secondly, there is no particular reason to assume that the climate change signal (i.e. the number of heat waves) should showing up 100 years ago, as the CO2 in the atmosphere started to be of significance in the last 50 years.

- This reads as a very selective analysis.

The statistical techniques used in the 2019 paper are highly non-standard for evaluating precipitation statistics, and the 2024 results have not been peer reviewed. The results reported in the NCA and IPCC and the underlying studies have clearly demonstrated that extreme precipitation is increasing.

The relevance of the paragraph on 19th century and paleoclimatic rainfall events is unclear. If anything, the fact that such extremes have occurred in the past argues for more aggressive adaptation measures to ensure that damages won't be as high if they happen again (even notwithstanding potential changes in risk due to climate change). Note, however, that additional context is provided in the box that follows two pages later.

averaged over all stations in the region, none of these trend parameters is statistically significant'. This is again how one would be looking at statistics of extremes in precipitation. As we know, precipitation is highly heterogenous and location dependent. As observed there are trends even with the approaches chosen by the authors of this study compared to NCA5 and averaging that over and saying there is no trend seems disingenuous.

The choice of looking at (only) 5-day precipitation totals is questionable. Most rainfall events, including the most extreme and damaging precipitation events, are typically on a daily timescale.

same argument as above

again the issue here is that we are talking about events that are happening in a short-time and over a small region. Averaging over time and space and develop statistics is probably not appropriate when we are talking about events such as the recent flooding event in Texas. If you average that event over 30 days, it will seem unremarkable.

As noted in NCA5 and shown in the analysis presented in figure 6.4.4 there is a clear trend of increase in extreme precipitation in 12 of the 27 stations analyzed. Again averaging over the entire region is not the best method for picking extreme precipitation events that are local both in time and space.

- Is this section suggesting that climate change is causing the decrease in tornadoes?
- This is another place where it would be better to reference newer literature and NCA5 instead of relying on NCA4. What has changed since then?

Perhaps this is more germane to the sea level rise section, but nuisance (high tide) flooding due to sea level rise has been observed across much of the united states.

- Figure 6.7.1, is this the same indicator used in AR6?

- “In summary there is no evidence of increasing drought frequency or intensity in the U.S. or globally over recent decades.”
- This statement is not supported by the citations presented here. AR6 finds that there are trends in agricultural and ecological droughts and this section does not contradict that finding, so this general statement about there being no evidence of increasing drought frequency or intensity is misleading.
- AR6 SPM A.3.2 – “Human-induced climate change has contributed to increases in agricultural and ecological droughts in some regions due to increased land evapotranspiration (medium confidence).”
Footnote, “Agricultural and ecological drought (depending on the affected biome): a period with abnormal soil moisture deficit, which results from combined shortage of precipitation and excess evapotranspiration, and during the growing season impinges on crop production or ecosystem function in general (see Annex VII: Glossary). Observed changes in meteorological droughts (precipitation deficits) and hydrological droughts (streamflow deficits) are distinct from those in agricultural and ecological droughts and are addressed in the underlying AR6 material (Chapter 11).”
- “The IPCC has not provided an assessment of wildfires.”
- AR6 WGII SPM discusses wildfire, and WG1 SPM discusses fire weather.
- AR6 WGII SPM – B.1.1, B.1.4, Fig. SPM.2, Fig. SPM.3, B.5.2, B.6.2, C.2.3, C.4.2, D.4.2,
- AR6 WGI SPM – C.2.4

However, there is evidence that flash droughts, which last weeks have increased over the Northeast and Midwest [28]

More recent analysis indicate forest loss due to wildfire accelerating over the recent past[29]. In particular there has been an increase in extreme fire radiative power with a 2 fold increase from 2003 to 2023 (period of satellite data availability)[30][31]. However, that these fires are not out-of-sample and there is a fire deficit as shown in figure 6.8.2 should be acknowledged. But this doesn't alleviate the concern that wildfire is leading to lot of socio-economic concerns and needs to be addressed[32]

The question of forest management and its role on wildfire frequency and intensity is interesting and deserves further study.

Patterns and variability of precipitation are poorly understood and models are particularly challenged to represent and predict precipitation, further research in this area is needed

The greater concern with hurricanes is the amount of moisture they carry and drop after landfall, i.e. flooding. While this is an active area of research, it is worth noting that this may be the bigger issue for hurricane change with warming

Combining heat and humidity as a measure of extreme heat is not unusual since it is this combination that is most impactful to human health - the body cannot cool if the humidity and heat are both high. Since a warmer atmosphere holds more moisture, and CO2 causes heat (and therefore moisture) to increase, this combination is a relevant assessment of the effects of CO2 on extreme conditions that make it difficult for people. Also combining with urban regions is helpful since that is where most people live. These points are not made in the report and are important.

The concern for many are the extreme hot temperatures without night-time cooling that provides an opportunity for recovery. This section doesn't address this. In this regards, the focus on the difference between extreme hot and extreme cold is misleading when generalizing about decrease in extremes (figure 6.3.4) - the decrease is because we don't have as much extreme cold.

Given the satellite observed ice sheet dynamics, I'm not sure that ice sheet retreat is that hypothetical.

- "The largest sea level increases along U.S. coasts are Galveston, New Orleans, and the Chesapeake Bay regions"

- How much? Are there quantifications?

The global mean sea-level was increasing at a rate of 0.08 inches/yr in 1993 compared 0.18 inches/yr in 2023, almost double the rate[33], which is approximately what is needed to meet the NOAA projections by mid-century and seems plausible.

This is the first time I have seen all sea-level rise estimates discredited by an offset from land subsidence. This is certainly a global dynamic that we would want to capture and understand in Earth system models, and DOE can look into this further with analyses and experiments if necessary.

The authors rightly describe the extreme uncertainty regarding sea level rise. It would be helpful to add some information about the complex, highly nonlinear and uncertain processes involved with ice-sheet dynamics and ice-sheet - ocean interactions that are especially key for Antarctica. **More research is needed to study and model ice sheet dynamics and ice-sheet ocean interactions.**

- "AR5 Working Group II (Cramer et al., 2014) makes the following statement:"
- What does AR6 say on this topic? The science on attribution has evolved since AR5.
- "There are three areas of substantive criticism of the IPCC's assessment of the causes of the recent warming..."
- Are there citations that can be included here or are these criticisms attributed to the authors?
- "IPCC's policy-informing..."
- This phrasing is a bit odd. IPCC reports are policy relevant but policy neutral. This phrasing the direct policy in some way.

"... datasets with high TSI variability can explain more than 70 percent of the temperature variability since preindustrial times." Variability on what time scales? 11-year cycles? How would this impact the well documented surface temperature increase of > 1 deg C global temperature change since 1900? This and other chapters cite the IPCC AR6. The IPCC provides a global process to address the issues and uncertainties in this report. Perhaps this report can set a path for pivoting research and re-engaging in a useful manner.

A possible reason the method used by IPCC (optimal fingerprinting) can be considered to address McKittrick criticism is in [34]

- “For comparison, the IPCC estimate of forcing from the increase in atmospheric CO₂ compared to preindustrial times is approximately 2.2 W/m² (Figure 3.1.2).”
- This is comparing different time periods. Figure 8.2 includes data for 2023 and 2024, IPCC wouldn’t include the most recent years since it was published in 2021. Suggest updating from the IPCC data to Forster et al. (2024) that updates these indicators.
- Forster et al. (2024) “Indicators of Global Climate Change 2024: annual update of key indicators of the state of the climate system and human influence” <https://doi.org/10.5194/essd-17-2641-2025>
- “It is not easy to justify a new positive low cloud feedback that began emerging in 2015.”
- Explain why this is not easy?

The role of clouds remains the single most uncertainty in climate modeling, I am glad this was acknowledged here.

- “Other columns of IPCC Table 12.12 report on whether anthropogenic signals are expected to emerge this century under RCP8.5, the most extreme forcing scenario. We have omitted these columns for several reasons.”
- Can this be done for RCPs other than 8.5? The point of these omitted columns in IPCC Table 12.12 was to show that for some extremes we may not see an effect now, but will in the future. As it is presented here it implies future changes won’t matter.

The substance of this paragraph was addressed in my comments in earlier sections. Some of the claims made here are either exaggerations of the usual practice (RCP8.5 being referred to as BAU) or the comment about not fit for purpose (a comment extracted from a publication that is calling out for cloud resolved models.

What are the numbers in the 3rd column? Please include in the caption.

The discussion neglects the influence of vertical mixing on surface temperature. The global ocean has warmed, as documented by increases in global ocean heat content, e.g. Murphy, et al (2009)
doi:10.1029/2009JD012105

This section is hardly a balanced and objective review. The report authors are known opponents of optimal fingerprint approaches and have been a loud, but very small minority voice. It is interesting that even they conclude "...these analyses do not falsify results of the IPCC optimal fingerprinting method..."

This discussion is incomplete and fails to take into account the extensive literature on cloud and surface albedo feedbacks.. There is an extensive discussion about the post 2015 period and recent record annual global surface temperatures, implying, but not stating, that recent changes are the result of natural variability, or decreases in atmospheric aerosol concentrations. Comparing the radiative forcing from these changes with long-term GHG forcing, particularly CO₂ is misleading to the average reader. If the reduction in albedo can be tied to atmospheric cloud and aerosol changes, as the authors state, they cannot ignore the likelihood that the changes are the result of anthropogenic factors, i.e. sulfate aerosol reduction through cleaner technology and the well-documented positive low-cloud feedback.

Further research is needed on climate variability, effects of ocean circulation, solar variability, model simulation of variability and statistical methods.

Anthropogenic aerosol effects may also influence the inflections in temperature in ways that are challenging to account for

Reductions in aerosol indirect (cloud brightening) effects may also explain the reduced planetary albedo, not just from ships but from reduced fossil fuel aerosols coming from the continents, China and east asia especially. This paper describes the reduction in aerosols <https://www.nature.com/articles/s41612-023-00488-y> **This is another area that needs further research!**

I don't see an inconsistency between what Ch 11 and Ch 8 say. The first is about precipitation, droughts, tropical cyclones, heat - overall. The 2nd (ch 8) is about specific types of events in specific regions. Suggest to soften the statement about the contrasting assertions.

The science of attribution of extreme events is challenging and computationally expensive; this is an area where further research is needed with potential to use AI to improve statistics

- IPCC AR6 WGII has an assessment of agriculture (both in the SPM and in Chapter 5), and the IPCC also has a special report on Climate Change and Land, that are more comprehensive, recent, and cover a larger range of the literature than discussed here. Please update this chapter to reflect these findings.

This chapter is silent on agricultural effects on atmosphere and climate. Sokal and Kachel 2025 (10.3390/en18092272) conclude: "The current study demonstrated that agricultural activities significantly contribute to the emissions of three main greenhouse gases: carbon dioxide, nitrous oxide, and methane."

The last sentence in this summary is not necessarily accurate. The authors cite Ainsworth and Long (2020) with respect to increased CO2 effects on crops. But, Ainsworth and Long (2020) indicate that increased temp and decreased precipitation will have negative impacts on crops, while increased CO2 may offset some of those negative effects. It is the impact of CO2 on climate and the subsequent impact on crops that is the concern; not the direct effect of CO2 on crops.

The section title is "Field studies..." but the next three paragraphs are primarily about responses of potted plants in growth chambers or greenhouses (with exception of Blandino, which was a field study and showed a much smaller response than the pot studies). There is an important reason that the field data cited in Blandino and in SoyFACE show a smaller response than data from the potted plants under artificial conditions. These data are barely relevant to the topic you are trying to address.

CO2Science.org website is a highly biased and unreliable website. They report on papers, or specific sentences in papers, that support their view and ignore conflicting reports or qualifying sentences. This is a report for the US. Why are you citing soybean studies in China when there is a major soyFACE study in Illinois? Yield in SoyFACE increased in eCO2 but declined with increasing temperature (10.1093/jxb/erac503). eCO2 did not result in water savings as hypothesized but rather increased soil moisture depletion. (10.1098/rstb.2024.0230)

The problem with both the FACE studies and the soybean, Maize and wheat studies cited here are that they don't consider the effect of increase in temperature, changes in precipitation, humidity and other environmental factors along with CO2 increases. These compound effects have been shown in wide number of studies [35] [36][37][38][39]. A meta analysis of the compound effects shows that increase in CO2, along with temperature, precipitation changes effects the quality of the cereal and yields[40]. Another recent study evaluated the impacts on C3 and C4 crops and conclude that CO2 enrichment for C3 crops could increase but will be compensated by temperature increase. CO2 fertilization effect was found to be small for C4 [41]. It is important therefore that we consider the compound effects on crop yields and not just focus on CO2 enrichment.

Ainsworth et al, 2025 (10.1093/jxb/erac503) state: "Early season stimulation in biomass at elevated [CO2] more than offset lower stomatal conductance in crops, resulting in greater depletion of soil moisture, instead of water savings [30,31]. Wheat tiller survival and grain filling was also reduced by elevated [CO2] under dry conditions [32]. The CO2 fertilization effect was also significantly reduced by growth at elevated temperatures in FACE experiments, again suggesting that high atmospheric [CO2] may not offset the negative impacts of global warming [29]. Thus, it may be that, to date, elevated [CO2] and the resultant climate changes have had net positive effects on agroecosystem productivity, but as climate change intensifies, there will be a shift towards net negative effects as has been suggested for natural ecosystems [3]."

Taylor and Schlenker 2021 also note that 'Climate change will likely have negative impact on agriculture in aggregate, especially in regions exposed to extreme heat. And the CO2-driven yield increases may be offset by effects on food nutrition and quality'. However, the results for US have been a net positive in the recent past but not true in the lower latitudes.

Another study that should be cited is Hultgren et al, 2025 (10.1038/s41586-025-09085-w). "For all crops except rice, we estimate that warming will likely reduce global yields by 2050 (probability of loss ranges from 0.701 (sorghum) to 0.946 (wheat)... " "We estimate that global production declines 5.5×10^{14} kcal annually per 1 °C global mean surface temperature (GMST) rise (120 kcal per person per day or 4.4% of recommended consumption per 1 °C; $P < 0.001$). We project that adaptation and income growth alleviate 23% of global losses in 2050 and 34% at the end of the century (6% and 12%, respectively; moderate-emissions scenario), but substantial residual losses remain for all staples except rice." Regions with moderate average temperatures tend to suffer the largest yield losses.

"Adjusting for CO₂ fertilization does not qualitatively alter the structure of our findings ... but it does reduce the central estimate for end-of-century yield losses by 5.0– 9.5 percentage points... and increases the likelihood of positive aggregate effects."

Reference to Ziska paper greatly mischaracterizes it. Ziska clearly indicates an effect of eCO₂ on nutritional quality, but he indicates there is uncertainty in some of the mechanisms behind that response, including the simple dilution hypothesis. He states up front: "However, it is the direct effect of rising [CO₂] on plant chemistry and stoichiometry that represents a continuing threat to the nutritional integrity of both human and animal systems."

Evidence for nutritional decline under eCO₂ is widespread: Toreti et al, 2020 10.1038/s43016-020-00195-4) "A meta-analysis of 228 pairs of experimental observations on barley, potato, rice and wheat reported reductions in protein concentrations ranging on average from –15.3% to –9.8% under eCO₂, while the reduction was relatively small (–1.4%) in soybean." AND Ainsworth et al 2025: "The positive 'CO₂ fertilization' effect on crop yield is complicated by a decrease in many nutrients in elevated [CO₂]. An analysis of 130 varieties of plants found that elevated [CO₂] decreased the concentration of 25 important minerals by 8% on average, and the carbohydrate to mineral ratio was increased in these plants."

URL for Taylor and Schlenker paper is incorrect. It should be <https://www.nber.org/papers/w29320>. "papers" not "paper. Was this report peer reviewed? I cannot find it in a peer-reviewed journal.

This section seems to gloss over many key points. Certainly there are limits where extreme heat and drought (water is key) could be detrimental to crops. It also depends on which crops. There is extensive literature on how certain crop viability is shifting. So a region with particular crop experience and cultivation may no longer be able to sustain that agriculture. There is cost to shifting crops across regions and farmers that needs to be accounted for. And water is key - is there enough water in various regions for energy (power plants need water) as well as agriculture when conditions are very hot for sustained periods?

- Section 10.1 rightly points out that technological advances have substantially reduced losses from extreme weather events. We are better at forecasting, preparing for, and adapting to extreme weather than we were in the past, and these advances all help reduce impacts. However, our ability to do this in the future depends upon preparing for future changes – we can't simply assume adaptation will continue, we need to provide information on future risks and how those risks are impacted by climate change in order to inform continued technological advances and future adaptation.

- "In May 2025 NOAA announced it has withdrawn the Billion Dollar Disaster product from publication (Pielke Jr., 2025)."

- It would be better to cite an official government source from NOAA announcing this withdraw instead of a blog post commenting on it.

- Losses from extreme weather and climate events are driven by climate risk (the physical events themselves), exposure to those risks (population and infrastructure impacted by these events), and adaptation to those events (technological advances in forecasting, warning systems, and resiliency). Understanding these risks requires disentangling these drivers in order to prepare for the future.

The criticism of NOAA Billion dollar map is likely warranted as it was never completely transparent how the data was generated. However, changing the baseline assumption and refactoring the data using a different set of criteria to make the effects smaller also doesn't make sense. We should continue collecting this data and make the data accessible for researchers and peer review. Figure 10.1 doesn't address these issues, it is another methodology with unknown province of the data.

- There are additional citations that focus on the net effects of heat and cold related mortality that would be worth discussing and citing in this section.

- EPA. 2024. Draft Technical Documentation for the Framework for Evaluating Damages and Impacts (FrEDI). U.S. Environmental Protection Agency, EPA 430-R-24-001.

- Cromar et al., 2022. Global Health Impacts for Economic Models of Climate Change: A Systematic Review and Meta-Analysis. *Ann. Am. Thorac. Soc.* 19, 1203–1212. <https://doi.org/10.1513/AnnalsATS.202110-1193OC>

- Hsiang et al., 2017. Estimating economic damage from climate change in the United States. *Science* 356, 1362–1369.

- Mills et al., 2015. Climate change impacts on extreme temperature mortality in select metropolitan areas in the United States. *Clim. Change* 131, 83–95.

- Section 10.3.2 has good points about the importance of energy affordability and how energy poverty impacts adaptation behaviors. Cong et al. (2022) is another good citation for this section.

- Cong, S., Nock, D., Qiu, Y.L. et al. Unveiling hidden energy poverty using the energy equity gap. *Nat Commun* 13, 2456 (2022). <https://doi.org/10.1038/s41467-022-30146-5>

Severe weather continues to disproportionately impact lower income communities as well as business that requires outdoor exposure. A thorough analysis would need to include these points and I recommend adding this caveat to the section.

An additional concern is if the grid fails under either extreme heat demands or cold/severe-storm events, the impact on communities will not be uniform and depends on grid infrastructure which can be better in more affluent communities

Related to my previous comment, deaths due to extreme heat conditions appear to be increasing, particularly for certain populations and for those with underlying health conditions (which can be hard to detangle when attributing mortality). For example

<https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2792389>

- “Economists have long considered climate a relatively unimportant factor in economic growth, a view echoed by the IPCC itself in AR5.”

- It would be good to update this to the relevant conclusions in AR6 to reflect the more recent literature.

- This section makes an important point, much of the public does not understand that climate impacts are relative to projected future without climate change and need to be put in the context of economic growth and a future where our decedents are better off than we are today. O'Neill (2023) is a useful citation here, finding that, "Climate change research and assessments, including the most recent IPCC report, paint an increasingly dire picture of the future. However, the assumption that the future will be worse than the present may be wrong for many aspects of human well-being."

- O'Neill, B.C. Envisioning a future with climate change. *Nat. Clim. Chang.* 13, 874–876 (2023).

<https://doi.org/10.1038/s41558-023-01784-4>

Climate change can be expected to increase cold weather extremes in regions that have not been exposed to these effect

The same Nobel prize winning model in its most recent study concludes that [42] "We conclude with three results relevant for climate policy. First, both current policies (base run) and the extended Paris Accord fall short of limiting global warming to 2 °C or to the cost–benefit optimal level. Second, the economic stakes in global climate policy are substantial, with estimated net present value of economic benefits around \$120 trillion from the cost–benefit optimal policy. Third, once differences in discounting are considered, the baseline DICE-2023 estimate of the social cost of carbon (\$66/tCO₂ for 2020) aligns closely with other recent estimates". Obviously this is the latest and the current version of the model and these results are quite different from thus discussed on this page. These conclusions should also be included in this discussion to avoid selective reading of the literature.

- "Overall they could not detect a temperature effect on GDP or GDP growth, and they estimated the 95 percent confidence interval for the impact on global growth as of 2100 even under the exaggerated RCP8.5 warming scenario spans .86 percent to +388 percent. In other words the net effect is likely positive but too uncertain to distinguish from zero."

- This finding from Newell et al. (2021) needs to be contextualized with their complementary finding that models excluding growth effects and relating temperatures to GDP levels are far more certain. This is important to include since the actual USG application of SC-GHG does not include growth effects.

- Newell et al. (20201) states, "Models relating temperature to GDP levels yield climate impact estimates that are far more certain. The best such models imply GDP losses by 2100 of 1–3%, consistent with damage functions currently embedded in the major integrated assessment models that underpin the U.S. social cost of carbon."

- "To summarize, economists consider climate a relatively unimportant factor in economic growth, a view echoed by the IPCC itself in the Fifth Assessment Report."

- Need to update to reflect AR6 findings.

Similar empirical studies [43] point to "Using an empirical approach that provides a robust lower bound on the persistence of impacts on economic growth, we find that the world economy is committed to an income reduction of 19% within the next 26 years independent of future emission choices (relative to a baseline without climate impacts, likely range of 11–29% accounting for physical climate and empirical uncertainty). These damages already outweigh the mitigation costs required to limit global warming to 2 °C by sixfold over this near-term time frame and thereafter diverge strongly dependent on emission choices. Committed damages arise predominantly through changes in average temperature, but accounting for further climatic components raises estimates by approximately 50% and leads to stronger regional heterogeneity. Committed losses are projected for all regions except those at very high latitudes, at which reductions in temperature variability bring benefits. The largest losses are committed at lower latitudes in regions with lower cumulative historical emissions and lower present-day income". This study uses more of the daily temperature precipitation from 1600 hundred locations world wide instead of global means. There are numerous studies that show the cost, many described in the COP 21 and follow-on meetings

- "The concepts of estimation and uncertainty do not readily apply to SCC calculations."
- The discussion of uncertainty in IAMs should reflect EPA (2023) which incorporated many major advances in the treatment of uncertainty in integrated assessment modeling. See section A.8 of EPA (2023) for a summary of the quantified sources of uncertainty in the SCGHG estimates presented in EPA (2023)
- EPA (2023) "Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, 'Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review' – EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances." Docket ID No. EPA-HQ-OAR-2021-0317. https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf
- "Economists use IAMs to compute the SCC. Two of the best known are the Climate Framework for Uncertainty, Negotiation and Distribution ("FUND", Tol 1997) and Nordhaus' DICE."
- This is old literature. Given the advances since FUND and DICE were the leading SCC models, and this report's focus on literature post-2020, this section should be updated to focus on the models used to calculate the SC-GHG in EPA (2023) which represents advances over the approach used in FUND and DICE in response to NAS (2017) recommendations.
- EPA (2023) "Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, 'Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review' – EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances." Docket ID No. EPA-HQ-OAR-2021-0317. https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf

- “and a ‘judgmental adjustment’ of 0.5 percent for excluded impacts at 3°C warming”
 - EPA (2023) is the most up to date representation of the USG approach to SC-GHG incorporating the recommendations of the NAS (2017), and EPA (2023) explicitly does not included an adder for missing damages. This is another example of this chapter focusing on older less relevant literature, and it really should be comprehensively updated to address EPA (2023).
 - EPA (2023), page 59 states, “the model runs performed for this report do not adopt a 25% adder (as used in the DICE model (e.g., Nordhaus 2017)) to account for unknown or missing damages for the meta-analysis-based damage module.”
 - EPA (2023) “Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, ‘Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review’ – EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances.” Docket ID No. EPA-HQ-OAR-2021-0317.
 - NAS (2017) “Valuing Climate Damages. Updating Estimation of the Social Cost of Carbon Dioxide” <https://nap.nationalacademies.org/catalog/24651/valuing-climate-damages-updating-estimation-of-the-social-cost-of>
-
- “Abatement costs: IAMs represent the cost to the economy of reducing CO2 emissions. If CO2 emission reductions are assumed to be inexpensive, then the model will conclude that the optimal policy should aim for deeper emission cuts and vice versa.”
 - IAMs such as DICE are often used in a mode of calculating the optimal level of abatement where they do indeed compare the SCC with the marginal cost of abatement to find the optimum. However, these abatement costs are not part of the SCC calculation, and as used by the USG for the purpose of regulatory impact analyses, this feature of these IAMs is not used. Instead the USG has other more appropriate tools for analyzing the costs of specific regulations. This distinction is important here as the assumptions about the cost of emissions abatement does not have any bearing on the calculation of the SCC.

- “For example, in 2023 the U.S. Environmental Protection Agency raised its preferred SCC value about 5-fold over the estimates it had issued ten years earlier. This is not because new data had been collected or better mathematical methods had been invented, but because new assumptions had been used, and the validity of those assumptions was a separate question.”
- This paragraph contains the only reference to EPA (2023). First, the characterization of this update as only because new assumptions are used is factually incorrect. EPA (2023) represents a significant change to how the USG calculated the SC-GHG moving away from the FUND/DICE/PAGE models used in the IWG era (see EPA (2016)) to a modular approach as recommended by NAS (2017). This chapter needs to address the NAS (2017) recommendations and the EPA (2023) response to those recommendations to capture how SC-GHG was actually used by the USG and how the literature has evolved over time.
- EPA (2016) “Technical Support Document: - Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis - Under Executive Order 12866 - Interagency Working Group on Social Cost of Greenhouse Gases, United States Government.” https://www.epa.gov/sites/default/files/2016-12/documents/sc_co2_tsd_august_2016.pdf
- EPA (2023) “Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, ‘Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review’ – EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances.” Docket ID No. EPA-HQ-OAR-2021-0317. https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf
- NAS (2017) “Valuing Climate Damages. Updating Estimation of the Social Cost of Carbon Dioxide” <https://nap.nationalacademies.org/catalog/24651/valuing-climate-damages-updating-estimation-of-the-social-cost-of>
- “It should also be noted that the SCC is focused on the social costs of CO2 emissions from fossil fuel use. It does not measure the private marginal benefits to consumers and society from the availability of fossil fuels. Public willingness to pay for fuels of all types indicates the value to society of reliable, abundant fossil energy.”
- Two points to make about this section. First, the benefits to society generally accrue from the provision of energy services, not from specific fuels.” Second, the private marginal benefit of the availability of fossil fuels or energy services is not part of the SCC because changes in the cost and provision of the energy services should be part of the other side of the cost benefit analysis. The SCC is used to characterize the benefits of GHG emissions abatement, and the changes in the costs and availability of energy services are part of the cost of emissions abatement.
- “the uncertainty of SCC estimates doesn’t mean that other regulatory measures are inherently better or more efficient.”
- ‘regulatory measures’ is a poor word choice here as it can be confused with the regulation itself instead of the metric used to assess the regulation.

- This chapter has a heavy focus on the relation between climate change and economic growth and how this relationship factors into the SCC. There is an ongoing debate in the literature about persistence in climate damages as measured by changes in economic growth over time; however, these persistence effects on economic growth are not considered in the most recent version of the SC-GHG used by the USG – EPA (2023). This chapter generally ignores the existence of EPA (2023), it is mentioned once in passing, and does not inform readers that these controversial areas of current research are not included in government estimates of the SC-GHG.

- EPA (2023) page 58 says, “Given the statistical insignificance of the estimated coefficient on the productivity indicator in the published Howard and Sterner meta-analysis, the SC-GHG estimates presented in this report do not rely on Howard and Sterner’s specifications that include productivity effects. This is consistent with the authors’ recommendations in the published paper, to only consider the inclusion of the productivity impact in sensitivity analysis. However, the question of whether the effects of climate change impacts (e.g., temperature, tropical cyclones, and other extreme weather events) on the economy are only temporary or persistent is an active area of research. Over the past decade, a host of empirical studies have found evidence of temperature changes having persistent effects on the economy (e.g., Dell et al. 2012; Burke et al. 2015; Moore and Diaz 2015; Deryugina and Hsiang 2017; Ricke et al. 2018; Burke and Tanutama 2019; Colacito et al. 2019; Henseler and Schumacher 2019; Kahn et al. 2021; Kumar and Khanna 2019; Bastien-Olvera et al. 2022); this is an important finding because even small changes in economic growth rates accumulate into large economic impacts over time.”

- Section 11.2.4 has an interesting discussion of tipping points. This is an active area of research in the field, however it needs to be noted that tipping points or tipping elements are not included in the SC-GHG as used by the USG, see EPA (2023).

- EPA (2023), page 3 says, “For example, the modeling in this report omits most of the consequences of changes in precipitation, damages from extreme weather events, the potential for nongradual damages from passing critical thresholds (e.g., tipping elements) in natural or socioeconomic systems, and nonclimate mediated effects of GHG emissions other than CO2 fertilization (e.g., ocean acidification due to CO2 emissions, tropospheric ozone formation due to CH4 emissions). Importantly, this update does not yet reflect interaction effects and feedback effects within, and across, natural and human systems. For example, it does not explicitly reflect potential interactions among damage categories, such as those stemming from the interdependencies of energy, water, and land use. These interactions and feedbacks, and others, were highlighted by the National Academies as an important area of future research for longer-term enhancements in the SC-GHG estimation framework.”

USG has actually implemented SCC. This chapter would benefit from a more explicit focus on EPA (2023) as it represents the latest state of the literature on SCC, shows how the USG applies the SCC for regulatory impact analyses, shows how the USG updated the older IAG era SCC as represented in EPA (2016) in response to the recommendations in NAS (2017), shows how uncertainty is incorporated into SCC estimates, and discusses areas of the literature that are on the frontier, highly uncertain, and not ready to be incorporated into the official USG estimates.

- As currently written much of this chapter is focused on these highly uncertain areas of the literature that are not incorporated into USG SCC estimates (e.g. tipping points, and climate change impacts on economic growth). EPA (2023) discusses these issues and why they are not ready to be incorporated into the official estimates.
 - Furthermore, EPA (2023) estimates of the SC-GHG does incorporate climate benefits such as CO₂ fertilization and reductions in cold mortality discussed elsewhere in this report, and this should be noted and discussed.
 - Finally, this section ignores areas of real controversy in the calculation of SC-GHG as it is used by the USG – namely the question of global vs. domestic benefits and the treatment of discounting when valuing benefits to future generations. These controversial issues deserve far more attention in this report instead of spending so much space on issues on the fringe of the literature that are not even included in USG SC-GHG estimates.
 - EPA (2016) “Technical Support Document: - Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis - Under Executive Order 12866 - Interagency Working Group on Social Cost of Greenhouse Gases, United States Government.” https://www.epa.gov/sites/default/files/2016-12/documents/sc_co2_tsd_august_2016.pdf
 - EPA (2023) “Supplementary Material for the Regulatory Impact Analysis for the Final Rulemaking, ‘Standards of Performance for New, Reconstructed, and Modified Sources and Emissions Guidelines for Existing Sources: Oil and Natural Gas Sector Climate Review’ – EPA Report on the Social Cost of Greenhouse Gases: Estimates Incorporating Recent Scientific Advances.” Docket ID No. EPA-HQ-OAR-2021-0317. https://www.epa.gov/system/files/documents/2023-12/epa_scghg_2023_report_final.pdf
- The impacts of sea level rise and coastal inundation on coastal communities and in low-lying countries should be addressed. This will have impacts particularly on certain countries/regions but also on world economics

Research is needed to better understand the likelihood and impacts of various tipping points, including changes to ocean circulation patterns, ice sheet acceleration, sea-ice loss, die-off of tropical forests, thawing of permafrost, etc.

- This chapter summary is misleading as it is describing change from today’s temperature rather than the effect on future temperatures.
- “And even if global emissions were to stop tomorrow, it would take decades or centuries to see a meaningful reduction in the global CO₂ concentration and hence human influences on the climate.”
- If global emissions were to go to zero tomorrow, CO₂ concentrations would stabilize, and global temperature would be approximately constant. Net negative CO₂ would result in reductions of concentrations and temperatures. In other words, while reducing global CO₂ emissions to zero might not lead to near-term reductions in CO₂ concentrations from today’s levels, it would stop future increases in CO₂ concentrations.

- This section is focused on unilateral action, but needs to address questions related to collective action and individual actions by multiple parties when the benefits of abatement and the damages from emissions have extensive externalities.
 - Kotchen (2021) provides a theoretical perspective on the global vs. domestic SCC question and addresses questions related to how an individual country's decisions are impacted by the expected choices of other countries given that GHG emissions are globally mixed the benefits of GHG abatement are shared widely.
 - Kotchen (2021). Which Social Cost of Carbon? A Theoretical Perspective. Journal of the Association of Environmental and Resource Economists. <https://doi.org/10.1086/697241>
- “Since that rate has been averaging about 50 percent of emissions in recent decades, a reduction of global emissions by 50 percent would (at least temporarily) halt the rise in atmospheric CO₂.”
- This may be addressed in earlier chapters that this reviewer has not read yet, but the standard framing is that stabilizing CO₂ concentrations requires balancing emissions by sources and removals by sinks. Removals by sinks includes both natural sequestration and anthropogenic sequestration (e.g. afforestation if it can be done that increases the overall land sink removal, or direct air capture). It could be useful to clarify this here.
- “Lomborg (2016) estimated that full compliance with the Paris Accord would not stop warming, it would only prevent about 0.1C warming and delay hitting the baseline year 2100 temperature levels by about a decade.”
- Note that the Lomborg (2016) finding is related to, “the of impact of the total, global emission cuts implied by all the submitted INDCs,” this is different than ‘full compliance with the Paris Accord’ as the INDCs analyzed in the paper are only the first round of Intended Nationally Determined Contributions submitted ahead of COP-21, and the Paris Accord requires additional Nationally Determined Contributions every 5 years. Since Lomborg (2016) only analyzes these INDCs, and parties have submitted two rounds of NDCs since then, suggest changing the wording here to, “... full compliance with initial commitments submitted ahead of the Paris Accord.”

comment, then again broken into multiple parts for ease of reading. This is PART 1]

- Section 12.1 is titled, “The scale of the problem,” is framed around contrasting criteria air contaminants with CO2 emissions; however, there really are three separate issues that need to be disentangled: Stock vs. Flow.

- The challenge here is that abatement costs borne by people today benefit people in the future – temporal externalities.

- CO2 emissions are a stock pollutant, so impacts depend on the cumulative emissions to date, whereas traditional criteria are a flow pollutant and impacts depend on the emissions at that point in time. Emissions of a stock pollutant going to zero will not affect impacts at that point in time, as the stock remains unchanged, but will have large impacts in the future as the stock will cease to increase until emissions resume; whereas emissions of a flow pollutant going to zero will immediately eliminate impacts, but impacts will have no lasting effects on future impacts as they will immediately resume when emissions resume.

- The sentence at the end of the second paragraph of this section reads, “And even if global emissions were to stop tomorrow, it would take decades or centuries to see a meaningful reduction in the global CO2 concentration and hence human influences on the climate.” This is conflating the expectations of how stock and flow pollutants work – if global emissions were to stop tomorrow, the fact that there would be no meaningful reduction in global CO2 concentrations is simply the nature of how stock pollutants work, but there would be very large impacts on future temperatures since the stock would cease to increase.

- The stock nature of CO2 emissions is a real challenge for justifying CO2 emissions reductions, but it is centered on justifying the costs of abatement being born by people today in order to create a stream of benefits that grow over time and primarily accrue to future generations. This leads to questions important questions about how decision-makers should think about inter-generational discounting, and what costs should or should not be borne today for the benefit of future generations that will potentially be substantially better off.

- It’s worth noting that since CH4 emissions are relatively short lived, they behave much more like a flow pollutant than a stock pollutant. If CH4 emissions went to zero, concentrations would fall relatively quickly,

- [NOTE: this comment is long and could be difficult to read in excel. It is included here once as a full comment, then again broken into multiple parts for ease of reading. This is PART 2]

Globally mixed.

- The challenge here is that abatement costs borne by people in one location benefit people in other locations – spatial externalities.

- The next issue is that GHG emissions are globally mixed, so that emissions in any one location have impacts all over the world. This is a distinct issue from the stock vs. flow issue, as the CH4 example above showed a flow pollutant can also be globally mixed. One can also consider a stock pollutant that is not globally mixed, such as groundwater pollution that depends on the stock of contaminants accumulating in the groundwater not just the flow in any one time period as one would see in surface water pollution.

- The globally mixed nature of GHG emissions is a real problem for justifying climate policy as the benefits of any one country’s actions primarily accrue to people in other countries. Conversely, a country benefits from the actions of every other country, not just their own actions. This is a problem of game theory and collective action, as unilateral action may not be justified even if collective action would be. It’s worth noting that this isn’t just a problem of scale – these same problems arise for globally mixed CH4 emissions even though as noted above they behave more as a flow pollutant, and would arise for any globally mixed pollutant regardless of the scale of the problem. There is an extensive literature on transboundary pollution that explores these issues.

comment, then again broken into multiple parts for ease of reading. This is PART 3]
Scale.

- The challenge here is that solving the problem requires enormous change and potentially enormous costs. However, individual actions need to be considered at the margin.

- As discussed in this section and the comments above, stabilizing CO2 concentrations and stopping the increase in global mean temperature requires balancing anthropogenic emissions and removals by sinks. The level global mean temperature stabilizes at would generally depend on the cumulative CO2 emissions up to the point CO2 emissions on net go to zero (the stock of CO2 emissions), and the level of short-lived non-CO2 emissions at that time (the flow of non-CO2 emissions). This would involve an enormous change to the global energy system. The costs of this change would be highly dependent on the timing of reaching net-zero CO2 emissions – in 10 years by 2035 would involve crippling costs to the global economy, in 100 years by 2125 would involve a far more gradual transition and likely impose minimal costs when discounted back to today.

- What is missing in this chapter is an acknowledgement that policy needs to be thought of at the margin. Do the marginal benefits of the policy equal or exceed the marginal costs? This is how economists think of these problems and the essence of how the government conducts cost-benefit analysis to inform decision-makers under E.O. 12866. The case study presented in section 12.2 poses the wrong two question. Showing that a policy's emissions reductions are small relative to global emissions, and the resulting impact on global temperatures is also small is tautological. No individual policy will solve climate change, but that does not mean that the individual policy is not justified. The relevant questions for decision makers are: (1) what is the cost of the marginal ton of emissions reductions by this policy? and (2) are those marginal costs smaller than the marginal benefit of a ton of emissions reductions? In practice, question (2) is very difficult to answer, and requires judgements about how to value benefits to Americans vs. benefits to the rest of the world, how to discount benefits that accrue to future generations, and how to consider the impacts U.S. actions have on the likelihood that other countries will take actions that benefit the U.S. These are difficult questions that need to be answered to justify climate policy, and this section would be improved by acknowledging that these are the questions that need to be addressed instead of the awkward argument

comment, then again broken into multiple parts for ease of reading. This is the full comment]


- Section 12.1 is titled, “The scale of the problem,” is framed around contrasting criteria air contaminants with CO2 emissions; however, there really are three separate issues that need to be disentangled:
 - Stock vs. Flow.
 - The challenge here is that abatement costs borne by people today benefit people in the future – temporal externalities.
 - CO2 emissions are a stock pollutant, so impacts depend on the cumulative emissions to date, whereas traditional criteria are a flow pollutant and impacts depend on the emissions at that point in time. Emissions of a stock pollutant going to zero will not affect impacts at that point in time, as the stock remains unchanged, but will have large impacts in the future as the stock will cease to increase until emissions resume; whereas emissions of a flow pollutant going to zero will immediately eliminate impacts, but impacts will have no lasting effects on future impacts as they will immediately resume when emissions resume.
 - The sentence at the end of the second paragraph of this section reads, “And even if global emissions were to stop tomorrow, it would take decades or centuries to see a meaningful reduction in the global CO2 concentration and hence human influences on the climate.” This is conflating the expectations of how stock and flow pollutants work – if global emissions were to stop tomorrow, the fact that there would be no meaningful reduction in global CO2 concentrations is simply the nature of how stock pollutants work, but there would be very large impacts on future temperatures since the stock would cease to increase.
 - The stock nature of CO2 emissions is a real challenge for justifying CO2 emissions reductions, but it is centered on justifying the costs of abatement being born by people today in order to create a stream of benefits that grow over time and primarily accrue to future generations. This leads to questions important questions about how decision-makers should think about inter-generational discounting, and what costs should or should not be borne today for the benefit of future generations that will potentially be substantially better off.
 - It’s worth noting that since CH4 emissions are relatively short lived, they behave much more like a flow pollutant than a stock pollutant. If CH4 emissions went to zero, concentrations would fall relatively quickly,

The following sentence is misleading: "And even if global emissions were to stop tomorrow, it would take decades or centuries to see a meaningful reduction in the global CO2 concentration and hence human influences on the climate." It is true that bringing global emissions immediately to zero would not reduce CO2 concentrations on short timescales. It is not true that this same scenario means no reduction in human influence on the climate. We must compare the zero emissions scenario to something, and the most meaningful comparison is the current trajectory of **increasing** emissions and the resulting **super-linear** increase in concentration. So by reducing emissions to zero we clearly avoid the expected increases in concentration and the expected increases in radiative forcing together with increased near surface air temperature and all the other expected consequences. So our change in action - going to zero emissions - does in fact have an immediate and growing human influence on the trajectory of the climate system. That is not an argument for reducing our emissions to zero tomorrow, it is just an argument for logic and clarity in how we talk about consequences of different actions.

Regarding the sentence "The practice of referring to unilateral U.S. reductions as "combatting climate change" or "taking action on climate" therefore reflects a profound misunderstanding of the scale of the issue." This makes the dubious assumption that when people refer to "combatting climate change" or "taking action on climate" they are ignorant of the scale of the problem. Instead, it is probably more justifiable to assume that it is well-understood by most people who advocate for these local and national changes that the major emitters must act together to make a big difference in global emissions. It is at least worth considering, in the context of this report, that a path toward action at the necessary scale will require local and national action from a broad consortium, and that individual advocacy is a useful and necessary starting point for shifting the inertia of global policies.

This chapter is highly distressing, but I suppose it encapsulates the whole thesis of the document. Apparently, there is no point in policies to reduce emissions (or make the necessary changes in energy policy to accomplish emission reductions) because it will make no difference to the atmospheric CO₂ concentration or climate. So why bother? Would you have said this in 1970 (CO₂= 326), 1990 (CO₂ = 354 ppm), or 2010 (CO₂=390)? Probably so. And in 30 years when CO₂ is 500 ppm, will it be fair to look back and evaluate your apparent call for inaction in the statement here "And even if global emissions were to stop tomorrow, it would take decades or centuries to see a meaningful reduction in the global CO₂ concentration and hence human influences on the climate." But I suppose this attitude is justified if there is no adverse consequence and the only important effect of rising CO₂ is to make little plants in greenhouse grow bigger (e.g., Fig. 2.3), and indirect effects on heat stress, drought, fires, sea level, etc. aren't worth discussing. While small in terms of global emissions (just from the US transport sector), over period this will be significant reduction. We need to remember that US is the second highest emitter of CO₂.

Do the authors think it worthwhile for private or public research, such as at DOE, to seek breakthrough technology that might effectively remove massive amounts of CO₂ from the atmosphere?



- [1] Rachel E. Mason et al. Evidence, causes, and consequences of declining nitrogen availability in terrestrial ecosystems. *Science* 376, eabh3767(2022).DOI:10.1126/science.abh3767
- [2] C.E. Moore, et al the effect of increasing temperature on crop photosynthesis: from enzymes to ecosystems. *Exp. Bot.*, 72 (8) (2021), pp. 2822-2844

[3]C.R. Schwalm, S. Glendon,& P.B. Duffy, RCP8.5 tracks cumulative CO2 emissions, Proc. Natl. Acad. Sci. U.S.A. 117 (33) 19656-19657, <https://doi.org/10.1073/pnas.2007117117> (2020).
[4] C.R. Schwalm, S. Glendon, & P.B. Duffy, Reply to Hausfather and Peters: RCP8.5 is neither problematic nor misleading, Proc. Natl. Acad. Sci. U.S.A. 117 (45) 27793-27794, <https://doi.org/10.1073/pnas.2018008117> (2020).

[5] Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., et al. (2020). An assessment of Earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics*, 58, e2019RG000678. <https://doi.org/10.1029/2019RG000678> [6]
Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., et al. (2020). Causes of higher climate sensitivity in CMIP6 models. *Geophysical Research Letters*, 47, e2019GL085782. <https://doi.org/10.1029/2019GL085782>

[7]Hausfather, Z., Drake, H. F., Abbott, T., & Schmidt, G. A. (2020). Evaluating the performance of past climate model projections. *Geophysical Research Letters*, 47, e2019GL085378. <https://doi.org/10.1029/2019GL085378>

[8] <https://www.cdc.gov/climate-health/php/effects/index.html>

[9] Xinyuan Wei et al 2025 Environ. Res. Lett. 20 064013

- [10]Liu, J., Wennberg, P.O. An emergent constraint on the thermal sensitivity of photosynthesis and greenness in the high latitude northern forests. *Sci Rep* 14, 6189 (2024). <https://doi.org/10.1038/s41598-024-56362-1>
- [11] Francesco S.R. Pausata, Marco Gaetani, Gabriele Missouri, Alexis Berg, Danielle Maia de Souza, Rowan F. Sage, Peter B. deMenocal, The Greening of the Sahara: Past Changes and Future Implications, *One Earth*, Volume 2, Issue 3, 2020, Pages 235-250, ISSN 2590-3322,<https://doi.org/10.1016/j.oneear.2020.03.002>.
- [12]Lesk, C., Anderson, W., Rigden, A. et al. Compound heat and moisture extreme impacts on global crop yields under climate change. *Nat Rev Earth Environ* 3, 872–889 (2022). <https://doi.org/10.1038/s43017-022-00368-8>

[13]Xin Chen, Tiexi Chen, Bin He, Shuci Liu, Shengjie Zhou, Tingting Shi, The global greening continues despite increased drought stress since 2000, Global Ecology and Conservation, Volume 49, 2024, e02791,ISSN 2351-9894,

[14] Ursula M. Ruiz-Vera, Matthew Siebers, Sharon B. Gray, David W. Drag, David M. Rosenthal, Bruce A. Kimball, Donald R. Ort, Carl J. Bernacchi, Global Warming Can Negate the Expected CO₂ Stimulation in Photosynthesis and Productivity for Soybean Grown in the Midwestern United States , *Plant Physiology*, Volume 162, Issue 1, May 2013, Pages 410–423, <https://doi.org/10.1104/pp.112.211938> [15] Lamba S, Hall M, Röntfors M, et al. Physiological acclimation dampens initial effects of elevated temperature and atmospheric CO₂ concentration in mature boreal Norway spruce. *Plant Cell Environ.* 2018; 41: 300–313. <https://doi.org/10.1111/pce.13079>

- [16]van Vuuren, D.P., Edmonds, J., Kainuma, M. et al. The representative concentration pathways: an overview. *Climatic Change* 109, 5 (2011). <https://doi.org/10.1007/s10584-011-0148-z>
- [17]Hausfather, Z. (2025). An assessment of current policy scenarios over the 21st century and the reduced plausibility of high-emissions pathways. *Dialogues on Climate Change*, 2(1), 26-32. <https://doi.org/10.1177/29768659241304854> (Original work published 2025)

[18] Parker, D. Large-scale warming is not urban. *Nature* 432, 290 (2004).

<https://doi.org/10.1038/432290a>

[19] Parker, D.E. (2010), Urban heat island effects on estimates of observed climate change. *WIREs Clim Change*, 1: 123-133. <https://doi.org/10.1002/wcc.21>.

[20] Parker, D. E., 2006: A Demonstration That Large-Scale Warming Is Not Urban. *J. Climate*, 19, 2882–2895, <https://doi.org/10.1175/JCLI3730.1>.

[21] Michael Mann, Our Fragile Moment: How lessons from Earth's past can help us survive climate crisis, Hach

[22]Hausfather, Z., Drake, H. F., Abbott, T., & Schmidt, G. A. (2020). Evaluating the performance of past climate model projections. *Geophysical Research Letters*, 47, e2019GL085378. <https://doi.org/10.1029/2019GL085378>

[23]Chung, ES., Kim, SJ., Sohn, BJ. et al. Multi-decadal climate variability and satellite biases have amplified model-observation discrepancies in tropical troposphere warming estimates. *Commun Earth Environ* 5, 342 (2024). <https://doi.org/10.1038/s43247-024-01510-8>

[24]Merlis, T. M., Guendelman, I., Cheng, K.-Y., Harris, L., Chen, Y.-T., Bretherton, C. S., et al. (2024). The vertical structure of tropical temperature change in global storm-resolving model simulations of climate change. *Geophysical Research Letters*, 51, e2024GL111549. <https://doi.org/10.1029/2024GL111549>

[25] B.D. Santer, S. Po-Chedley, L. Zhao, C. Zou, Q. Fu, S. Solomon, D.W.J. Thompson, C. Mears, & K.E. Taylor, Exceptional stratospheric contribution to human fingerprints on atmospheric temperature, *Proc. Natl. Acad. Sci. U.S.A.* 120 (20) e2300758120, <https://doi.org/10.1073/pnas.2300758120> (2023).

[26]Cohn, T. A., and H. F. Lins (2005), Nature's style: Naturally trendy, Geophysics. Res. Lett., 32, L23402, doi:10.1029/2005GL024476.

[27]<https://www.epa.gov/climate-indicators/climate-change-indicators-heat-waves>

[28]Ronald D. Leeper, Rocky Bilotta, Bryan Petersen, Crystal J. Stiles, Richard Heim, Brian Fuchs, Olivier P. Prat, Michael Palecki, Steve Ansari

[29]Tyukavina Alexandra , Potapov Peter , Hansen Matthew C. , Pickens Amy H. , Stehman Stephen V. , Turubanova Svetlana , Parker Diana , Zalles Viviana , Lima André , Kommareddy Indrani , Song Xiao-Peng , Wang Lei , Harris Nancy; Global Trends of Forest Loss Due to Fire From 2001 to 2019, *Frontiers in Remote Sensing*.Volume 3 – 2022,2022.

[30]Cunningham, C.X., Williamson, G.J. & Bowman, D.M.J.S. Increasing frequency and intensity of the most extreme wildfires on Earth. *Nat Ecol Evol* 8, 1420–1425 (2024).

<https://doi.org/10.1038/s41559-024-02452-2>

[31]Patrick H. Freeborn, W. Matt Jolly, Mark A. Cochrane, Gareth Roberts, Large wildfire driven increases in nighttime fire activity observed across CONUS from 2003–2020, *Remote Sensing of Environment*, Volume 268, 2022,12777

[32]Parks, S.A., Guiterman, C.H., Margolis, E.Q. et al. A fire deficit persists across diverse North American forests despite recent increases in area burned. *Nat Commun* 16, 1493 (2025).

<https://doi.org/10.1038/s41467-025-56333-8>

[33]Hamlington, B.D., Bellas-Manley, A., Willis, J.K. et al. The rate of global sea level rise doubled during the past three decades. *Commun Earth Environ* 5, 601 (2024). <https://doi.org/10.1038/s43247-024-01761-5>

[34]Chen, H., Chen, S.X. & Mu, M. A statistical review on the optimal fingerprinting approach in climate change studies. *Clim Dyn* 62, 1439–1446 (2024). <https://doi.org/10.1007/s00382-023-06975-5>

[35] Ruiz-Vera UM, Siebers M, Gray SB, Drag DW, Rosenthal DM, Kimball BA, Ort DR, Bernacchi CJ. Global warming can negate the expected CO₂ stimulation in photosynthesis and productivity for soybean grown in the Midwestern United States. *Plant Physiol.* 2013 May;162(1):410-23

[36] Levine LH, Richards JT, Wheeler RM. Super-elevated CO₂ interferes with stomatal response to ABA and night closure in soybean (*Glycine max*). *J Plant Physiol.* 2009 Jun 1;166(9):903-13.

[37] Večeřová K, Oravec M, Puranik S, Findurová H, Veselá B, Opoku E, Ofori-Amanfo KK, Klem K, Urban O, Sahu PP. Single and interactive effects of variables associated with climate change on wheat metabolome. *Front Plant Sci.* 2022 Oct 10;13:1002561.

[38] Thomas, J.M.G., Boote, K.J., Allen, L.H., Gallo-Meagher, M. and Davis, J.M. (2003), Elevated Temperature and Carbon Dioxide Effects on Soybean Seed Composition and Transcript Abundance. *Crop Sci.*, 43: 1548-1557. <https://doi.org/10.2135/cropsci2003.1548>

[39] Lesk, C., Anderson, W., Rigden, A. et al. Compound heat and moisture extreme impacts on global crop yields under climate change. *Nat Rev Earth Environ* 3, 872–889 (2022). <https://doi.org/10.1038/s43017-022-00368-8>

[40] Ben Mariem S, Soba D, Zhou B, Loladze I, Morales F, Aranjuelo I. Climate Change, Crop Yields, and Grain Quality of C3 Cereals: A Meta-Analysis of [CO₂], Temperature, and Drought Effects. *Plants.* 2021; 10(6):1052. <https://doi.org/10.3390/plants10061052>

[41] Rezaei, E.E., Webber, H., Asseng, S. et al. Climate change impacts on crop yields. *Nat Rev Earth Environ* 4, 831–846 (2023). <https://doi.org/10.1038/s43017-023-00491-0>

[42]L. Barrage, & W. Nordhaus, Policies, projections, and the social cost of carbon: Results from the DICE-2023 model, Proc. Natl. Acad. Sci. U.S.A. 121 (13) e2312030121, <https://doi.org/10.1073/pnas.2312030121> (2024).

[43] Kotz, M., Levermann, A. & Wenz, L. The economic commitment of climate change. *Nature* 628, 551–557 (2024). <https://doi.org/10.1038/s41586-024-07219-0>



ette Group Inc., 2024.

From: Judith Curry <curry.judith@gmail.com>
Sent: Thursday, July 31, 2025 5:24 PM
To: Fisher, Travis
Cc: Roy Spencer; Park, Charles; Ross McKittrick; John Christy; Steven Koonin; Loucks, Joshua; Cohen, Seth
Subject: Re: [EXTERNAL] DOE reviews

Thanks. I think Ross' version in msword is easier to work with, including the color coded comments.

On Thu, Jul 31, 2025 at 2:18 PM Fisher, Travis <travis.fisher@hq.doe.gov> wrote:

I'm attaching the version of the internal review comments that was compiled by the Office of Science. Note that I haven't checked to see if they numbered the reviewers the same way we did.

Even though we haven't had our introductory call yet, I'm also including Charles Park. Let me know if you're available tomorrow afternoon for a call to introduce Charles.

And I just saw Steve's email—yes, we should compile the anonymous reviews and your responses, and we should strongly consider releasing that document (and I believe we would have to if anyone requests it).

From: Judith Curry <curry.judith@gmail.com>
Sent: Thursday, July 31, 2025 5:06 PM
To: Roy Spencer <roywspencer@hotmail.com>
Cc: Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>; Steven Koonin <steven.koonin@googlemail.com>; Josh Loucks <loucksj14@gmail.com>; Cohen, Seth <seth.cohen@hq.doe.gov>; Fisher, Travis <travis.fisher@hq.doe.gov>
Subject: [EXTERNAL] DOE reviews

Ross, if you can send me the latest version of the 8 DOE review responses, i can clean those up, verify changes were made, and send around for approval. Would be good to get those out of the way before the public comments

On Thu, Jul 31, 2025 at 12:27 PM Roy Spencer <roywspencer@hotmail.com> wrote:

Ironical that no one cared when MY work was mischaracterized by the 97-Percenters' as being part of the 3%, even though by their criteria I was part of the 97% consensus.

I complained, but all they said was, "well, we put you in the 3 percent category".

The bias is rampant.

-Roy

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 31, 2025 11:27 AM
To: Roy W. Spencer <roywspencer@hotmail.com>; Judith Curry <curry.judith@gmail.com>; John Christy <climateman60@gmail.com>; Steven Koonin <steven.koonin@googlemail.com>; Josh Loucks <loucksj14@gmail.com>; Cohen, Seth <seth.cohen@hq.doe.gov>; Fisher, Travis <travis.fisher@hq.doe.gov>
Subject: Fwd: Fw: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

John and I got this from Tim Vogelsang at MSU. Carbon Brief is contacting everyone whose work was cited trying to gather up claims of misrepresentation. As Tim notes, there is no option on the form for saying your work was correctly characterized.

From: Vogelsang, Timothy <tjv@msu.edu>
Sent: Thursday, July 31, 2025 12:19 PM
To: Ross McKittrick <rmckitri@uoguelph.ca>; John Christy <climateman60@gmail.com>
Subject: RE: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

CAUTION: This email originated from outside of the University of Guelph. Do not click links or open attachments unless you recognize the sender and know the content is safe. If in doubt, forward suspicious emails to IThelp@uoguelph.ca.

If you look at the pdf I was sent with the instructions, there is no option to report that someone's research has been correctly characterized. That's not a very scientific factchecking exercise, is it!

Tim

From: Ross McKittrick <rmckitri@uoguelph.ca>
Sent: Thursday, July 31, 2025 12:15 PM
To: Vogelsang, Timothy <tjv@msu.edu>; John Christy <climateman60@gmail.com>
Subject: Re: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

I've heard of Carbon Brief but only read snippets of their articles cited elsewhere. They seem to be heavily into promoting renewables.

It's interesting to watch the machine swing into gear. There is a public comment docket opening up and I expect there will be quite the avalanche of material submitted, which we will need to prepare responses to, all of which will form part of the public record. If the comments are substantive we'll readily make revisions. We responded to a lot of comments from the internal DoE staff during the pre-publication review stage. But some of the most hostile comments didn't provide any data or citations in support so we couldn't do anything with them.

Ross

Dr. Ross McKittrick

Professor of Economics

Department of Economics and Finance

rossmckitrick.com



University of Guelph | Guelph ON Canada N1G 3P9

From: Vogelsang, Timothy <tjv@msu.edu>
Sent: Thursday, July 31, 2025 12:02 PM
To: Ross McKittrick <rmckitri@uoguelph.ca>; John Christy <climateman60@gmail.com>
Subject: FW: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

CAUTION: This email originated from outside of the University of Guelph. Do not click links or open attachments unless you recognize the sender and know the content is safe. If in doubt, forward suspicious emails to IThep@uoguelph.ca.

Ross and John,

I just got this email. Maybe you know about this group. I'm sure this is just the tip of the iceberg of pushback coming.

Tim

From: Leo Hickman <leo.hickman@carbonbrief.org>

Sent: Thursday, July 31, 2025 11:49 AM

To: Leo Hickman <leo.hickman@carbonbrief.org>

Subject: Carbon Brief factcheck - your study has been cited in new US DoE 'critical review'

Hello,

I'm contacting you because we have noticed that at least one of your studies has been cited in a new report published this week by the US Dept of Energy titled, "A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate."

A number of scientists cited in the report have already publicly highlighted that their findings have been falsely or misleadingly characterised. It seems that the report is intended to be used as scientific justification by the Trump administration in its efforts to revoke the US Environmental Protection Agency's "endangerment finding" on greenhouse gases.

Carbon Brief is now seeking your help to assess whether your own study, or indeed your area(s) of expertise more widely, have been mischaracterised in the report. This is part of a wider 'crowdsourced' fact-checking effort we are now convening to draw on inputs from experts in their respective fields.

I have attached a PDF with some instructions on how you can participate with your own insights and reflections, should you wish. (The instructions include ways to do this anonymously and securely, if that is your preference.)

Additionally, if you know of a colleague or co-author who might wish to participate, please feel free to forward this email to them.

Our deadline for submissions is Monday 11 August 2025 - 0900 GMT.

Many thanks in advance.

Best wishes, Leo

--

Leo Hickman

Editor, Carbon Brief


Phone: + 44 (0) 7731 403979

Social: Twitter; BlueSky; LinkedIn

Web: www.carbonbrief.org

Need a digest of the latest climate change-related headlines sent to your inbox? You can sign up to Carbon Brief's free daily, weekly, Cropped and China Briefing newsletters here.

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

This message does not originate from a known Department of Energy email system.
Use caution if this message contains attachments, links or requests for information.

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

From: John Christy <climateman60@gmail.com>
Sent: Monday, June 30, 2025 5:24 PM
To: Cohen, Seth
Cc: Steven Koonin; Travis Fisher; Josh Loucks; Ross McKittrick; Judith Curry; Roy W. Spencer
Subject: Re: [EXTERNAL] NYTimes on warming

All

I was in Fairbanks Alaska when the Heat Advisory was announced. Didn't bother us (85F is a cool June day in Alabama - and their dewpoint was 24F - crisp, bone dry air).

Station, (Year Began), June 2025 Max, Hottest ever June temp, All Time Hottest
Aurora, (2002), 86F, 93F, 93F
College Observatory, (1948), 86F, 94F, 94F
Fairbanks Intl AP, (1929), 85F, 96F, 96F
Fairbanks RAWs, (1988), M, 95F, 100F
Univ Exp Stn, (1904), M (report after end of month), 95F, 95F

June usually has the hottest temperature since the interior (Fairbanks) is so dry, and solar radiation is the big deal there, so June usually generates the hottest annual temperature. This First Ever heat advisory warned of temperatures 11F to 7F below the max temperatures we know have visited the region.

From the AP story on Alaska Heat Advisories, "It's not the first instance of unusually high temperatures in what many consider the nation's coldest state, but the National Weather Service only recently allowed for heat advisories to be issued there."

I guess the NY Times doesn't appreciate historical perspective.

For the global tropospheric temperatures, the situation looks more like a step function - it's a little warmer after each El Nino than before. Short time series allow all kinds of mischief with trends. The UAH global atmospheric trend for 2010 to 2025 is +0.36 °C/decade. But starting in 1998, the trend is cut to half of that (+0.17 °C/decade). If you are going to end on a peak, you need to start on a peak - and 15 years is way too short for a climate signal. To chuckle at the absurdity, the global trend since April 2024 is -3.3 °C/decade ... we're going to freeze to death!

By the way, Vermont's flood-of-record occurred in 1927.

"Nothing is defying our big picture about the physics of the climate system," Dr. Bitz said. Uh, Dr. Bitz, ... how about models warming much too quickly?

That's enough of that.

John C.

On Mon, Jun 30, 2025 at 2:31 PM Cohen, Seth <seth.cohen@hq.doe.gov> wrote:

Y axis is overweighted up, x axis has a skewed tail to the left (90 years), versus 50 on the right.

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Monday, June 30, 2025 3:30 PM
To: 'Travis Fisher' <travis.scott.fisher@gmail.com>; 'Josh Loucks' <loucksj14@gmail.com>; Cohen, Seth <seth.cohen@hq.doe.gov>
Cc: 'Ross McKittrick' <ross.mckittrick@gmail.com>; 'Judith Curry' <curry.judith@gmail.com>; 'Roy W. Spencer' <roywspencer@hotmail.com>; 'John Christy' <climateman60@gmail.com>
Subject: [EXTERNAL] NYTimes on warming

A quiz for our non-scientist colleagues:

Critique the lead graphics in this recent NYT article:

https://www.nytimes.com/2025/06/26/climate/climate-heat-intensity.html?unlocked_article_code=1.S08.1xcl.nm658aV-Q3kU&smid=url-share

Steve Koonin

PS Bonus question- critique the article itself

This message does not originate from a known Department of Energy email system.
Use caution if this message contains attachments, links or requests for information.

From: Ison, Jeremy T. <jeremy.ison@hq.doe.gov>
Sent: Thursday, August 21, 2025 6:45 PM
To: Steven Koonin
Cc: 'Judith Curry'; 'Ross McKittrick'; 'Roy Spencer'; 'John Christy'; Barrios, Audrey
Subject: Re: [EXTERNAL] RE: catchup call?

Steve,

Thank you for reaching out and apologies on the delay, as I was out of the office today.

I was waiting for the completion of some internal discussions before setting up this meeting, but happy to get together and address what I can.

I can meet tomorrow from 9-10:30am and then again at 4pm ET.

I can generally make myself available all of Monday for this discussion and may have more to share at that time.

Thank you,
Jeremy

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Thursday, August 21, 2025 5:54 PM
To: Ison, Jeremy T. <jeremy.ison@hq.doe.gov>
Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; 'Roy Spencer' <roywspencer@hotmail.com>; 'John Christy' <climateman60@gmail.com>; Barrios, Audrey <audrey.barrios@hq.doe.gov>
Subject: [EXTERNAL] RE: catchup call?

Jeremy – following up on this as the CWG sees some urgency in having a discussion. After on last week's call, we thought you were going to arrange one this week.

Here are some of the issues we'd like to discuss:

- Release all the public comments submitted so far on the CWG report (we have spotted 200 that have been released in the Federal Register docket, but there are more than 200 still to go)
- Post (on the DOE website?) the internal DOE and lab peer review comments (and our responses) that occurred before the report was released. This needed to counter the allegation that the report wasn't peer reviewed before release.
- We would like an update on the status of external review committee for the CWG Report
- We need an update on the lawsuit filed against the DOE and CWG, and whether there are any constraints (or even just cautions) on our public statements
- We would like clarification on the schedule for revising the CWG Report and responding to comments

- We would like an update on the status of the NCA5 Review, and whether the charge for that has changed.
- Curry does not yet have a contract
- Christy and Spencer's SGE appointments have just ended

Steve

From: Steven Koonin <steven.koonin@gmail.com>

Sent: Thursday, August 21, 2025 11:34 AM

To: Ison, Jeremy T. <jeremy.ison@hq.doe.gov>

Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; 'Roy Spencer' <roywspencer@hotmail.com>; 'John Christy' <climateman60@gmail.com>; 'Barrios, Audrey' <audrey.barrios@hq.doe.gov>

Subject: catchup call?

Jeremy –

Could we do a call sometime soon (today, tomorrow, or early next week) to catch up on recent events and map out plans going forward?

Steve Koonin

This message does not originate from a known Department of Energy email system.
Use caution if this message contains attachments, links or requests for information.

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Monday, July 28, 2025 9:06 AM
To: Roy Spencer; Ross McKittrick; John Christy
Cc: Steven Koonin; Judith Curry; Loucks, Joshua; Cohen, Seth
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Hi Roy,

The report should be released tomorrow, likely in the afternoon. I'll make sure everyone here is in the loop when it goes out. I believe the Secretary is aware that he owes you all some glowing public comments and an open bar tab at some point.

On media issues, my preference is to let the professionals take the lead. I suggest you forward media requests to Andrea Woods in DOE's Office of Public Affairs so she's aware of the questions: andrea.woods@hq.doe.gov

That said, people will want to talk to the CWG authors. If you're willing to talk to reporters, just let Andrea know. She should have a good sense of which ones are coming to this issue in good faith and can set up interviews.

Best,
Travis

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Monday, July 28, 2025 6:01 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis:

Any word on when the Report will be released?
I'm leaning toward not responding to any media queries, unless others think it's needed at some point.

-Roy

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Friday, July 25, 2025 3:52 PM
To: Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>
Cc: Roy Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth

<seth.cohen@hq.doe.gov>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Hi all,

I have a request for a minor change in the preface. The last line presently reads: “We are also grateful to a team of anonymous peer reviewers whose input helped improve the final report.”

Would it cause any heartburn to change the line to read “We are also grateful to a team of anonymous **DOE and national lab** reviewers whose input helped improve the final report.”?

Best,

Travis

PS – the Secretary is reading the updated version tonight.

From: Ross McKittrick <ross.mckittrick@gmail.com>

Sent: Thursday, July 24, 2025 7:12 PM

To: John Christy <climateman60@gmail.com>

Cc: Roy Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Didn’t even think of that. If it comes up I guess I’d invoke McShane and Weiner and say the ipcc has never addressed the problem that the millennial scale back-projections at the hemispheric scale are simply too noisy to draw precise conclusions. And nothing core to our report depended one way or the other on those issues.

On Thu, Jul 24, 2025 at 5:59 PM John Christy <climateman60@gmail.com> wrote:

Ross

What will you say when asked why we didn't deal with all the Hockey Sticks from Mann to IPCC AR6?

John C.

Sent from my iPhone

On Jul 24, 2025, at 4:27 PM, Roy Spencer <roywspencer@hotmail.com> wrote:

Dang! Wish I had thought of getting a Canadian phone!

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 24, 2025 3:20 PM
To: Steven Koonin <steven.koonin@gmail.com>
Cc: John Christy <climateman60@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

They'll have trouble reaching me. My Canadian phone died and I won't be able to replace it and reactivate the number until I'm back home next week.

On Thu, Jul 24, 2025 at 3:14 PM Steven Koonin <steven.koonin@gmail.com> wrote:

Got a similar call from Scott Waldmann, but let it go to voicemail.

He said he wanted to speak on background about the endangerment finding and "some of the climate work" I've been doing.

I've not returned his call.

From: John Christy <climateman60@gmail.com>
Sent: Thursday, July 24, 2025 3:13 PM
To: Ross McKitrick <ross.mckitrick@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Just got a call from Scott at Politico stating he knows I've worked on a DoE science report. I said no comment. (He uses his wife's cell phone, so you don't know it's him when you pick up.)

John C.

On Thu, Jul 24, 2025 at 10:24 AM Ross McKitrick <ross.mckitrick@gmail.com> wrote:

PS here are the report files now with a hi-res logo included.

On Thu, Jul 24, 2025 at 10:18 AM Steven Koonin <steven.koonin@gmail.com> wrote:

If you want to shorten Ross' proposed title a bit, it could read:

Department of Energy Releases Report That Challenges Common Misconceptions About Climate Change

From: Ross McKitrick <ross.mckitrick@gmail.com>
Sent: Thursday, July 24, 2025 11:16 AM
To: John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

The title might need to say a bit more, e.g.

Department of Energy Releases Report of Climate Working Group That Challenges Common Misconceptions About Climate Change

and then the remainder follows the text as shown in Steve's edit, with "renowned" replaced by "eminent", there being no adjective that won't draw scorn from the haters, but we might as well pick the word we like best.

On Thu, Jul 24, 2025 at 9:09 AM John Christy <climateman60@gmail.com> wrote:

Looks good.

John C.

On Thu, Jul 24, 2025 at 8:59 AM Steven Koonin <steven.koonin@gmail.com> wrote:

And with me.

SEK

From: Roy Spencer <roywspencer@hotmail.com>

Sent: Thursday, July 24, 2025 9:49 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>; Steven Koonin <steven.koonin@gmail.com>; 'John Christy' <climateman60@gmail.com>

Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Fine with me.

Sent from my Verizon, Samsung Galaxy smartphone

----- Original message -----

From: "Fisher, Travis" <travis.fisher@hq.doe.gov>

Date: 7/24/25 8:42 AM (GMT-06:00)

To: Steven Koonin <steven.koonin@gmail.com>, 'John Christy' <climateman60@gmail.com>

Cc: 'Judith Curry' <curry.judith@gmail.com>, 'Ross McKittrick' <ross.mckittrick@gmail.com>, "Loucks, Joshua" <Joshua.Loucks@hq.doe.gov>, "Cohen, Seth" <seth.cohen@hq.doe.gov>, "Roy W. Spencer" <roywspencer@hotmail.com>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Ok I'm going with eminent for now. All of this is subject to review by the office of public affairs, so we'll see what sticks.

In this next iteration I added your bios at the end (straight copy/paste from the report). Are you all comfortable if we decide to take that approach?

From: Steven Koonin <steven.koonin@gmail.com>

Sent: Thursday, July 24, 2025 9:21 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>

Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'Roy W. Spencer' <roywspencer@hotmail.com>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Perhaps “eminent” .

Although in some eyes, “notorious” would be appropriate.

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Thursday, July 24, 2025 9:07 AM
To: John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

I don’t want anyone to think of you all as circus performers! I was looking for a synonym for “distinguished,” with the bonus of being well-known. I’m open to ideas: RENOWNED Synonyms: 104 Similar and Opposite Words | Merriam-Webster Thesaurus

From: John Christy <climateman60@gmail.com>
Sent: Thursday, July 24, 2025 8:53 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis

“Renowned” sounds a little like a circus performer.

I wonder how many reporters will actually read the whole document? I suspect one of our common answers to their questions is, “just read the report!”

John C.

Sent from my iPhone

On Jul 23, 2025, at 8:28 PM, Fisher, Travis <travis.fisher@hq.doe.gov> wrote:

It's a heck of a report. I can't wait to see where we go from here!

Here's my first attempt at a press release—please make any edits or suggestions you like.

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Wednesday, July 23, 2025 5:50 PM
To: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>; 'Roy W. Spencer' <roywspencer@hotmail.com>
Subject: [EXTERNAL] RE: July 23rd version - final (?)

Second that! Amazing effort, Ross.

SEK

From: Judith Curry <curry.judith@gmail.com>
Sent: Wednesday, July 23, 2025 5:49 PM
To: Ross McKittrick <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; John Christy <climateman60@gmail.com>; Roy W. Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>
Subject: Re: July 23rd version - final (?)

congrats to us all, and special thanks to RM for pulling it all together

On Wed, Jul 23, 2025 at 2:19 PM Ross McKittrick
<ross.mckittrick@gmail.com> wrote:


Attached in MSWord and PDF formats.

All of Steve's corrections incorporated, also John's stray L and everything else I'm aware of.

The only remaining correction I have not been able to make is using a high-res version of the DoE logo. If that becomes the main focus of public criticism I will say we did well.

Ross

--

 Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

This message does not originate from a known Department of Energy email system.

Use caution if this message contains attachments, links or requests for information.

<CWG Press Release.docx>

From: Fisher, Travis [travis.fisher@hq.doe.gov]
Sent: 7/28/2025 2:45:32 PM
To: Roy Spencer [roywspencer@hotmail.com]; Ross McKittrick [ross.mckittrick@gmail.com]; John Christy [climatemanager60@gmail.com]
CC: Steven Koonin [steven.koonin@gmail.com]; Judith Curry [curry.judith@gmail.com]; Loucks, Joshua [Joshua.Loucks@hq.doe.gov]; Cohen, Seth [seth.cohen@hq.doe.gov]
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)
Attachments: Draft Memo from CWG.docx

Hi all,

This is a matter of administrative housekeeping—I want to make sure you're all aware of the process and comfortable with it.

The attached memo presents an action item to the Secretary (to publish the report). It's a formality, but please let me know if you'd like to see any changes to the text.

Best,
Travis

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Monday, July 28, 2025 9:21 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climatemanager60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

In my experience, any of the right-leaning news outlets have always treated me well. But (unlike John), I'm no longer willing to talk to the "usual suspects"... CNN, MSNBC, NPR and its derivatives, and the British press

Years ago the science reporter at BBC, Richard Black, put some *ad hominem* in his report on my interview, and he took a lot of flak for it from readers... he tried to get me to do a follow-up to save face, and I said sorry, buddy, you had your chance.

-Roy

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Monday, July 28, 2025 8:05 AM
To: Roy Spencer <roywspencer@hotmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climatemanager60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Hi Roy,

The report should be released tomorrow, likely in the afternoon. I'll make sure everyone here is in the loop when it goes out. I believe the Secretary is aware that he owes you all some glowing public comments and an open bar tab at some point.

On media issues, my preference is to let the professionals take the lead. I suggest you forward media requests to Andrea Woods in DOE's Office of Public Affairs so she's aware of the questions: andrea.woods@hq.doe.gov

That said, people will want to talk to the CWG authors. If you're willing to talk to reporters, just let Andrea know. She should have a good sense of which ones are coming to this issue in good faith and can set up interviews.

Best,
Travis

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Monday, July 28, 2025 6:01 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis:

Any word on when the Report will be released?

I'm leaning toward not responding to any media queries, unless others think it's needed at some point.

-Roy

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Friday, July 25, 2025 3:52 PM
To: Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>
Cc: Roy Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Hi all,

I have a request for a minor change in the preface. The last line presently reads: “We are also grateful to a team of anonymous peer reviewers whose input helped improve the final report.”

Would it cause any heartburn to change the line to read “We are also grateful to a team of anonymous **DOE and national lab** reviewers whose input helped improve the final report.”?

Best,
Travis

PS – the Secretary is reading the updated version tonight.

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 24, 2025 7:12 PM
To: John Christy <climateman60@gmail.com>
Cc: Roy Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Didn't even think of that. If it comes up I guess I'd invoke McShane and Weiner and say the ipcc has never addressed the problem that the millennial scale back-projections at the hemispheric scale are simply too noisy to draw precise conclusions. And nothing core to our report depended one way or the other on those issues.

On Thu, Jul 24, 2025 at 5:59 PM John Christy <climateman60@gmail.com> wrote:

Ross

What will you say when asked why we didn't deal with all the Hockey Sticks from Mann to IPCC AR6?

John C.

Sent from my iPhone

On Jul 24, 2025, at 4:27 PM, Roy Spencer <roywspencer@hotmail.com> wrote:

Dang! Wish I had thought of getting a Canadian phone!

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 24, 2025 3:20 PM
To: Steven Koonin <steven.koonin@gmail.com>

Cc: John Christy <climateman60@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

They'll have trouble reaching me. My Canadian phone died and I won't be able to replace it and reactivate the number until I'm back home next week.

On Thu, Jul 24, 2025 at 3:14 PM Steven Koonin <steven.koonin@gmail.com> wrote:

Got a similar call from Scott Waldmann, but let it go to voicemail.

He said he wanted to speak on background about the endangerment finding and "some of the climate work" I've been doing.

I've not returned his call.

From: John Christy <climateman60@gmail.com>
Sent: Thursday, July 24, 2025 3:13 PM
To: Ross McKittrick <ross.mckitrick@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Just got a call from Scott at Politico stating he knows I've worked on a DoE science report. I said no comment. (He uses his wife's cell phone, so you don't know it's him when you pick up.)

John C.

On Thu, Jul 24, 2025 at 10:24 AM Ross McKittrick <ross.mckitrick@gmail.com> wrote:

PS here are the report files now with a hi-res logo included.

On Thu, Jul 24, 2025 at 10:18 AM Steven Koonin <steven.koonin@gmail.com> wrote:

If you want to shorten Ross' proposed title a bit, it could read:

Department of Energy Releases Report That Challenges Common Misconceptions About Climate Change

From: Ross McKittrick <ross.mckitrick@gmail.com>
Sent: Thursday, July 24, 2025 11:16 AM
To: John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis

<travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>;
Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

The title might need to say a bit more, e.g.

Department of Energy Releases Report of Climate Working Group That Challenges Common
Misconceptions About Climate Change

and then the remainder follows the text as shown in Steve's edit, with "renowned" replaced by "eminent",
there being no adjective that won't draw scorn from the haters, but we might as well pick the word we like
best.

On Thu, Jul 24, 2025 at 9:09 AM John Christy <climateman60@gmail.com> wrote:

Looks good.

John C.

On Thu, Jul 24, 2025 at 8:59 AM Steven Koonin <steven.koonin@gmail.com> wrote:

And with me.

SEK

From: Roy Spencer <roywspencer@hotmail.com>

Sent: Thursday, July 24, 2025 9:49 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>; Steven Koonin <steven.koonin@gmail.com>; 'John Christy'
<climateman60@gmail.com>

Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua
<Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Fine with me.

Sent from my Verizon, Samsung Galaxy smartphone

----- Original message -----

From: "Fisher, Travis" <travis.fisher@hq.doe.gov>

Date: 7/24/25 8:42 AM (GMT-06:00)

To: Steven Koonin <steven.koonin@gmail.com>, 'John Christy' <climateman60@gmail.com>

Cc: 'Judith Curry' <curry.judith@gmail.com>, 'Ross McKittrick' <ross.mckittrick@gmail.com>, "Loucks, Joshua" <Joshua.Loucks@hq.doe.gov>, "Cohen, Seth" <seth.cohen@hq.doe.gov>, "Roy W. Spencer" <roywspencer@hotmail.com>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Ok I'm going with eminent for now. All of this is subject to review by the office of public affairs, so we'll see what sticks.

In this next iteration I added your bios at the end (straight copy/paste from the report). Are you all comfortable if we decide to take that approach?

From: Steven Koonin <steven.koonin@gmail.com>

Sent: Thursday, July 24, 2025 9:21 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>

Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'Roy W. Spencer' <roywspencer@hotmail.com>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Perhaps "eminent" .

Although in some eyes, "notorious" would be appropriate.

From: Fisher, Travis <travis.fisher@hq.doe.gov>

Sent: Thursday, July 24, 2025 9:07 AM

To: John Christy <climateman60@gmail.com>

Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

I don't want anyone to think of you all as circus performers! I was looking for a synonym for "distinguished," with the bonus of being well-known. I'm open to ideas: [RENOWNED Synonyms: 104 Similar and Opposite Words | Merriam-Webster Thesaurus](#)

From: John Christy <climateman60@gmail.com>

Sent: Thursday, July 24, 2025 8:53 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>

Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth

<seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>

Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis

“Renowned” sounds a little like a circus performer.

I wonder how many reporters will actually read the whole document? I suspect one of our common answers to their questions is, “just read the report!”

John C.

Sent from my iPhone

On Jul 23, 2025, at 8:28 PM, Fisher, Travis <travis.fisher@hq.doe.gov> wrote:

It’s a heck of a report. I can’t wait to see where we go from here!

Here’s my first attempt at a press release—please make any edits or suggestions you like.

From: Steven Koonin <steven.koonin@gmail.com>

Sent: Wednesday, July 23, 2025 5:50 PM

To: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckitrick@gmail.com>

Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'John Christy' <climatemanager60@gmail.com>; 'Roy W. Spencer' <roywspencer@hotmail.com>

Subject: [EXTERNAL] RE: July 23rd version - final (?)

Second that! Amazing effort, Ross.

SEK

From: Judith Curry <curry.judith@gmail.com>

Sent: Wednesday, July 23, 2025 5:49 PM

To: Ross McKittrick <ross.mckitrick@gmail.com>

Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; John Christy <climatemanager60@gmail.com>; Roy W. Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>

Subject: Re: July 23rd version - final (?)

congrats to us all, and special thanks to RM for pulling it all together

On Wed, Jul 23, 2025 at 2:19 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

Attached in MSWord and PDF formats.

All of Steve's corrections incorporated, also John's stray L and everything else I'm aware of.

The only remaining correction I have not been able to make is using a high-res version of the DoE logo.
If that becomes the main focus of public criticism I will say we did well.

Ross

--

<image001.jpg>

Judith Curry, President
CFAN - Climate Forecast Applications
Network
Reno, NV USA
curry.judith@cfanclimate.com |
+1.404.803.2012
<http://www.cfanclimate.net>

This message does not originate from a known Department of Energy email system.
Use caution if this message contains attachments, links or requests for information.

<CWG Press Release.docx>

DOCUMENT WITHHELD FOR PRIVILEGE

From: John Christy <climateman60@gmail.com>
Sent: Monday, July 28, 2025 1:14 PM
To: Ross McKitrick
Cc: Roy Spencer; Fisher, Travis; Steven Koonin; Judith Curry; Loucks, Joshua; Cohen, Seth
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

If Ross likes it, I'm good.

John C.

On Mon, Jul 28, 2025 at 9:59 AM Ross McKitrick <ross.mckitrick@gmail.com> wrote:
Looks good to me.

On Mon, Jul 28, 2025 at 9:57 AM Roy Spencer <roywspencer@hotmail.com> wrote:
Fine with me.
-Roy

Sent from my Verizon, Samsung Galaxy smartphone

----- Original message -----

From: "Fisher, Travis" <travis.fisher@hq.doe.gov>
Date: 7/28/25 9:46 AM (GMT-06:00)
To: Roy Spencer <roywspencer@hotmail.com>, Ross McKitrick <ross.mckitrick@gmail.com>, John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>, Judith Curry <curry.judith@gmail.com>, "Loucks, Joshua" <Joshua.Loucks@hq.doe.gov>, "Cohen, Seth" <seth.cohen@hq.doe.gov>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Hi all,

This is a matter of administrative housekeeping—I want to make sure you're all aware of the process and comfortable with it.

The attached memo presents an action item to the Secretary (to publish the report). It's a formality, but please let me know if you'd like to see any changes to the text.

Best,

Travis

From: Roy Spencer <roywspencer@hotmail.com>

Sent: Monday, July 28, 2025 9:21 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>; Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>

Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

In my experience, any of the right-leaning news outlets have always treated me well. But (unlike John), I'm no longer willing to talk to the "usual suspects"... CNN, MSNBC, NPR and its derivatives, and the British press

Years ago the science reporter at BBC, Richard Black, put some *ad hominem* in his report on my interview, and he took a lot of flak for it from readers... he tried to get me to do a follow-up to save face, and I said sorry, buddy, you had your chance.

-Roy

From: Fisher, Travis <travis.fisher@hq.doe.gov>

Sent: Monday, July 28, 2025 8:05 AM

To: Roy Spencer <roywspencer@hotmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>

Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Hi Roy,

The report should be released tomorrow, likely in the afternoon. I'll make sure everyone here is in the loop when it goes out. I believe the Secretary is aware that he owes you all some glowing public comments and an open bar tab at some point.

On media issues, my preference is to let the professionals take the lead. I suggest you forward media requests to Andrea Woods in DOE's Office of Public Affairs so she's aware of the questions:

andrea.woods@hq.doe.gov

That said, people will want to talk to the CWG authors. If you're willing to talk to reporters, just let Andrea know. She should have a good sense of which ones are coming to this issue in good faith and can set up interviews.

Best,

Travis

From: Roy Spencer <roywspencer@hotmail.com>

Sent: Monday, July 28, 2025 6:01 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>; Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>

Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis:

Any word on when the Report will be released?

I'm leaning toward not responding to any media queries, unless others think it's needed at some point.

-Roy

From: Fisher, Travis <travis.fisher@hq.doe.gov>

Sent: Friday, July 25, 2025 3:52 PM

To: Ross McKittrick <ross.mckittrick@gmail.com>; John Christy <climateman60@gmail.com>

Cc: Roy Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Hi all,

I have a request for a minor change in the preface. The last line presently reads: “We are also grateful to a team of anonymous peer reviewers whose input helped improve the final report.”

Would it cause any heartburn to change the line to read “We are also grateful to a team of anonymous **DOE and national lab** reviewers whose input helped improve the final report.”?

Best,

Travis

PS – the Secretary is reading the updated version tonight.

From: Ross McKittrick <ross.mckittrick@gmail.com>

Sent: Thursday, July 24, 2025 7:12 PM

To: John Christy <climateman60@gmail.com>

Cc: Roy Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>

Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Didn't even think of that. If it comes up I guess I'd invoke McShane and Weiner and say the ipcc has never addressed the problem that the millennial scale back-projections at the hemispheric scale are simply too noisy to draw precise conclusions. And nothing core to our report depended one way or the other on those issues.

On Thu, Jul 24, 2025 at 5:59 PM John Christy <climateman60@gmail.com> wrote:

Ross

What will you say when asked why we didn't deal with all the Hockey Sticks from Mann to IPCC AR6?

John C.

Sent from my iPhone

On Jul 24, 2025, at 4:27 PM, Roy Spencer <roywspencer@hotmail.com> wrote:

Dang! Wish I had thought of getting a Canadian phone!

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 24, 2025 3:20 PM
To: Steven Koonin <steven.koonin@gmail.com>
Cc: John Christy <climateman60@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

They'll have trouble reaching me. My Canadian phone died and I won't be able to replace it and reactivate the number until I'm back home next week.

On Thu, Jul 24, 2025 at 3:14 PM Steven Koonin <steven.koonin@gmail.com> wrote:

Got a similar call from Scott Waldmann, but let it go to voicemail.

He said he wanted to speak on background about the endangerment finding and “some of the climate work” I’ve been doing.

I’ve not returned his call.

From: John Christy <climateman60@gmail.com>
Sent: Thursday, July 24, 2025 3:13 PM
To: Ross McKittrick <ross.mckittrick@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Just got a call from Scott at Politico stating he knows I've worked on a DoE science report. I said no comment. (He uses his wife's cell phone, so you don't know it's him when you pick up.)

John C.

On Thu, Jul 24, 2025 at 10:24 AM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

PS here are the report files now with a hi-res logo included.

On Thu, Jul 24, 2025 at 10:18 AM Steven Koonin <steven.koonin@gmail.com> wrote:

If you want to shorten Ross’ proposed title a bit, it could read:

Department of Energy Releases Report That Challenges Common Misconceptions About Climate Change

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 24, 2025 11:16 AM
To: John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

The title might need to say a bit more, e.g.

Department of Energy Releases Report of Climate Working Group That Challenges
Common Misconceptions About Climate Change

and then the remainder follows the text as shown in Steve's edit, with "renowned" replaced by "eminent", there being no adjective that won't draw scorn from the haters, but we might as well pick the word we like best.

On Thu, Jul 24, 2025 at 9:09 AM John Christy <climateman60@gmail.com> wrote:

Looks good.

John C.

On Thu, Jul 24, 2025 at 8:59 AM Steven Koonin <steven.koonin@gmail.com> wrote:

And with me.

SEK

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Thursday, July 24, 2025 9:49 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; Steven Koonin <steven.koonin@gmail.com>; 'John Christy' <climateman60@gmail.com>
Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Fine with me.

Sent from my Verizon, Samsung Galaxy smartphone

----- Original message -----

From: "Fisher, Travis" <travis.fisher@hq.doe.gov>

Date: 7/24/25 8:42 AM (GMT-06:00)

To: Steven Koonin <steven.koonin@gmail.com>, 'John Christy' <climateman60@gmail.com>

Cc: 'Judith Curry' <curry.judith@gmail.com>, 'Ross McKittrick' <ross.mckittrick@gmail.com>, "Loucks, Joshua" <Joshua.Loucks@hq.doe.gov>, "Cohen, Seth" <seth.cohen@hq.doe.gov>, "Roy W. Spencer" <roywspencer@hotmail.com>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Ok I'm going with eminent for now. All of this is subject to review by the office of public affairs, so we'll see what sticks.

In this next iteration I added your bios at the end (straight copy/paste from the report). Are you all comfortable if we decide to take that approach?

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Thursday, July 24, 2025 9:21 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>
Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'Roy W. Spencer' <roywspencer@hotmail.com>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Perhaps “eminent” .

Although in some eyes, “notorious” would be appropriate.

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Thursday, July 24, 2025 9:07 AM
To: John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

I don't want anyone to think of you all as circus performers! I was looking for a synonym for “distinguished,” with the bonus of being well-known. I'm open to ideas: [RENOWNED Synonyms: 104 Similar and Opposite Words | Merriam-Webster Thesaurus](#)

From: John Christy <climateman60@gmail.com>
Sent: Thursday, July 24, 2025 8:53 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis

“Renowned” sounds a little like a circus performer.

I wonder how many reporters will actually read the whole document? I suspect one of our common answers to their questions is, “just read the report!”

John C.

Sent from my iPhone

On Jul 23, 2025, at 8:28 PM, Fisher, Travis <travis.fisher@hq.doe.gov> wrote:

It’s a heck of a report. I can’t wait to see where we go from here!

Here’s my first attempt at a press release—please make any edits or suggestions you like.

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Wednesday, July 23, 2025 5:50 PM
To: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>; 'Roy W. Spencer' <roywspencer@hotmail.com>
Subject: [EXTERNAL] RE: July 23rd version - final (?)

Second that! Amazing effort, Ross.

SEK

From: Judith Curry <curry.judith@gmail.com>
Sent: Wednesday, July 23, 2025 5:49 PM

To: Ross McKittrick <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua
<Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; John
Christy <climateman60@gmail.com>; Roy W. Spencer
<roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>
Subject: Re: July 23rd version - final (?)

congrats to us all, and special thanks to RM for pulling it all together

On Wed, Jul 23, 2025 at 2:19 PM Ross McKittrick
<ross.mckittrick@gmail.com> wrote:


Attached in MSWord and PDF formats.

All of Steve's corrections incorporated, also John's stray L and
everything else I'm aware of.


The only remaining correction I have not been able to make is using a
high-res version of the DoE logo. If that becomes the main focus of
public criticism I will say we did well.

Ross

--

 Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

This message does not originate from a known Department of Energy
email system.
Use caution if this message contains attachments, links or requests for
information.



<CWG Press Release.docx>

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Thursday, July 24, 2025 9:42 AM
To: Steven Koonin; 'John Christy'
Cc: 'Judith Curry'; 'Ross McKittrick'; Loucks, Joshua; Cohen, Seth; 'Roy W. Spencer'
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)
Attachments: CWG Press Release v2.docx

Ok I'm going with eminent for now. All of this is subject to review by the office of public affairs, so we'll see what sticks.

In this next iteration I added your bios at the end (straight copy/paste from the report). Are you all comfortable if we decide to take that approach?

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Thursday, July 24, 2025 9:21 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>
Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckitrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'Roy W. Spencer' <roywspencer@hotmail.com>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Perhaps "eminent" .

Although in some eyes, "notorious" would be appropriate.

From: Fisher, Travis <travis.fisher@hq.doe.gov>
Sent: Thursday, July 24, 2025 9:07 AM
To: John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckitrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

I don't want anyone to think of you all as circus performers! I was looking for a synonym for "distinguished," with the bonus of being well-known. I'm open to ideas: [RENOWNED Synonyms: 104 Similar and Opposite Words | Merriam-Webster Thesaurus](#)

From: John Christy <climateman60@gmail.com>
Sent: Thursday, July 24, 2025 8:53 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>
Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckitrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis

“Renowned” sounds a little like a circus performer.

I wonder how many reporters will actually read the whole document? I suspect one of our common answers to their questions is, “just read the report!”

John C.

Sent from my iPhone

On Jul 23, 2025, at 8:28 PM, Fisher, Travis <travis.fisher@hq.doe.gov> wrote:

It’s a heck of a report. I can’t wait to see where we go from here!

Here’s my first attempt at a press release—please make any edits or suggestions you like.

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Wednesday, July 23, 2025 5:50 PM
To: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>; 'Roy W. Spencer' <roywspencer@hotmail.com>
Subject: [EXTERNAL] RE: July 23rd version - final (?)

Second that! Amazing effort, Ross.

SEK

From: Judith Curry <curry.judith@gmail.com>
Sent: Wednesday, July 23, 2025 5:49 PM
To: Ross McKittrick <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; John Christy <climateman60@gmail.com>; Roy W. Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>
Subject: Re: July 23rd version - final (?)

congrats to us all, and special thanks to RM for pulling it all together


On Wed, Jul 23, 2025 at 2:19 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

Attached in MSWord and PDF formats.

All of Steve's corrections incorporated, also John's stray L and everything else I'm aware of. The only remaining correction I have not been able to make is using a high-res version of the DoE logo. If that becomes the main focus of public criticism I will say we did well.

Ross

--



Judith Curry, President
CFAN - Climate Forecast Applications Network
Reno, NV USA
curry.judith@cfanclimate.com | +1.404.803.2012
<http://www.cfanclimate.net>

This message does not originate from a known Department of Energy email system.
Use caution if this message contains attachments, links or requests for information.

<CWG Press Release.docx>

Department of Energy Releases Report from Distinguished Climate Working Group

The Department of Energy's Climate Working Group report highlights key differences between the scientific literature and the conventional narrative on climate change

Energy.gov

July 29, 2025

The Department of Energy's Climate Working Group report highlights key differences between the scientific literature and the conventional narrative on climate change

WASHINGTON—The U.S. Department of Energy (DOE) today released its report “A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate,” authored by five eminent climate experts assembled by Secretary of Energy Chris Wright.

From Secretary Wright's foreword:

“The rise of human flourishing over the past two centuries is a story worth celebrating. Yet we are told—relentlessly—that the very energy systems that enabled this progress now pose an existential threat. Hydrocarbon-based fuels, the argument goes, must be rapidly abandoned or else we risk planetary ruin. That view demands scrutiny. That's why I commissioned this report: to encourage a more thoughtful and science-based conversation about climate change and energy.”

Using peer-reviewed scientific literature and government sources of data, the report:

- Reviews scientific certainties and uncertainties in how anthropogenic emissions of carbon dioxide (CO₂) and other greenhouse gases have affected, or will affect, the Nation's climate, extreme weather events, and metrics of societal well-being.
- Assesses the near-term impacts of elevated concentrations of CO₂, including enhanced plant growth and reduced ocean alkalinity.
- Evaluates data and projections regarding long-term impacts of elevated concentrations of CO₂, including estimates of future warming.
- Finds that claims of increased frequency or intensity of hurricanes, tornadoes, floods, and droughts are not supported by U.S. historical data.
- Asserts that CO₂-induced warming might be less damaging economically than commonly believed, and that aggressive mitigation policies could prove more detrimental than beneficial.

The DOE report is intended to add clarity and balance to the debate over climate policy in the United States and abroad. In the spirit of open public discourse, a notice of availability and public comment will be published in the Federal Register—all interested parties are encouraged to submit comments.

The Climate Working Group consists of the following climate experts:

John Christy, Ph.D. is the Distinguished Professor of Atmospheric and Earth Sciences and Alabama's State Climatologist at the University of Alabama in Huntsville. He has a B.A. (Mathematics) from CSU Fresno and an M.S. and Ph.D. (Atmospheric Sciences) from the University of Illinois in Urbana-Champaign. He is a Fellow of the American Meteorological Society and recipient of The NASA Medal for Exceptional Scientific Achievement for constructing, with Dr. Roy Spencer, the first global atmospheric temperature dataset from polar orbiting satellites. He served as Lead Author of the 3rd IPCC Assessment Report (2001), he coauthored the National Academy of Science expert report on Surface Temperature Reconstructions (2006), and he has served on the NOAA Climate Change Science Program, NASA's Earth Science Subcommittee, the National Research Council Space Studies Board and the EPA Science Advisory Board. In addition, Dr. Christy has given testimony in 20 Congressional Hearings.

Judith Curry, Ph.D. is Professor Emerita at the Georgia Institute of Technology, where she served as Chair of Earth and Atmospheric Sciences for 13 years. She is President and co-founder of Climate Forecast Applications Network (CFAN). Curry has a PhD in Geophysical Sciences from the University of Chicago. She has authored or coauthored 192 peer-reviewed papers in atmospheric and climate sciences, two textbooks, and most recently the book *Climate Uncertainty and Risk*. Curry is a Fellow of the American Meteorological Society, the American Association for the Advancement of Science, American Geophysical Union, and a Member of the American Academy of Sciences and Letters. She has served on several Science Steering Committees of the World Climate Research Program and also the DOE Biological & Environmental Research Advisory Committee, Earth Science Subcommittee of the NASA Advisory Council, and the NRC Space Studies Board and Climate Research Committee. Dr. Curry has given testimony in 13 Congressional Hearings.

The Honorable Steven E. Koonin, Ph.D. is the Edward Teller Senior Fellow at Stanford's Hoover Institution. He has served as Under Secretary for Science at the Department of Energy (2009-2011), as Chief Scientist for British Petroleum (2004-2009), and as a professor at Caltech (1975 – 2004, the last nine years as Vice President and Provost). Koonin is a member of the National Academy of Sciences and the JASON group of government advisors, as well as a Governor of Lawrence Livermore National Laboratory, having served in similar capacities for the Los Alamos, Sandia, Brookhaven, and Argonne National Laboratories. Other prior roles include a University

Professor at New York University, a non-resident Senior Fellow at the American Enterprise Institute, and a Trustee of the Institute for Defense Analyses. Koonin has a BS in Physics from Caltech and a PhD in Theoretical Physics from Massachusetts Institute of Technology. He authored the 2021 bestseller *Unsettled: What Climate Science Tells Us, What It Doesn't, and Why It Matters*, the 1985 textbook *Computational Physics*, and some 200 peer-reviewed papers in physics and astrophysics, scientific computation, energy technology and policy, and climate science.

Ross McKittrick, Ph.D., is a Professor of Environmental Economics at the University of Guelph in Ontario, Canada. He holds a BA in economics from Queen's University, and an MA and Ph.D. in economics from the University of British Columbia. He has published widely in both economics and physical science journals on topics related to climate change, pollution and public policy. His book *Economic Analysis of Environmental Policy* was published by the University of Toronto Press in 2010. His background in applied statistics has led him to author or coauthor peer-reviewed publications across a wide range of topics in the physical sciences including paleoclimate reconstruction, malaria transmission, surface temperature measurement, climate attribution methodology and climate model evaluation. Professor McKittrick has made many invited academic presentations around the world, was an invited contributor to the 2006 National Academy of Sciences report on Surface Temperature Reconstructions, has served as an expert reviewer for the last three IPCC Assessment Reports (Working Groups I and II) and has testified before the U.S. Congress and committees of the Canadian House of Commons and Senate.

Roy W. Spencer, Ph.D., is a Principal Research Scientist at the University of Alabama in Huntsville. He holds a BS in Atmospheric and Oceanic Science from the University of Michigan, and MS and PhD in Meteorology from the University of Wisconsin -Madison. Spencer's published research has focused on monitoring of weather processes and climate change from Earth-orbiting satellites, as well as using simple physical models for diagnosing climate sensitivity and climate feedbacks. As a NASA Senior Scientist for Climate Studies he served as the U.S. Science Team Leader for the Advanced Microwave Scanning Radiometer (AMSR-E) flying on the Aqua satellite, providing our most accurate global measurements of sea surface temperatures, sea ice, and several other features of the climate system. Along with Dr. John Christy, Spencer developed the first satellite-based technique for global temperature monitoring, for which they received NASA's Exceptional Scientific Achievement Medal and the American Meteorological Society's Special Award.

From: Ross McKittrick [ross.mckittrick@gmail.com]
Sent: 7/24/2025 3:23:47 PM
To: Steven Koonin [steven.koonin@gmail.com]
CC: John Christy [climateman60@gmail.com]; Roy Spencer [roywspencer@hotmail.com]; Fisher, Travis [travis.fisher@hq.doe.gov]; Judith Curry [curry.judith@gmail.com]; Loucks, Joshua [Joshua.Loucks@hq.doe.gov]; Cohen, Seth [seth.cohen@hq.doe.gov]
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)
Attachments: DoE.Report.July23.pdf; DoE.Report.July23.docx

PS here are the report files now with a hi-res logo included.

On Thu, Jul 24, 2025 at 10:18 AM Steven Koonin <steven.koonin@gmail.com> wrote:

If you want to shorten Ross' proposed title a bit, it could read:

Department of Energy Releases Report That Challenges Common Misconceptions About Climate Change

From: Ross McKittrick <ross.mckittrick@gmail.com>
Sent: Thursday, July 24, 2025 11:16 AM
To: John Christy <climateman60@gmail.com>
Cc: Steven Koonin <steven.koonin@gmail.com>; Roy Spencer <roywspencer@hotmail.com>; Fisher, Travis <travis.fisher@hq.doe.gov>; Judith Curry <curry.judith@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

The title might need to say a bit more, e.g.

Department of Energy Releases Report of Climate Working Group That Challenges Common Misconceptions About Climate Change

and then the remainder follows the text as shown in Steve's edit, with "renowned" replaced by "eminent", there being no adjective that won't draw scorn from the haters, but we might as well pick the word we like best.

On Thu, Jul 24, 2025 at 9:09 AM John Christy <climateman60@gmail.com> wrote:

Looks good.

John C.

On Thu, Jul 24, 2025 at 8:59 AM Steven Koonin <steven.koonin@gmail.com> wrote:

And with me.

SEK

From: Roy Spencer <roywspencer@hotmail.com>
Sent: Thursday, July 24, 2025 9:49 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; Steven Koonin <steven.koonin@gmail.com>; 'John Christy' <climateman60@gmail.com>
Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Fine with me.

Sent from my Verizon. Samsung Galaxy smartphone

----- Original message -----

From: "Fisher, Travis" <travis.fisher@hq.doe.gov>
Date: 7/24/25 8:42 AM (GMT-06:00)
To: Steven Koonin <steven.koonin@gmail.com>, 'John Christy' <climateman60@gmail.com>
Cc: 'Judith Curry' <curry.judith@gmail.com>, 'Ross McKittrick' <ross.mckittrick@gmail.com>, "Loucks, Joshua" <Joshua.Loucks@hq.doe.gov>, "Cohen, Seth" <seth.cohen@hq.doe.gov>, "Roy W. Spencer" <roywspencer@hotmail.com>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Ok I'm going with eminent for now. All of this is subject to review by the office of public affairs, so we'll see what sticks.

In this next iteration I added your bios at the end (straight copy/paste from the report). Are you all comfortable if we decide to take that approach?

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Thursday, July 24, 2025 9:21 AM
To: Fisher, Travis <travis.fisher@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>
Cc: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'Roy W. Spencer' <roywspencer@hotmail.com>
Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

Perhaps “eminent” .

Although in some eyes, “notorious” would be appropriate.

From: Fisher, Travis <travis.fisher@hq.doe.gov>

Sent: Thursday, July 24, 2025 9:07 AM

To: John Christy <climateman60@gmail.com>

Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>

Subject: RE: [EXTERNAL] RE: July 23rd version - final (?)

I don’t want anyone to think of you all as circus performers! I was looking for a synonym for “distinguished,” with the bonus of being well-known. I’m open to ideas: [RENOWNED Synonyms: 104 Similar and Opposite Words | Merriam-Webster Thesaurus](#)

From: John Christy <climateman60@gmail.com>

Sent: Thursday, July 24, 2025 8:53 AM

To: Fisher, Travis <travis.fisher@hq.doe.gov>

Cc: Steven Koonin <steven.koonin@gmail.com>; Judith Curry <curry.judith@gmail.com>; Ross McKittrick <ross.mckittrick@gmail.com>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; Roy W. Spencer <roywspencer@hotmail.com>

Subject: Re: [EXTERNAL] RE: July 23rd version - final (?)

Travis

“Renowned” sounds a little like a circus performer.

I wonder how many reporters will actually read the whole document? I suspect one of our common answers to their questions is, “just read the report!”

John C.

Sent from my iPhone

On Jul 23, 2025, at 8:28 PM, Fisher, Travis <travis.fisher@hq.doe.gov> wrote:

It’s a heck of a report. I can’t wait to see where we go from here!

Here’s my first attempt at a press release—please make any edits or suggestions you like.

From: Steven Koonin <steven.koonin@gmail.com>
Sent: Wednesday, July 23, 2025 5:50 PM
To: 'Judith Curry' <curry.judith@gmail.com>; 'Ross McKittrick' <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; 'John Christy' <climateman60@gmail.com>; 'Roy W. Spencer' <roywspencer@hotmail.com>
Subject: [EXTERNAL] RE: July 23rd version - final (?)

Second that! Amazing effort, Ross.

SEK

From: Judith Curry <curry.judith@gmail.com>
Sent: Wednesday, July 23, 2025 5:49 PM
To: Ross McKittrick <ross.mckittrick@gmail.com>
Cc: Fisher, Travis <travis.fisher@hq.doe.gov>; Loucks, Joshua <Joshua.Loucks@hq.doe.gov>; Cohen, Seth <seth.cohen@hq.doe.gov>; John Christy <climateman60@gmail.com>; Roy W. Spencer <roywspencer@hotmail.com>; Steven Koonin <steven.koonin@gmail.com>
Subject: Re: July 23rd version - final (?)

congrats to us all, and special thanks to RM for pulling it all together

On Wed, Jul 23, 2025 at 2:19 PM Ross McKittrick <ross.mckittrick@gmail.com> wrote:

Attached in MSWord and PDF formats.

All of Steve's corrections incorporated, also John's stray L and everything else I'm aware of.

The only remaining correction I have not been able to make is using a high-res version of the DoE logo. If that becomes the main focus of public criticism I will say we did well.

Ross

--

<image001.jpg>

Judith Curry, President
CFAN - Climate Forecast Applications
Network
Reno, NV USA
curry.judith@cfanclimate.com |
+1.404.803.2012
<http://www.cfanclimate.net>

This message does not originate from a known Department of Energy email system.

Use caution if this message contains attachments, links or requests for information.

<CWG Press Release.docx>



U.S. DEPARTMENT *of* **ENERGY**

A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate

Climate Working Group
United States Department of Energy
July 23, 2025

A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate

Report to U.S. Energy Secretary Christopher Wright
July 23, 2025

Climate Working Group:

John Christy, Ph.D.

Judith Curry, Ph.D.

Steven Koonin, Ph.D.

Ross McKittrick, Ph.D.

Roy Spencer, Ph.D.

Copyright © 2025 United States

Suggested citation:

Climate Working Group (2025) A Critical Review of Impacts of Greenhouse Gas Emissions on the U.S. Climate. Washington DC: Department of Energy, July 23, 2025

Table of Contents

SECRETARY’S FOREWORD	VII
EXECUTIVE SUMMARY	VIII
PREFACE.....	IX
PART I: DIRECT HUMAN INFLUENCE ON ECOSYSTEMS AND THE CLIMATE.....	1
1 CARBON DIOXIDE AS A POLLUTANT	2
2 DIRECT IMPACTS OF CO ₂ ON THE ENVIRONMENT	3
2.1 CO ₂ AS A CONTRIBUTOR TO GLOBAL GREENING	3
2.1.1 <i>Measurement of global greening</i>	3
2.1.2 <i>Photosynthesis and CO₂ levels</i>	4
2.1.3 <i>Rising CO₂ and crop water use efficiency</i>	6
2.1.4 <i>CO₂ fertilization benefits in IPCC Reports</i>	6
2.2 THE ALKALINE OCEANS	6
2.2.1 <i>Changing pH</i>	6
2.2.2 <i>Coral reef changes</i>	7
3 HUMAN INFLUENCES ON THE CLIMATE.....	11
3.1 COMPONENTS OF RADIATIVE FORCING AND THEIR HISTORY.....	11
3.1.1 <i>Historical radiative forcing</i>	11
3.1.2 <i>Change in atmospheric CO₂ since 1958</i>	13
3.2 FUTURE EMISSION SCENARIOS AND THE CARBON CYCLE.....	14
3.2.1 <i>Emission scenarios</i>	14
3.2.2 <i>The carbon cycle relating emissions and concentrations</i>	17
3.3 URBANIZATION INFLUENCE ON TEMPERATURE TRENDS	20
PART II: CLIMATE RESPONSE TO CO ₂ EMISSIONS.....	24
4 CLIMATE SENSITIVITY TO CO ₂ FORCING.....	25
4.1 INTRODUCTION.....	25
4.2 MODEL-BASED ESTIMATES OF CLIMATE SENSITIVITY	26
4.3 DATA-DRIVEN ESTIMATES OF CLIMATE SENSITIVITY.....	27
4.4 TRANSIENT CLIMATE RESPONSE	29
5 DISCREPANCIES BETWEEN MODELS AND INSTRUMENTAL OBSERVATIONS.....	31
5.1 INTRODUCTION.....	31
5.2 SURFACE WARMING	33
5.3 TROPOSPHERIC WARMING	34
5.4 VERTICAL TEMPERATURE PROFILE MISMATCH	36

5.5	STRATOSPHERIC COOLING	38
5.6	SNOW COVER MISMATCH	39
5.7	HEMISPHERIC SYMMETRY OF THE PLANETARY ALBEDO	41
5.8	U.S. CORN BELT	42
6	EXTREME WEATHER	46
6.1	INTRODUCTION.....	46
6.2	HURRICANES AND TROPICAL CYCLONES.....	48
6.3	TEMPERATURE EXTREMES.....	52
6.3.1	<i>Temperatures in the U.S. are becoming less extreme</i>	<i>53</i>
6.3.2	<i>Exceedances of a heat threshold.....</i>	<i>56</i>
6.3.3	<i>Heatwaves.....</i>	<i>57</i>
6.4	EXTREME PRECIPITATION	61
6.5	TORNADOES	66
6.6	FLOODING.....	67
6.7	DROUGHTS.....	68
6.8	WILDFIRES.....	69
7	CHANGES IN SEA LEVEL	75
7.1	GLOBAL SEA LEVEL RISE	75
7.2	U.S. SEA LEVEL RISE.....	75
7.3	PROJECTED SEA LEVEL RISE.....	79
8	UNCERTAINTIES IN CLIMATE CHANGE ATTRIBUTION	82
8.1	INTRODUCTION.....	82
8.2	ATTRIBUTION METHODS	83
8.3	ATTRIBUTION OF GLOBAL WARMING	84
8.3.1	<i>Natural climate variability</i>	<i>84</i>
8.3.2	<i>Optimal fingerprinting</i>	<i>88</i>
8.3.3	<i>Time series methods.....</i>	<i>89</i>
8.4	DECLINING PLANETARY ALBEDO AND RECENT RECORD WARMTH	90
8.5	ATTRIBUTION OF CLIMATE IMPACT DRIVERS.....	92
8.6	EXTREME EVENT ATTRIBUTION (EEA).....	95
8.6.1	<i>Case study – 2021 Western North America heat wave.....</i>	<i>96</i>
	PART III: IMPACTS ON ECOSYSTEMS AND SOCIETY.....	103
9	CLIMATE CHANGE AND U.S. AGRICULTURE	104
9.1	ECONOMETRIC ANALYSES	104
9.2	FIELD AND LABORATORY STUDIES OF CO ₂ ENRICHMENT	105

9.3	CROP MODELING META-ANALYSES.....	106
9.4	CO ₂ FERTILIZATION AND NUTRIENT LOSS.....	107
10	MANAGING RISKS OF EXTREME WEATHER.....	110
10.1	SOCIOECONOMIC CONTEXT.....	110
10.2	DATA CHALLENGES.....	111
10.3	MORTALITY FROM TEMPERATURE EXTREMES.....	111
10.3.1	<i>Heat and cold risks.....</i>	<i>111</i>
10.3.2	<i>Mortality risks and energy costs.....</i>	<i>113</i>
11	CLIMATE CHANGE, THE ECONOMY, AND THE SOCIAL COST OF CARBON	116
11.1	CLIMATE CHANGE AND ECONOMIC GROWTH	116
11.1.1	<i>Overview.....</i>	<i>116</i>
11.1.2	<i>Empirical analysis of climate change and economic growth</i>	<i>119</i>
11.2	MODELS OF THE SOCIAL COST OF CARBON	120
11.2.1	<i>Estimating the SCC.....</i>	<i>121</i>
11.2.2	<i>Variations in the SCC.....</i>	<i>122</i>
11.2.3	<i>Evidence for low SCC.....</i>	<i>123</i>
11.2.4	<i>Tipping points</i>	<i>123</i>
11.2.5	<i>Are there alternatives?</i>	<i>125</i>
12	GLOBAL CLIMATE IMPACTS OF U.S. EMISSIONS POLICIES.....	129
12.1	THE SCALE PROBLEM	129
12.2	CASE STUDY: U.S. MOTOR VEHICLE EMISSIONS	129
12.3	CONCLUDING THOUGHTS	130
	GLOSSARY	132
	METADATA FOR FIGURES AND TABLES	136
	ABOUT THE AUTHORS.....	140

SECRETARY'S FOREWORD

Energy, Integrity, and the Power of Human Potential

Over my lifetime, I've had the privilege of working as an energy entrepreneur across a range of fields—nuclear, geothermal, natural gas, and more—and I now serve as Secretary of Energy under President Donald Trump. But above all, I'm a physical scientist who sees modern energy as nothing short of miraculous. It powers every aspect of modern life, drives every industry, and has made America an energy powerhouse with the ability to fuel global progress.

The rise of human flourishing over the past two centuries is a story worth celebrating. Yet we are told—relentlessly—that the very energy systems that enabled this progress now pose an existential threat. Hydrocarbon-based fuels, the argument goes, must be rapidly abandoned or else we risk planetary ruin.

That view demands scrutiny. That's why I commissioned this report: to encourage a more thoughtful and science-based conversation about climate change and energy. With my technical background, I've reviewed reports from the Intergovernmental Panel on Climate Change, the U.S. government's assessments, and the academic literature. I've also engaged with many climate scientists, including the authors of this report.

What I've found is that media coverage often distorts the science. Many people—even well-meaning ones—walk away with a view of climate change that is exaggerated or incomplete. To provide clarity, I asked a diverse team of independent experts to summarize the current state of climate science, with a focus on how it relates to the United States.

I didn't select these authors because we always agree—far from it. In fact, they may not always agree with each other. But I chose them for their rigor, honesty, and willingness to elevate the debate. I exerted no control over their conclusions. What you'll read are their words, drawn from the best available data and scientific assessments.

I've reviewed the report carefully and challenged its findings. I believe it faithfully represents the state of climate science today. Still, many readers may be surprised by its conclusions—which differ in important ways from the mainstream narrative. That's a sign of how far the public conversation has drifted from the science itself.

To correct course, we need open, respectful, and informed debate. That's why I'm inviting public comment on this report. Honest scrutiny and scientific transparency should be at the heart of our policymaking.

Climate change is real, and it deserves attention. But it is not the greatest threat facing humanity. That distinction belongs to global energy poverty. As someone who values data, I know that improving the human condition depends on expanding access to reliable, affordable energy. Climate change is a challenge—not a catastrophe. But misguided policies based on fear rather than facts could truly endanger human well-being.

We stand at the threshold of a new era of energy leadership. If we empower innovation rather than restrain it, America can lead the world in providing cleaner, more abundant energy—lifting billions out of poverty, strengthening our economy, and improving our environment along the way.

EXECUTIVE SUMMARY

This report reviews scientific certainties and uncertainties in how anthropogenic carbon dioxide (CO₂) and other greenhouse gas emissions have affected, or will affect, the Nation's climate, extreme weather events, and selected metrics of societal well-being. Those emissions are increasing the concentration of CO₂ in the atmosphere through a complex and variable carbon cycle, where some portion of the additional CO₂ persists in the atmosphere for centuries.

Elevated concentrations of CO₂ directly enhance plant growth, globally contributing to “greening” the planet and increasing agricultural productivity [Section 2.1, Chapter 9]. They also make the oceans less alkaline (lower the pH). That is possibly detrimental to coral reefs, although the recent rebound of the Great Barrier Reef suggests otherwise [Section 2.2].

Carbon dioxide also acts as a greenhouse gas, exerting a warming influence on climate and weather [Section 3.1]. Climate change projections require scenarios of future emissions. There is evidence that scenarios widely-used in the impacts literature have overstated observed and likely future emission trends [Section 3.1].

The world's several dozen global climate models offer little guidance on how much the climate responds to elevated CO₂, with the average surface warming under a doubling of the CO₂ concentration ranging from 1.8°C to 5.7°C [Section 4.2]. Data-driven methods yield a lower and narrower range [Section 4.3]. Global climate models generally run “hot” in their description of the climate of the past few decades – too much warming at the surface and too much amplification of warming in the lower- and mid-troposphere [Sections 5.2-5.4]. The combination of overly sensitive models and implausible extreme scenarios for future emissions yields exaggerated projections of future warming.

Most extreme weather events in the U.S. do not show long-term trends. Claims of increased frequency or intensity of hurricanes, tornadoes, floods, and droughts are not supported by U.S. historical data [Sections 6.1-6.7]. Additionally, forest management practices are often overlooked in assessing changes in wildfire activity [Section 6.8]. Global sea level has risen approximately 8 inches since 1900, but there are significant regional variations driven primarily by local land subsidence; U.S. tide gauge measurements in aggregate show no obvious acceleration in sea level rise beyond the historical average rate [Chapter 7].

Attribution of climate change or extreme weather events to human CO₂ emissions is challenged by natural climate variability, data limitations, and inherent model deficiencies [Chapter 8]. Moreover, solar activity's contribution to the late 20th century warming might be underestimated [Section 8.3.1].

Both models and experience suggest that CO₂-induced warming might be less damaging economically than commonly believed, and excessively aggressive mitigation policies could prove more detrimental than beneficial [Chapters 9, 10, Section 11.1]. Social Cost of Carbon estimates, which attempt to quantify the economic damage of CO₂ emissions, are highly sensitive to their underlying assumptions and so provide limited independent information [Section 11.2].

U.S. policy actions are expected to have undetectably small direct impacts on the global climate and any effects will emerge only with long delays [Chapter 12].

PREFACE

This document originated in late March 2025 when Secretary Wright assembled an independent group to write a report on issues in climate science relevant for energy policymaking, including evidence and perspectives that challenge the mainstream consensus. We agreed to undertake the work on the condition that there would be no editorial oversight by the Secretary, the Department of Energy, or any other government personnel. This condition has been honored throughout the process and the writing team has worked with full independence.

The group began working in early April with a May 28 deadline to deliver a draft for internal DoE review. The short timeline and the technical nature of the material meant that we could not comprehensively review all topics. Rather, we chose to focus on topics that are treated by a serious, established academic literature; that are relevant to our charge; that are downplayed in, or absent from, recent assessment reports; and that are within our competence.

While the report is intended to be accessible to non-experts, we have omitted some introductory or explanatory material that can easily be accessed elsewhere. Nor have we attempted to survey the entire literature related to the topics covered. We have focused as much as possible on literature published since 2020 and referenced previous IPCC and NCA assessment reports. We have also used data through 2024 where possible.

The writing team is grateful to Secretary Wright for the opportunity to prepare this report and for his support of independent scientific assessment and open scientific debate. We are also grateful to a team of anonymous peer reviewers whose input helped improve the final report.

John Christy, Ph.D.

Judith Curry, Ph.D.

Steven Koonin, Ph.D.

Ross McKittrick, Ph.D.

Roy Spencer, Ph.D.

PART I: DIRECT HUMAN INFLUENCE ON ECOSYSTEMS AND THE CLIMATE

1 CARBON DIOXIDE AS A POLLUTANT

Chapter summary:

Carbon dioxide (CO₂) differs in many ways from the so-called Criteria Air Pollutants. It does not affect local air quality and has no human toxicological implications at ambient levels. It is an issue of concern because of its effects on the global climate.

The Clean Air Act of 1970 defined six so-called Criteria Air Contaminants subject to regulation (EPA): particulate matter, ground-level ozone, sulfur dioxide, nitrogen dioxide, lead, and carbon monoxide. In 2007, the Supreme Court ruled that greenhouse gases (CO₂ among them) were also “pollutants” subject to regulation under Clean Air Act (*Mass. v. EPA*, 2007). While the definition of “pollutant” is ultimately a legal matter, there are important scientific distinctions between CO₂ and the Criteria Air Contaminants. The latter are subject to regulatory control because they cause local problems depending on concentrations that include nuisances (odor, visibility), damage to plants, and, at high enough exposure levels, toxicological effects in humans. In contrast, CO₂ is odorless, does not affect visibility and has no toxicological effects at ambient levels. It is a naturally occurring part of the atmosphere and a key component of human and plant respiration. CO₂ is essential for plant photosynthesis and higher levels are beneficial for vegetation. In these aspects, CO₂ is similar to water vapor.

Ambient outdoor air today contains about 430 parts per million (ppm) CO₂, increasing at about 2 ppm per year. The U.S. Occupational Safety and Health Administration issues guidelines for indoor workplaces in which elevated CO₂ might be encountered, such as where dry ice is used. The Permissible Exposure Limit is 5,000 ppm over 8 hours (OSHA, 2024). Allen *et al.* (2015) reported evidence of diminished performance on some cognitive tasks among workers in office cubicles when exposed to CO₂ levels above 1,000-1,500 ppm. These levels are far larger than any plausible ambient outdoor value through the end of the 22nd century.

The growing amount of CO₂ in the atmosphere directly influences the earth system by promoting plant growth (global greening), thereby enhancing agricultural yields, and by neutralizing ocean alkalinity. But the primary concern about CO₂ is its role as a greenhouse gas (GHG) that alters the earth’s energy balance, warming the planet. How the climate will respond to that influence is a complex question that will occupy much of this report.

References

- Allen, J., Macnaughton, P., Satish, U., *et al.* (2015). Associations of cognitive function scores with carbon dioxide, ventilation, and volatile organic compound exposures in office workers: A controlled exposure study of green and conventional office environments. *Environmental Health Perspectives*. 124. <https://doi.org/10.1289/ehp.1510037>
- Massachusetts v. Environmental Protection Agency, 549 U.S. 497 (2007). <https://www.oyez.org/cases/2006/05-1120>
- U.S. Environmental Protection Agency. (n.d.). Criteria air pollutants. <https://www.epa.gov/criteria-air-pollutants>
- U.S. Occupational Safety and Health Administration. (2024). OSHA occupational chemical database: Carbon dioxide. <https://www.osha.gov/chemicaldata/183>

2 DIRECT IMPACTS OF CO₂ ON THE ENVIRONMENT

Chapter summary:

CO₂ enhances photosynthesis and improves plant water use efficiency, thereby promoting plant growth. Global greening due in part to increased CO₂ levels in the atmosphere is well-established on all continents.

CO₂ absorption in sea water makes the oceans less alkaline. The recent decline in pH is within the range of natural variability on millennial time scales. Most ocean life evolved when the oceans were mildly acidic. Decreasing pH might adversely affect corals, although the Australian Great Barrier Reef has shown considerable growth in recent years.

2.1 CO₂ as a contributor to global greening

The growing CO₂ concentration in the atmosphere has the important positive effect of promoting plant growth by enhancing photosynthesis and improving water use efficiency. That is evident in the “global greening” phenomenon discussed below, as well as in the improving agricultural yields discussed in Chapter 10. Here we focus just on CO₂ fertilization; research on combined effects due to temperature and precipitation changes are discussed in Chapter 10.

2.1.1 Measurement of global greening

“Greening” refers to an increase in the fraction of the Earth’s surface covered by plants. It can be quantified by the “Leaf Area Index” (LAI) measured by satellite. Many studies over the past decade have confirmed a global greening pattern (increase in LAI) attributable in part to rising CO₂ levels. Zhu *et al.* (2016) was one of the first studies to report that global greening was detectable using satellite sensors. From 1982 to 2011 they detected greening over 25-50 percent of the Earth versus “browning” over only four percent and attributed 70 percent of the greening to rising CO₂ levels (see Figure 2.1). Other contributors included land-use changes, warming and nitrogen. The fraction attributable to CO₂ was largest in the tropics; other factors played more dominant roles in CONUS.

Zeng *et al.* (2017) confirmed the pattern of greening, noting that over thirty years it had added 8 percent to global leaf area and that greening was mitigating warming. Greening has been observed globally. Chen *et al.* (2019) show that in China and India much of it is driven by land management changes. Thus, while China accounts for only 6.6 percent of global vegetated area it accounts for 25 percent of global net increase in LAI. Piao *et al.* (2020) noted that greening was even observable in the Arctic. CO₂ fertilization effects are influenced by local temperature and nutrient and water availability, all of which vary regionally.

While plant models predict increased photosynthesis in response to rising CO₂, Haverd *et al.* (2020) reported a CO₂ fertilization rate much larger than model predictions. That is, CO₂ fertilization had driven an increase in observed global photosynthesis by 30 percent since 1900, versus 17 percent predicted by plant models. If true it would indicate that global models of the socioeconomic impacts of rising CO₂ have understated the benefits to crops and agriculture. Keenan *et al.* (2023), however, estimated a lower fertilization rate more in line with models. The connection between CO₂ fertilization and agriculture will be discussed in Chapter 9.




Figure 2.1: Trends in average Leaf Area Index (LAI). Source: Zhu *et al.* 2016 Figure 3.

Piao *et al.* (2020) and Chen *et al.* (2024) report that the greening trend continues with no evidence of slowdown, and CO₂ fertilization remains the dominant driver

2.1.2 Photosynthesis and CO₂ levels

Plants build biomass through *photosynthesis*, a process that converts carbon dioxide, water, and light into sugar. The plant enzyme responsible for photosynthesis is Ribulose-1,5-bisphosphate-carboxylase/oxygenase or “Rubisco”. Photosynthesis is initiated when CO₂ is available at the surface of the Rubisco enzyme where it is converted to a molecule with 3 carbon atoms and thereafter incorporated into plant mass. This is referred to as the “C3” process.

Rubisco is estimated to have evolved about 3 billion years ago. Over geological time the Earth’s atmospheric CO₂ levels were usually many times higher than they are today. About 400 million years ago CO₂ levels were an estimated 2,000-4,000 ppm and were at or above 1,000 ppm for much of the interval from 200 to 50 million years ago (Berner 2006, Judd *et al.* 2024). Over the past 35 million years the level of atmospheric CO₂ has been steadily declining, falling to as low as 170 ppm during glaciations (Gerhart and Ward 2010). While the modern rate of change in CO₂ may be high compared to prior intervals, the geological evidence is that plants and animals evolved under much higher CO₂ levels than at present.

In response to low-CO₂ conditions some plants evolved another photosynthetic pathway called C4, in which CO₂ is concentrated in the vicinity of Rubisco, allowing for the C3 process to function more efficiently. For agricultural purposes the plant categories are:

- C3: rice, wheat, soybeans and most other crops
- C4: maize (corn), sugar cane, millet, sorghum

Had atmospheric CO₂ levels continued declining, plant growth would have declined and eventually ceased. Below 180 ppm, the growth rates of many C3 species are reduced 40-60 percent relative to 350 ppm (Gerhart and Ward 2010) and growth has stopped altogether under experimental conditions of 60—140 ppm CO₂. Some C4 plants are still able to grow at levels even as low as 10 ppm, albeit very slowly (Gerhart and Ward 2010).




Figure 2.2: growth of *Abutilon theophrasti* after 14 days under identical conditions but for the indicated variations in CO₂ levels. Source: Gerhart and Ward (2010). Note “Current” corresponds to 1988 in image.

Current CO₂ levels are about 430 ppm, up from 280 ppm in the early 1800s. The positive response of plants to extra CO₂ is illustrated in Figure 2.2, reproduced from Gerhart and Ward (2010). It shows the growth effect of CO₂ on Velvetleaf (*Abutilon theophrasti*) seedlings over 14 days under controlled conditions where only the CO₂ exposure is varied. The gains induced by increasing CO₂ from 150 ppm to 350 ppm continue under a further doubling to 700 ppm.

Over the past 60+ years there have been thousands of studies on the response of plants to rising CO₂ levels. The overwhelming theme is that plants, especially C3 plants, benefit from extra CO₂. There are two mechanisms by which CO₂ confers a growth benefit:

- Enhanced photosynthesis via the metabolic pathways described above.
- Increased water use efficiency. This arises because plants draw in CO₂ by opening the stomata (pores) on the leaf surface. When CO₂ is scarce the stomata must be kept wide open for long periods, allowing water to evaporate. Under enriched CO₂ conditions the stomata remain closed for longer periods, thus helping the plant retain water longer, and so increasing water use efficiency.

Specific effects of climate change on U.S. agriculture will be reviewed in Chapter 9.

2.1.3 *Rising CO₂ and crop water use efficiency*

Deryng *et al.* (2016) surveyed evidence on crop water productivity (CWP), the yield per unit of water used, drawing attention to the potential for CO₂ both to enhance photosynthesis and to reduce leaf-level transpiration (water loss during leaf respiration). They surveyed all available FACE data (Free Air CO₂ Enrichment—see Chapter 9) on crop yield changes for maize (corn), wheat, rice, and soybean and combined it with crop model data simulating yield responses as of 2080 under the extreme RCP8.5 emissions scenario in four growing regions (Tropics, Arid, Temperate and Cold) each of which were split into rainfed and irrigated sub-regions. They reported that models without CO₂ fertilization predicted CWP losses in every region, but those were more than offset by CO₂ fertilization so that all regions showed a net CWP gain. Deryng *et al.* (2016) also reported that negative impacts of warming on wheat and soybean yields were fully offset by CWP gains and mitigated by up to 90 percent for rice and 60 percent for maize.

Similarly, Cheng *et al.* (2017) noted that increased Gross Primary Production from 1982 to 2011 due to rising CO₂ uptake was accompanied by such large gains in CWP that global water use by plants had not increased, despite the extra biomass.

Deryng *et al.* (2016) assumed that climate change would “exacerbate water scarcity”. Yet while models do predict that drylands will expand under climate warming, current data show the opposite: greening is happening even in arid areas. Zhang *et al.* (2024) report that due to increased CO₂ levels “increasing aridity in drylands won’t lead to a general loss of vegetation productivity”; at most only 4 percent of currently arid areas will see increased desertification.

2.1.4 *CO₂ fertilization benefits in IPCC Reports*

The IPCC has only minimally discussed global greening and CO₂ fertilization of agricultural crops. The topic is briefly acknowledged in a few places in the body of the IPCC 6th and earlier Assessment Reports but is omitted in all Summary documents. Section 2.3.4.3.3 of the AR6 Working Group I report, entitled “global greening and browning,” points out that the IPCC Special Report on Climate Change and Land had concluded with *high confidence* that greening had increased globally over the past 2-3 decades. It then discusses that there are variations in the greening trend among data sets, concluding that while they have *high confidence* greening has occurred, they have *low confidence* in the magnitude of the trend. There are also brief mentions of CO₂ fertilization effects and improvements in water use efficiency in a few other chapters in the AR6 Working Groups I and II Reports.

Overall, however, the Policymaker Summaries, Technical Summaries, and Synthesis Reports of AR5 and AR6 do not discuss the topic.

2.2 The Alkaline Oceans

2.2.1 *Changing pH*

A neutral aqueous solution has a pH of 7.0, while one with pH greater than 7.0 is alkaline (also termed basic) and with pH less than 7.0 is acidic. The modern-day global average pH of surface sea water is estimated to be 8.04 (Copernicus Marine Service 2025, Figure 2.3), down from an estimated value of 8.2 in pre-Industrial times (Gattuso and Hansson, 2011). As CO₂ concentrations in the atmosphere increased, the oceans absorbed more, which decreases their pH. Depending upon the oceans’ buffering capacity, they are expected to become somewhat less alkaline over time, consistent with the observed decrease in pH.




Figure 2.3: Ocean pH 1985 – 2022. Source: Copernicus Marine Service 2025

While this process is often called “ocean acidification”, that is a misnomer because the oceans are not expected to become acidic; “ocean neutralization” would be more accurate. Even if the water were to turn acidic, it is believed that life in the oceans evolved when the oceans were mildly acidic with pH 6.5 to 7.0 (Krissansen-Totton *et al.*, 2018). On the time scale of thousands of years, boron isotope proxy measurements show that ocean pH was around 7.4 or 7.5 during the last glaciation (up to about 20,000 years ago) increasing to present-day values as the world warmed during deglaciation (Rae *et al.*, 2018). Thus, ocean biota appear to be resilient to natural long-term changes in ocean pH since marine organisms were exposed to wide ranges in pH.

2.2.2 Coral reef changes

There are concerns that a decreasing pH of sea water will reduce the calcification rate of coral reefs. But coral reefs already endure large swings in pH, partly due to daily photosynthetic activity in the reef; measured pH values range from 9.4 during the day to 7.5 at night (Revelle and Fairbridge, 1957). De'ath *et al.* (2009) reported that a portion of Australia's Great Barrier Reef (GBR, the world's largest coral reef ecosystem) had experienced a 14 percent decline in calcification since 1990. This was tentatively attributed to increasing water temperature and decreasing pH. But Ridd *et al.* (2013) showed that report to have resulted from a biased data analysis that, when corrected, showed no change in calcification rates. Nevertheless, the alarm produced by the original paper has persisted as evidenced by the large number of published citations (541) to the original study compared to only 11 citations to the correction (as of 30 April 2025).

The most recent annual summary of GBR conditions from the Australian Institute of Marine Science indicates that coral production has rebounded strongly (AIMS, 2023). Figure 2.4 shows the results of the AIMS surveys of hard coral cover, expressed as a percentage of the reef area. Much of the decline in the GBR before 2011 turned out to be due to intense tropical cyclone activity (Beeden *et al.*, 2015) as well as a string of marine heatwaves, agricultural runoff and invasive species (Woods Hole, 2023). Given the reported declines in GBR calcification between 1990 and 2009 and the continued increase in atmospheric CO₂ levels, the rebound has surprised some observers.




Figure 2.4 Hard coral cover of three regions of the Great Barrier Reef 1985 to 2023. Source: AIMS 2023.

It is being increasingly recognized that publication bias (alarming ocean acidification results preferred by high-impact research publications) exaggerates the reported impacts of declining ocean pH. An ICES Journal of Marine Science Special Issue addressed this problem with an article entitled, *Towards a Broader Perspective on Ocean Acidification Research*. In the Introduction to that Special Issue, H. I. Browman stated, “As is true across all of science, studies that report no effect of ocean acidification are typically more difficult to publish.” (Browman, 2016).

Similarly, a meta-analysis (Clements *et al.*, 2021) of the negative effects of ocean acidification on reef fish behavior found what they called a “decline effect”: initially dramatic conclusions published in prominent journals showing apparently large impacts of acidification tended to be followed up by subsequent studies on larger sample sizes yielding much smaller and typically non-existent effects. They call for their colleagues to improve research practices to counter the “decline effect”:

[The] vast majority of studies with large effect sizes in this field tend to be characterized by low sample sizes, yet are published in high-impact journals and have a disproportionate influence on the field in terms of citations. We contend that ocean acidification has a negligible direct impact on fish behavior, and we advocate for improved approaches to minimize the potential for a decline effect in future avenues of research (Clements *et al.*, 2021).

In summary, ocean life is complex and much of it evolved when the oceans were acidic relative to the present. The ancestors of modern coral first appeared about 245 million years ago. CO₂ levels for more than 200 million years afterward were many times higher than they are today. Much of the public discussion of the effects of ocean “acidification” on marine biota has been one-sided and exaggerated.

References

- AR6: *Intergovernmental Panel on Climate Change Sixth Assessment Report* (2021) Working Group I Contribution. www.ipcc.ch.
- Australian Institute of Marine Science. (2022). Continued coral recovery leads to 36-year highs across two-thirds of the Great Barrier Reef. https://www.aims.gov.au/sites/default/files/2022-08/AIMS_LTMP_Report_on%20GBR_coral_status_2021_2022_040822F3.pdf
- Beeden, R., Maynard, J., Puotinen, M., Marshall, P., Dryden, J., Goldberg, J., and Williams, G. (2015). Impacts and recovery from Severe Tropical Cyclone Yasi on the Great Barrier Reef. *PLOS ONE*, 10, e0121272. <https://doi.org/10.1371/journal.pone.0121272>
- Berner, R. A. (2006). GEOCARBSULF: A combined model for Phanerozoic atmospheric O₂ and CO₂. *Geochimica et Cosmochimica Acta*, 70, 5653–5664.

- Browman, H. I. (2016). Applying organized scepticism to ocean acidification research. *ICES Journal of Marine Science*, 73(3), 529.1–536. <https://doi.org/10.1093/icesjms/fsw010>
- Chen, C., Park, T., Wang, X., Piao, S., Xu, B., Chaturvedi, R. K., and Myneni, R. B. (2019). China and India lead in greening of the world through land-use management. *Nature Sustainability*, 2, 122–129. <https://www.nature.com/articles/s41893-019-0220-7>
- Chen, X., Wang, Y., Liu, Y., and Piao, S. (2024). The global greening continues despite increased drought stress since 2000. *Global Ecology and Conservation*, 49, e02791. <https://www.sciencedirect.com/science/article/pii/S2351989423004262>
- Cheng, L., Zhang, L., Wang, Y. P., *et al.* (2017). Recent increases in terrestrial carbon uptake at little cost to the water cycle. *Nature Communications*, 8, 110. <https://doi.org/10.1038/s41467-017-00114-5>
- Clements, J. C., Sundin, J., Clark, T. D., and Jutfelt, F. (2022). Meta-analysis reveals an extreme “decline effect” in the impacts of ocean acidification on fish behavior. *PLOS Biology*, 20(2), e3001511. <https://doi.org/10.1371/journal.pbio.3001511>
- Copernicus Marine Service. (2025). Global ocean acidification – Mean sea water pH time series and trend from multi-observations reprocessing. https://data.marine.copernicus.eu/product/GLOBAL_OMI_HEALTH_carbon_ph_area_averaged/description
- De'ath, G., Lough, J., and Fabricius, K. (2009). Declining coral calcification on the Great Barrier Reef. *Science*, 323, 116–119. <https://doi.org/10.1126/science.1165283>
- Deryng, D., Conway, D., Ramankutty, N., Price, J., Warren, R., Jones, R., ... and Elliott, J. (2016). Regional disparities in the beneficial effects of rising CO₂ concentrations on crop water productivity. *Nature Climate Change*. <https://doi.org/10.1038/nclimate2995>
- Gattuso, J. P., and Hansson, L. (Eds.). (2011). *Ocean acidification: Background and history*. Oxford University Press.
- Gerhart, L. M., and Ward, J. K. (2010). Plant responses to low [CO₂] of the past. *New Phytologist*, 188, 674–695. <https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-8137.2010.03441.x>
- Haverd, V., B. Smith, J. G. Canadell, *et al.* (2020). Higher than expected CO₂ fertilization inferred from leaf to global observations. *Global Change Biology*, 26, 2390–2402. <https://doi.org/10.1111/gcb.14950>
- Keenan, T. F., X. Luo, B. D. Stocker, *et al.* (2023). A constraint on historic growth in global photosynthesis due to rising CO₂. *Nature Climate Change* 13(12): 1376-1381 DOI: 10.1038/s41558-023-01867-2.
- Judd, E. J., Scotese, C. R., Young, S. A., *et al.* (2024). A 485-million-year history of Earth's surface temperature. *Science*, 385(6715). <https://doi.org/10.1126/science.adk3705>
- Krissansen-Totton, J., Arney, G. N., and Catling, D. C. (2018). Constraining the climate and ocean pH of the early Earth with a geological carbon cycle model. *Proceedings of the National Academy of Sciences*, 115(6), 4105–4110. <https://doi.org/10.1073/pnas.1721296115>
- Piao, S., X. Wang, T. Park, *et al.* (2020). Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth & Environment* 1(1): 14-27 DOI: 10.1038/s43017-019-0001-x
- Rae, J. W. B., Burke, A., Robinson, L. F., *et al.* (2018). CO₂ storage and release in the deep Southern Ocean on millennial to centennial timescales. *Nature*, 562, 569–573. <https://doi.org/10.1038/s41586-018-0614-0>
- Revelle, R., and Fairbridge, R. W. (1957). Carbonate and carbon dioxide. In J. W. Hedgpeth (Ed.), *Treatise on marine ecology and paleoecology* (Vol. 1). *Geological Society of America*.
- Ridd, P., Silva, E., and Stieglitz, T. (2013). Have coral calcification rates slowed in the last twenty years? *Marine Geology*, 346, 392–399. <https://doi.org/10.1016/j.margeo.2013.09.002>
- Woods Hole Oceanographic Institution. (2023). Is the Great Barrier Reef making a comeback? <https://www.whoi.edu/oceanus/feature/is-the-great-barrier-reef-making-a-comeback/>
- Zeng, Z., Piao, S. Li, L., *et al.* (2017). Climate mitigation from vegetation biophysical feedbacks during the past three decades. *Nature Climate Change*. <https://doi.org/10.1038/nclimate3299>

- Zhang, Y., Liu, Y., Chen, X., *et al.* (2024). Less than 4% of dryland areas are projected to desertify despite increased aridity under climate change. *Nature Communications Earth and Environment*, 5. <https://www.nature.com/articles/s43247-024-01463-y>
- Zhu, Z., Piao, S., Myneni, R. B., *et al.* (2016). Greening of the Earth and its drivers. *Nature Climate Change*, 6, 791–795. <https://www.nature.com/articles/nclimate3004>

3 HUMAN INFLUENCES ON THE CLIMATE

Chapter Summary:

The global climate is naturally variable on all time scales. Anthropogenic CO₂ emissions add to that variability by changing the total radiative energy balance in the atmosphere.

The IPCC has downplayed the role of the sun in climate change but there are plausible solar irradiance reconstructions that imply it contributed to recent warming.

Climate projections are based on IPCC emission scenarios that have tended to exceed observed trends. Most academic climate impact studies in recent years are based upon the extreme RCP 8.5 scenario that is now considered implausible; its use as a business-as-usual scenario has been misleading.

Carbon cycle models connect annual emissions to growth in the atmospheric CO₂ stock. While models disagree over the rate of land and ocean CO₂ uptake, all agree that it has been increasing since 1959.

There is evidence that urbanization biases in the land warming record have not been completely removed from climate data sets.

3.1 Components of radiative forcing and their history

3.1.1 Historical radiative forcing

A changing climate has been the norm throughout the Earth's 4.6-billion-year history. The Earth's temperature and weather patterns change naturally over timescales ranging from decades to millions of years. Natural variations in the surface climate originate in two ways. Internal climate fluctuations associated with circulations in the atmosphere and ocean exchange energy, water, and carbon between the atmosphere, oceans, land, and ice. External influences on the climate system include variations in the energy received from the sun and the effects of volcanic eruptions. Human activities influence climate through changing land use and land cover. Humans are also changing the composition of the atmosphere by emissions of CO₂ and other greenhouse gases and by altering the concentration of aerosol particles in the atmosphere.

The earth is warmed by the sunlight it absorbs and is cooled by the heat it radiates to space. Averaged over the Earth's surface, each of these processes involve power flows of about 240 Watts per square meter (W/m²). When they are in balance, there are no net external causes of warming or cooling. Both human and natural influences on the climate alter this balance and so cause the climate to change.

Influences on the Earth's energy balance at the top of the atmosphere are quantified by "radiative forcing", the extent to which they disturb the warming/cooling balance; a positive forcing warms while a negative forcing cools. The IPCC's estimated history of major components of radiative forcing since 1750 is shown in the following two figures from its AR6.




Figure 3.1.1: IPCC estimates of radiative forcing components over time. Shading indicates uncertainty ranges. Source: AR6 WGI Ch2 Fig. 10




Figure 3.1.2 IPCC estimates of radiative forcing component changes from 1750 to 2019. Source: AR6 WGI Ch 7 Fig. 7-6.

These graphs show that the total radiative forcing is comprised of both natural and anthropogenic components. Carbon dioxide is the largest human influence on the climate and the one most relevant to the influence of fossil fuel use. It exerts a warming influence by decreasing the cooling power of the atmosphere. Emissions of CO_2 are accumulating in the atmosphere, as described in the following section, so that the warming influence is growing. Other greenhouse gases (methane, nitrous oxide, halogens, and ozone) act similarly, currently adding another 75 percent to CO_2 's warming. Aerosols exert an overall cooling effect, although with large uncertainties in the way they catalyze the formation of reflective clouds. As a result, understanding the causes of recent warming requires not just identifying the warming effects of CO_2 , but also the more uncertain cooling effects of aerosols.

The IPCC assesses the change in the radiative forcing by the sun to be negligible, based on their preference for data reconstructions that imply minimal solar change since preindustrial times. But Connolly *et al.* (2021) reviewed sixteen different Total Solar Irradiance (TSI) reconstructions in the literature

covering the years 1600-2000; the reconstructions vary from almost no change in TSI to a relatively large upward trend. Those authors note that the variation in TSI reconstructions combined with variations in surface temperature reconstructions allows for inferences consistent with either no or most 20th century warming being attributable to the sun.

A particularly thorny issue is the gap in TSI data between 1989 and 1991 due to a delay in the launch of a monitor following the Space Shuttle Challenger disaster on January 28 1986. This delay prevented a replacement satellite from being launched in time to overlap with, and its readings to be intercalibrated with, the prior system (Zacharias 2014, Scafetta *et al.* 2019). This is called the ACRIM (Active Cavity Radiometer Irradiance Monitor) gap problem. The question of whether there is an upward trend in TSI over 1978 to 2018 hangs on how the ACRIM data gap is filled. Connolly *et al.* (2021) found that the IPCC's consensus statements on solar forcing were formulated prematurely through the suppression of dissenting scientific opinions.

Another natural radiative forcing component is volcanic aerosols, which exert an episodic cooling influence. Box 4.1 in the IPCC AR6 Report addresses the climate impact of volcanic eruptions, noting three explosive volcanic eruptions that occurred in the first half of the 19th century. This included the 1815 Tambora eruption that resulted in the 'year without summer', with multiple harvest failures across the Northern Hemisphere. There is uncertainty about the sign of the relatively small forcing due to the submarine volcano Hunga Tonga which erupted in 2022 (Jenkins *et al.* 2023, Schoeberl *et al.* 2024).

Figure 3.1.1 shows that the anthropogenic forcing component was negligible before about 1900 and has increased steadily since, rising to almost 3 W/m² today. However, this is still only about 1 percent of the unperturbed radiation flows, making it a challenge to isolate the effects of anthropogenic forcing; state-of-the-art satellite estimates of global radiative energy flows are only accurate to a few W/m².

Natural sources of global energy imbalance other than volcanoes and total solar irradiance (TSI) are not included in these graphs because they remain largely unknown.

3.1.2 Change in atmospheric CO₂ since 1958

Carbon dioxide's warming influence depends on how much "extra" CO₂ accumulates in the atmosphere- *i.e.*, its concentration above the preindustrial value of 280 ppm. The CO₂ level as recorded at the Mauna Loa observatory in Hawaii, generally used as the representative global average concentration, is available online at <https://gml.noaa.gov/ccgg/trends/index.html>. The concentration was about 316 ppm at the start of the record in 1959 and is now about 430 ppm, a 36 percent increase. At the end of the last glaciation CO₂ levels had fallen to about 180 ppm. As discussed in Chapter 2, C3 plants begin dying at CO₂ levels below about 140 ppm and C4 plants at levels below 100 ppm, so if CO₂ levels had continued falling plant life would have been imperiled.




Fig. 3.1.3. Yearly average atmospheric CO₂ concentrations (1959-2025) in ppm measured at Mauna Loa (blue). C3 Threshold: Level below which C3 plants begin dying (140 ppm, see Chapter 2). C4 Threshold: Level below which C4 plants begin dying (100 ppm, see Chapter 2). Glacial minimum: Minimum level during recent glaciations (purple arrow). CO₂ data source: <https://gml.noaa.gov/ccgg/trends/index.html>

The annual increase in concentration is only about half of the CO₂ emitted because land and ocean processes currently absorb “excess” CO₂ at a rate approximately 50 percent of the human emissions. Future concentrations, and hence future human influences on the climate, therefore depend upon two components: (1) future rates of global human CO₂ emissions, and (2) how fast the land and ocean remove extra CO₂ from the atmosphere. We discuss each of these in turn.

3.2 Future emission scenarios and the carbon cycle

3.2.1 Emission scenarios

Assessing the dangers of future GHG emissions requires assumptions about what those emissions will be. Future emissions, and hence human influences on the climate, will depend upon future demographics, economic activity, regulation, and energy and agricultural technologies. Various assumptions about each of those lead to projections of greenhouse gas emissions and concentrations, aerosol concentrations, and changes in land use, which ultimately can be combined into assumptions about anthropogenic radiative forcing.

The great uncertainties about these many factors make it impossible to precisely predict future emissions. Instead, the IPCC has used various sets of scenarios meant to span a plausible range of possibilities for population, economy, and technologies. Recent versions of the scenarios are labeled by a number indicating the anthropogenic radiative forcing expected in 2100 under that scenario. Thus, a scenario labeled with a “6” corresponds to 6 W/m² of human-induced radiative forcing (warming) at the end of the century. (Recall current anthropogenic radiative forcing is about 2.7 W/m².)